

279-3155

03-14-2011

1/10



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND  
POLLUTION PREVENTION

March 14, 2011

Christopher Davis  
FMC Corporation  
1735 Market Street  
Philadelphia, PA 19103

Subject: Notification -- Per PR Notices 98-10, PRN 97-6 and Agency's June 4,  
2009 letter  
EPA Registration Number: 279-3155  
Date of Submission: November 5, 2010

Dear Mr. Davis:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration (PRN) 98-10 dated November 5, 2010, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact BeWanda Alexander at (703) 305-7460 or e-mail [alexander.bewanda@epa.gov](mailto:alexander.bewanda@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "BeWanda Alexander for".

Richard Gebken  
Product Manager 10  
Insecticide Branch  
Registration Division (7505P)

Enclosures: Copy of Label Stamped "Notification"

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

**Application for Pesticide - Section I**

1. Company/Product Number 279-3155	2. EPA Product Manager Gebken	3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted
4. Company/Product (Name) Talstar Nursery, Flowable Insecticide/Miticide	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) FMC Corporation 1735 Market Street Philadelphia, PA 19103 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____  Product Name _____

**Section - II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

**NOTIFICATION**

**MAR 14 2011**

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification per PR Notices 98-10, 97-6, and EPA letter of June 4, 2009  
 On behalf of FMC Corporation, I certify that this notification is consistent with the provisions of PR Notice 98-10, PR-Notice 97-6 and the EPA letter of June 4, 2009. and EPA regulations at 40 CFR 152.46. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10, 97-6, the EPA letter of June 4, 2009 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

**Section - III**

**1. Material This Product Will Be Packaged In:**

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted If "Yes" Unit Packaging wgt.      No. per container		If "Yes" Package wgt      No. per container	

3. Location of Net Contents Information  
 Label       Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product  
 Lithograph Paper glued Stenciled       Other \_\_\_\_\_


**Section - IV**

**1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)**

Name Christopher Davis	Title Registration Manager	Telephone No: (Include Area Code) 215-299-6334
---------------------------	-------------------------------	---

**Certification**

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Registration Manager	6. Date Application Received (Stamped) _____ _____ _____ _____ _____
4. Typed Name Christopher Davis	5. Date 11-5-10	

# Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
279-3155	11/5/10	000279-03155.20101105.Notification.pdf

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

  
 \_\_\_\_\_  
 Signature

11-5-10  
 \_\_\_\_\_  
 Date

Christopher Davis  
 \_\_\_\_\_  
 Name (typed)

Registration Manager  
 \_\_\_\_\_  
 Title

# FMC Agricultural Products

FMC Corporation  
1735 Market Street  
Philadelphia, PA 19103  
215.299.6000 Phone  
www.fmc.com

November 5, 2010

Mr. Richard Gebken (PM-13)  
United States Environmental Protection Agency  
Office of Pesticide Programs (NOTIF)  
Registration Division (H7504P)  
Document Processing Desk  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202

Dear Mr. Gebken:

**Subject: Talstar Nursery Flowable Insecticide/Miticide  
EPA Reg. No. 279-3155  
Notification per PR Notice 98-10 and Letter of June 4, 2009**

FMC is interested in revising labeling for the subject product. FMC believes that the changes are in full compliance with PR Notice 98-10, Section II Labeling Notifications; part N Other Revisions; PR-Notice 97-6, and the EPA letter of June 4, 2009 for Non-Ag Outdoor Use Pyrethroids.

Specifically, we have made the following changes:


- Changed "Inert Ingredients" to "Other Ingredients"
- Amended the warranty statement to be consistent with the changes you have requested for all of our labels
- Revised the environmental hazards language and directions for use for the subject product per EPA's letter dated June 4, 2009.

To complete this notification, enclosed please find the following:

- Application for Pesticide (EPA Form 8570-1).
- Final Print Labeling (2 copies), including one hi-lited version for ease of review.
- Completed Certification with Respect to Label Integrity form
- A compact disk containing the label as amended
- Certification of Compliance with PR Notice 98-10, 97-6 and letter of June 4, 2009

Thank you for your attention to this matter, please feel free to contact me at (215) 299-6334 or e-mail at [chris.davis@fmc.com](mailto:chris.davis@fmc.com).

Sincerely,



Christopher Davis, Registration Manager  
APG Specialty Products Business



510

**RESTRICTED USE PESTICIDE**

Toxic to fish and aquatic organisms.  
For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

1658

Net Contents

**Talstar**<sup>®</sup>  
Nursery Flowable  
Insecticide/Miticide

**NOTIFICATION**

**MAR 14 2011**

**For Commercial Non-Food Use on Indoor and Outdoor Ornamentals, Greenhouses, Nurseries, Turf on Golf Courses and Sod Farms.**

EPA Reg. No. 279-3155

EPA Est. 279-

<b>Active Ingredient:</b>	<b>By Wt.</b>
Bifenthrin:*	7.9%
<b>Other Ingredients:</b>	92.1%
	100.0%

\*Cis isomers 97% minimum, trans isomers 3% maximum.  
Talstar® Nursery Flowable insecticide/miticide contains 2/3 pound active ingredient per gallon.  
U.S. Patent No. 4,238,505

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See other panels for additional precautionary information.

**NOTE: USERS OF THIS PRODUCT IN CALIFORNIA MUST BE IN POSSESSION OF STATE SPECIFIC SUPPLEMENTAL LABELING**

**DO NOT USE THIS PRODUCT ON GOLF COURSES AND SOD FARMS IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK.**



FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia PA 19103

**FIRST AID**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**HOTLINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.

**NOTE TO PHYSICIAN**

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

**For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362).**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans (and Domestic Animals)**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling.

**Personnel Protective Equipment:**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions or category C on an EPA chemical resistance category selection chart.

**Applicators and other handlers (other than mixers and loaders) must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.
- Shoes plus socks

**Mixers and Loaders must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## User Safety Recommendations

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help to avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow to drift to blooming crops if bees are visiting the treatment area.

## Use Directions for Tip-N-Measure Container

1. Remove the measuring chamber cap and induction seal. Replace the cap and securely tighten. Tip container until liquid fills measuring chamber.
2. Return container to level position. No adjustment is needed.
3. Remove measuring chamber cap and dispense into proper application equipment.

For multiple dose measuring: Remove fill chamber cap and dispense according to markings on side of bottle.

## DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not water treated area to the point of run-off.

Do not make applications during rain.

Do not apply this product through any kind of irrigation system.

## Agriculture Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as, plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton.
- Shoes plus socks

## Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

Do not allow people or pets on treated surfaces until the spray has

## STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use replace lids and close tightly. Do not put concentrate or dilute material into food or drink container.

**In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.**

**To Confine Spill:** If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal:** Pesticide wastes are toxic. Do not contaminate water, food or feed by storage or disposal. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. Dispose of excess or waste pesticide by use according to label directions, or contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### Container Disposal

**Metal or Plastic Container:** Non-refillable container. Do not reuse or refill this container. **Triple rinse as follows:** Empty the contents into application equipment or a mix tank and drain for 10 seconds after flow begins to drip. Fill container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. **Then offer for recycling, if available or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill.**

## General Applications Instructions

Talstar® Nursery Flowable insecticide/miticide formulation mixes readily with water and other aqueous carriers, and controls a wide spectrum of insects and mites on trees, shrubs, foliage plants, non-bearing fruit and nut trees, and flowers in greenhouses and outdoor nurseries, and turf on golf courses and sod farms. Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application.

Talstar Nursery Flowable may be tank-mixed with other products, including insect growth regulators. When tank mixing Talstar Nursery Flowable with other products, observe all precautions and limitations on each separate product label. The addition of spreader stickers is not necessary. The physical compatibility of Talstar Nursery Flowable may vary with different sources of pesticide products, and local cultural practices. Any tank mixture which has not been previously tested should be prepared on a small scale (pint or quart jar), using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture.

The following procedure is recommended for preparation of a new tank mix, unless specified otherwise in label directions: (1) Add wettable powders to tank water, (2) Agitate, (3) Add liquids and flowables, (4) Agitate, (5) Add emulsifiable concentrates, and (6) Agitate. If a mixture is found to be incompatible following this order of addition, try reversing the order of addition, or increase the volume of water. Note: If the tank-mixture is found to be compatible after increasing the amount of water, then the sprayer will need to be recalibrated for a higher volume application. Do not allow tank mix to stand overnight.

Maximum rates: Do not apply more than 0.2 lb ai/acre (40 fl. ozs. of Talstar Nursery Flowable) in a single application or per year for outdoor applications.

**Resistance:** Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

### Talstar Nursery Flowable Dilution Chart

Applic. Volume: Gallons Per Acre	Applic. Rate: Lbs ai per Acre	Fluid Ounces* of Talstar Nursery Flowable Diluted to these Volumes of Finished Spray			
		1 Gallon	25 Gallons	50 Gallons	100 Gallons
50	0.025	0.1	2.5	5.0	10.0
50	0.05	0.2	5.0	10.0	20.0
50	0.1	0.4	10.0	20.0	40.0
50	0.2	0.8	20.0	40.0	80.0
100	0.025	0.05	1.25	2.5	5.0
100	0.05	0.1	2.5	5.0	10.0
100	0.1	0.2	5.0	10.0	20.0
100	0.2	0.4	10.0	20.0	40.0
150	0.025	0.03	0.83	1.67	3.3
150	0.05	0.07	1.67	3.33	6.7
150	0.1	0.133	3.33	6.67	13.3
150	0.2	0.266	6.67	13.33	26.7
200	0.025	0.025	0.63	1.25	2.5
200	0.05	0.05	1.25	2.5	5.0
200	0.1	0.1	2.5	5.0	10.0
200	0.2	0.2	5.0	10.0	20.0
250	0.025	—	0.5	1.0	2.0
250	0.05	—	1.0	2.0	4.0
250	0.1	—	2.0	4.0	8.0
250	0.2	—	4.0	8.0	16.0
300	0.025	—	0.42	0.83	1.7
300	0.05	—	0.83	1.67	3.3
300	0.1	—	1.67	3.33	6.7
300	0.2	—	3.33	6.67	13.3

\*To convert to milliliters, multiply by 29.57  
 1 fl. oz. = 29.57 ml = 2 tablespoons = 6 teaspoons  
 Do not use household utensils to measure Talstar Nursery Flowable.

**Formula for Determining the Active Ingredient Content of the Finished Spray Mixture:** The following formula may be used to determine the percent active ingredient that is in the spray tank after mixing Talstar Nursery Flowable:

$$\frac{(7.9)(\text{Fl. Oz. of Talstar added to tank})}{(\text{Gallons of finished spray mix})(128)} = \text{Percent Active Ingredient of spray mix}$$

### APPLICATION RECOMMENDATIONS

#### Ornamentals in Greenhouses, Lath Houses, Shade Houses and Outdoor Nurseries, including Non-Bearing Fruit and Nut Trees

Apply 0.025 to 0.2 lbs ai/A (5 to 40 fl. ozs.) of Talstar Nursery Flowable insecticide/miticide. Talstar Nursery Flowable may be diluted and applied in various volumes of water providing that the maximum label rate (0.2 lbs ai/A or 40 fl. ozs.) is not exceeded (refer to Dilution Chart for specific instructions). Talstar Nursery Flowable may be applied through low volume application equipment by dilution with water or other carriers and providing that the maximum label rate (0.2 lbs ai/A or 40 fl. ozs.) is not exceeded.

### ORNAMENTAL APPLICATION RATES

The application rates listed in the following table will provide control of the respective pests under typical conditions. However, at the discretion of the applicator, Talstar Nursery Flowable may be applied at up to 0.2 lbs ai/A (40 fl ozs) to control each of the pest listed in this Table.

Pest	Application Rate Talstar Nursery Flowable	
	lbs ai/A	Fluid Ounces per Acre
Aphids Bagworms <sup>1</sup> Cutworms Elm Leaf Beetles Fall Webworms Lace Bugs Leaf Feeding Caterpillars Plant Bugs (Including Lygus spp.) Tent Caterpillars	0.025 - 0.05	5 - 10
Beet Armyworm Black Vine Weevil (Adults) Brown Soft Scales Broad Mites Budworms California Red Scale (Crawlers) <sup>2</sup> Centipedes Citrus Thrips Clover Mites Crickets Diaprepes (Adults) Earwigs European Red Mite Flea Beetles Fungus Gnats (Adults) Grasshoppers Gypsy Moth Caterpillars Leafhoppers Leafrollers Mealybugs Millipedes Mites Orchid Weevil Pillbugs Pine Needle Scales (Crawlers) <sup>2</sup> San Jose Scales (Crawlers) <sup>2</sup> Sowbugs Spider Mites Spiders Thrips Tip Moths Twig Borers <sup>2</sup> Weevils Whiteflies	0.05 - 0.1	10 - 20
Ants Imported Fire Ants** Japanese Beetle (Adult) Leafminers Pecan Leaf Scorch Mite Pine Shoot Beetle (Adults)	0.1 - 0.2	20 - 40

<sup>1</sup>Bagworms: Apply when larvae begin to hatch and spray larvae directly. Applications when larvae are young will be most effective.  
<sup>2</sup>Scale Crawlers and Twig Borers: Treat trunks, stems and twigs in addition to plant foliage.  
 \*\*For foraging ants.

Certain cultivars may be sensitive to the final spray solution. A small number of plants should be treated and observed for one week prior to application to the entire planting.

Apply with ground equipment only.

Do not apply when wind direction favors downwind drift towards nearby water bodies.

Do not apply when wind velocity exceeds 10 mph.

Avoid application when wind gusts approach 10 mph.

Do not apply when a temperature inversion exists.

Apply using the largest nozzle size compatible with adequate coverage.

Do not apply if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).

Do not apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. When treating tall trees (>15 feet) from the ground with high pressure sprays or during any application with air

assisted equipment (mist blower) do not apply within 150 feet of aquatic areas.

## APPLICATION RECOMMENDATIONS

### Turf (Golf Courses and Sod Farms)

#### NOT FOR USE ON SOD FARMS IN THE STATE OF NEW YORK.

Apply Talstar® Nursery Flowable Insecticide/Miticide as a surface or sub-surface treatment. Use application volumes of up to 10 gallons per 1000 square feet to get uniform coverage when treating dense and or long turf foliage.

For low volume applications, less than 2 gallons/1000 square feet, immediate irrigation of treated area with at least 0.25 inches of water following application to ensure efficacy of sub-surface pests such as, but not limited to, Mole Crickets, is recommended.

## TURF (Golf Courses and Sod Farms) APPLICATION RATES

The application rates listed in the following table will provide control of the respective pests under typical conditions. However, at the discretion of the applicator a, Talstar Nursery Flowable may be applied at up to 0.1 lb ai/A (20 fl. ozs.) to control each of the pests listed in this Table. (0.2 lb ai/A or 40 fl. ozs. of Talstar Nursery Flowable for ants, imported fire ants and mole crickets).

aDuring periods of high pest pressure or for maximum residual control.

Pest	Active Ingredient lbs. per acre	Application Rate Talstar Nursery Flowable	
Armyworms <sup>3</sup> Cutworms <sup>3</sup> Sod Webworm <sup>3</sup>	0.05 lbs ai per acre	10 fl. oz per acre	0.25 fl. oz. per 1000 sq.ft.
Annual Bluegrass Weevil (Hyperodes) (Adult) <sup>4</sup> Ants Billbugs (Adult) <sup>5</sup> Black Turfgrass Ataenius (Adult) <sup>6</sup> Centipedes Chinch Bugs <sup>7</sup> Crickets Earwigs Fleas (Adult) Grasshoppers Leafhoppers Mealybugs Millipedes Mites <sup>8</sup> Mole Cricket (Adult) <sup>9</sup> Mole Cricket (Nymph) <sup>10</sup> Pillbugs Sowbugs	0.05 - 0.1 lbs ai per acre	10 - 20 fl. oz per acre	0.25 - 0.5 fl. oz. per 1000 sq.ft.
Fleas (Larvae) <sup>11</sup> Imported Fire Ants Japanese Beetle (Adult) Ticks <sup>12</sup>	0.1 lbs ai per acre	20 fl. oz per acre	0.5 fl. oz. per 1000 sq.ft.
Ants Imported Fire Ants <sup>14</sup> Mole Crickets	0.2 <sup>13</sup> lbs ai per acre	40 <sup>13</sup> fl. oz per acre	1 fl. oz. <sup>13</sup> per 1000 sq.ft.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Talstar Nursery if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

### Comments

<sup>3</sup>Armyworms, Cutworms and Sod Webworms: To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application. If the turf area is being maintained at a mowing height of greater than 1 inch, then higher application rates (Up to 0.1 lb ai/A or 20 fl. ozs. of Talstar Nursery Flowable) may be required during periods of high pest pressure.

<sup>4</sup>Annual Bluegrass Weevil (*Hyperodes*) adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into turf areas. This movement generally begins when *Forsythia* is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

<sup>5</sup>Billbug adults: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

<sup>6</sup>Black Turfgrass Ataenius adults: Applications should be made during May and July to control the first and second generation of black turfgrass ataeenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).

<sup>7</sup>Chinch Bugs: Chinch Bugs infest the base of turf plants and are often found in the thatch layer. Irrigation of the turf area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained. Chinch Bugs can be one of the most difficult pests to control in grasses and the higher application rates (Up to 0.1 lb ai/A or 20 fl. ozs. of Talstar Nursery Flowable) may be required to control populations that contain both nymphs and adults during the middle of the summer.

<sup>8</sup>Mites: To ensure optimal control of eriophyid mites, apply in combination with the labeled application rate of a surfactant. A second application, five to seven days after the first, may be necessary to achieve control.

<sup>9</sup>Mole Cricket adults: Achieving control of adult mole crickets is difficult because preferred turf areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Turf areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

<sup>10</sup>Mole Cricket nymphs: Turf areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

<sup>11</sup>Flea larvae: Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Use a higher volume application when treating these areas to ensure penetration of the insecticide into the soil. Note: if the lawn area is being treated with Talstar Nursery Flowable at 0.05 lb. ai/A (10 fl. ozs.) for adult flea control, then the larval application rate may be achieved by doubling the application volume.

<sup>12</sup>Ticks: Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. A repeat application, seven days after the first, may be necessary to achieve control. Do not allow public use of treated areas during application or until sprays have dried.

Deer ticks (*Ixodes sp.*) have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or turf above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered.

<sup>13</sup>Note: For large infestations of ants, imported fire ants, and mole crickets, a single application of 0.2 lb ai/A (40 fl. ozs. of Talstar Nursery Flowable) may be applied once per year.

<sup>14</sup>Imported Fire Ants: Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application or use a high volume application. Broadcast treatments should apply 0.2 lb ai/A (40 fl. ozs. of Talstar Nursery Flowable). Mounds should be treated by diluting 1 teaspoon of Talstar Nursery Flowable per gallon of water and applying 1 to 2 gallons of finished spray per mound. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. A four foot diameter circle around the mound should also be treated. For best results, apply in cool weather (65 - 80°F) or in early morning or late evening hours. Note: a spray rig that is calibrated to apply 0.2 lb ai/A (40 fl. ozs.) of Talstar Nursery Flowable in 5 gallons per 1,000 square feet contains the approximate dilution (1 teaspoon per gallon) that is required for fire ant mound drenches in the spray tank.

Apply with ground application equipment only (and apply with nozzles not more than two feet above the turf).

Do not apply when wind conditions favor downwind drift to nearby water bodies.

Do not apply when wind velocity exceeds 10 mph.

Avoid application when wind gusts approach 10 mph.

Do not apply when a temperature inversion exists.

Apply using nozzles that provide the largest droplet size compatible with adequate coverage.

Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).

Do not apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not apply when turf areas are water-logged or soil is saturated with water (i.e. will not accept irrigation).

### Imported Fire Ant Quarantine Treatment

Against Imported Fire Ants (IFA) in Potting Media (including balled and containerized nursery grown ornamental trees, shrubs, plants, flowers, conifers, bushes, Christmas trees, and non-bearing fruit and nut-trees). Talstar Nursery Flowable insecticide/miticide is approved and can be used in accordance with the USDA Imported Fire Ant Quarantine Program. Talstar Nursery Flowable may be applied either soil incorporated, as a topical application, or as a high volume drench treatment.

**Soil Incorporation:** Incorporate the appropriate volume of Talstar Nursery Flowable (see table below) per cubic yard of potting media by diluting it in water (typically 1 quart to 1 gallon per cubic yard of media) and sprinkling or spraying it onto the media. The applications are based on the dry bulk density of the potting media. When used in accordance with USDA guidelines, this application will provide a 6 month certification period.

### Recommended Soil Incorporation Rate of Talstar Nursery Flowable for Control of IFA in Potting Media.

Potting Media Bulk Density (lbs. cubic yard)	Fluid ounces of Talstar Nursery Flowable in one cubic yard
200	1.9
400	3.8
600	5.7
800	7.6
1000	9.5
1200	11.4
1400	13.3

Use proportional amounts of Talstar Nursery Flowable for potting media with bulk densities not listed.

**Topical Application:** Mix Talstar Nursery Flowable in 1,000 ounces of water based on container size and bulk density of the potting media (see table below). Apply one (1) ounce of the mix to each container evenly distributed over the surface of the potting media. Irrigate all treated containers with 1.5 inches of water following application. When used in accordance with USDA guidelines, this application will provide a 6 month certification period.

### Recommended Topical Drench Application Rate of Talstar Nursery Flowable for Control of IFA in Potting Media.

Potting Media Bulk Density (lbs. cubic yard)	Fluid ounces of Talstar Nursery Flowable per 1,000 ounces of water	
	3 Qt. Container	4 Qt. Container
200	3.6	5.2
400	7.2	10.4
600	10.8	15.6
800	14.4	20.8
1000	18.0	26.0
1200	21.6	31.2
1400	25.2	36.4

Use proportional amounts of Talstar Nursery Flowable for potting media with bulk densities not listed.

**High Volume Drench:** Apply Talstar Nursery Flowable as a high volume drench by mixing the appropriate amount of product based on the bulk density in 100 gallons of water (see table below). Apply mix to individual containers to the point of saturation. The amount of mix used for each plant is generally 1/5 volume of the container. When used in accordance with USDA guidelines, this application will provide a 6 month certification period.

### Recommended High Drench Application Rate of Talstar Nursery Flowable for Control of IFA in Potting Media.

Potting Media Bulk Density (lbs. cubic yard)	Fluid ounces of Talstar Nursery Flowable in 100 Gallons
200	2.4
400	4.8
600	7.2
800	9.6
1000	12.0
1200	14.4
1400	16.8

Use proportional amounts of Talstar Nursery Flowable for potting media with bulk densities not listed.

### Larval Control in Potting Media of Containerized Plants.

**Black Vine Weevil Larval Control - Preventative Treatment - Topical Drench:** For preventative control of black vine weevil larvae in containerized plants, dilute Talstar Nursery Flowable at the rate of 10 to 40 fl. ozs. (0.05 to 0.2 lb AI) per 100 gallons and apply as a drench at the rate of 4 to 8 fluid ounces of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container. Diluting 10 fluid ounces of Talstar Nursery Flowable per 100 gallons and applying 8 fluid ounces of finished spray per 6 inch (diameter) container will provide black vine weevil larval control for one growing season when the application is made in the spring. Diluting 20 to 40 fluid ounces of Talstar Nursery Flowable per 100 gallons and applying 8 fluid ounces of finished spray per 6 inch (diameter) container will provide black vine weevil larval control for two growing seasons when the application is made in the spring.

**White Grub Control - Preventative Treatment - Topical Drench:** For preventative control of white grubs (including, but not limited to, Japanese beetle, oriental beetle and European chafer) in containerized plants, dilute Talstar Nursery Flowable at the rate of 40 to 80 fluid ounces (0.2 to 0.4 lb AI) per 100 gallons and apply as a drench at the rate of 4 to 8 fluid ounces of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container. Use the higher application rate for a longer period of control.

**Black Vine Weevil and White Grub Larval Control - Preventative Treatment - Media Incorporation:** For preventative control of black vine weevil and white grub larvae in containerized plants, incorporate the appropriate volume of Talstar Nursery Flowable (see table below) per cubic yard of potting media by diluting it in water (typically 1 quart to 1 gallon per cubic yard of media) and sprinkling or spraying it onto the media. Use the higher application rates for longer periods of control.

Potting Media Bulk Density (lbs. per cubic yard)	Fluid ounces of Talstar Nursery Flowable in one cubic yard			
	10 PPM	15 PPM	20 PPM	25 PPM
200	0.4	0.6	0.8	1.0
300	0.6	0.9	1.2	1.5
400	0.8	1.2	1.6	2.0
500	1.0	1.5	2.0	2.5
600	1.2	1.8	2.4	3.0
700	1.4	2.1	2.8	3.5
800	1.6	2.4	3.2	4.0
900	1.8	2.7	3.6	4.5
1000	2.0	3.0	4.0	5.0

The application rates listed above are based on the dry bulk density of the potting media. Use proportional volumes of Talstar Nursery Flowable for potting media with dry bulk densities that are not listed above.

**Black Vine Weevil Larval Control - Curative Treatment - Topical Drench:** To control black vine weevil larvae infesting containerized plants, dilute Talstar Nursery Flowable at the rate of 10 to 40 fl. ozs. (0.05 to 0.2 lb AI) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container.

**Bare-root Treatment for Preventative Root Weevil Larval Control:** To protect treated roots of field grown nursery stock from feeding by root weevil larvae, dilute one gallon of Talstar Nursery Flowable in 100 gallons of water and treat the bare roots of plants that are being transplanted into the field either by dipping the roots into the insecticide solution for ten seconds or by spraying the insecticide solution onto the roots.

**Diaprepes Weevil Larval Control - Curative Treatment - Topical Drench:** To control Diaprepes weevil larvae infesting containerized plants, dilute Talstar Nursery Flowable at the rate of 10 to 40 fl. ozs (0.05 to 0.2 lb AI) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or

12  
10

greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container.

**Fungus Gnat Larval Control - Preventative Treatment - Topical Drench:** For preventative control of fungus gnat larvae in containerized plants, dilute Talstar Nursery Flowable at the rate of 20 to 40 fl. ozs (0.1 to 0.2 lb AI) per 100 gallons and apply as a drench at the rate of 4 to 8 fl. ozs of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container. Use the higher application rate for a longer period of control.

**Fungus Gnat Larval Control - Curative Treatment - Topical Drench:** To control fungus gnat larvae infesting containerized plants, dilute Talstar Nursery Flowable at the rate of 10 to 40 fl. ozs. (0.05 to 0.2 lb AI) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container. Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter. Ideally, the media should be treated to the point of saturation, which generally requires 1/5 the volume of the container.

**Conditions of Sale and Limitation of Warranty and Liability:**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. EXCEPT AS WARRANTER BY THIS LABEL, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.**

To the extent consistent with applicable law FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Talstar and FMC—Trademarks of FMC Corporation

## Supplemental Label

**RESTRICTED USE PESTICIDE  
TOXIC TO FISH AND AQUATIC ORGANISMS**  
For retail sale to and use by certified applicators, or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

### TALSTAR NURSERY FLOWABLE INSECTICIDE/MITICIDE

**ADDITIONAL PERSONAL PROTECTIVE EQUIPMENT  
AND EXTENDED  
REAPPLICATION INTERVALS REQUIREMENTS FOR  
GREENHOUSE USE IN CALIFORNIA.  
EPA Reg. No. 279-3155**

#### DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This supplemental labeling must be in possession of all users of the product in greenhouses in California.

#### CALIFORNIA SPECIFIC REQUIREMENTS FOR GREENHOUSE APPLICATORS AND HARVESTERS:

In addition to following all applicable precautionary statements on the label on the product container, the following is required for greenhouse applicators and harvesters:

**Greenhouse Applicator:** Greenhouse applicators must wear a full body chemical-resistant protective suit (such as barrier laminate, butyl rubber, nitrile rubber, polyvinyl chloride, or equivalent).

**Reapplication Interval:** Reapplications to greenhouses must be at intervals of 30 days or longer.

**Greenhouse Harvesters:** Greenhouse harvesters must wear either regular-length gloves plus a long sleeved shirt or elbow-length (gauntlet type) gloves during the 30 days following application.

PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE CONTAINER LABEL.



FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia, PA 19103