



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 1, 2021

Shannon Cavanaugh
Senior Registration Manager
FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Subject: Label Amendment – Revised the existing label rates for canola
Product Name: Command 3ME Microencapsulated Herbicide
EPA Registration Number: 279-3158
Application Date: July 20, 2020
Decision Number: 565021

Dear Ms. Cavanaugh:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. The next label printing of this product must use this labeling unless subsequent changes have been approved. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 279-3158
Decision No. 565021

with FIFRA section 6. If you have any questions, please contact Aleah Holt at 703-347-0482 or by email at holt.aleah@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Shaja B. Joyner". The signature is written in a cursive style with a large initial 'S'.

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

ACCEPTED

07/01/2021

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 279-3158

Clomazone Group 13 Herbicide



**For Agricultural or Commercial Use Only
NOT FOR USE IN CALIFORNIA**

For control of weeds on Bean (dry and succulent), Brassica head and stem vegetable group 5-16, Chinese broccoli, Cottonseed subgroup 20C, Cucurbit Vegetables Crop Group 9, Herb Subgroup 19A, Kohlrabi, Peas (Succulent), Peppermint tops, Peppers, Rapeseed (canola varieties only), Rhubarb, Rice, Soybeans, Spearmint tops, Stalk and Stem Vegetable Subgroup 22A except kohlrabi, Sugarcane, Tobacco, Tuberos and Corm Vegetables subgroup 1D except Potato, and Vegetable Soybean (Edamame).

EPA Reg. No. 279-3158 **EPA Est.** _____

Active Ingredient:	By Wt.
Clomazone: 2-(2-Chlorophenyl)methyl-4, 4-dimethyl-3-isoxazolidinone	31.1%
Other Ingredients:	<u>68.9%</u>
	100.0%

Contains 3.0 pounds of active ingredient per gallon
U.S. Patent No. 38,675

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass by mouth of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give anything to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

For Emergency Assistance Call (800) 331-3148.

See other panels for additional precautionary information.

Sold By



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

Net Contents: _____

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Caution

Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves; Barrier Laminate or Butyl Rubber, or Nitrile Rubber or Viton;
- Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from the area treated. **DO NOT** apply where runoff is likely to occur. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as specified on this label.

Physical/Chemical Hazards

DO NOT use or store near heat or open flame.

Non-target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

TABLE OF CONTENTS

Read all information on pages 1-X before referring to specific crop use.

	Page
Ingredients.....	X
First Aid	X
Precautionary Statements.....	X
Directions for Use	X
Storage and Disposal	X
Resistance Management.....	X
Mixing and Handling Instructions	
for Bulk/Mini-Bulk Containers.....	X
Application Directions	X
Buffer and Application Restrictions.....	X
Mandatory Spray Drift	X
Rotational Cropping Precautions.....	X
Sprayer Cleanup.....	X
Mixing Instructions.....	X
Product Use Rate to Active Ingredient Conversion Chart.....	X
Application Instructions.....	X
Command Geographical Areas.....	X
ROW CROPS	
Cottonseed subgroup 20C (Cotton).....	X
Rice.....	X
Soybeans	X
Sugarcane.....	X
Tobacco.....	X

VEGETABLE CROPS

Stalk and Stem Vegetable Subgroup 22A (Asparagus and other crops in this subgroup except kohlrabi)	X
Beans (dry and succulent, except lima)	X
Brassica head and stem vegetable group 5-16 (Cabbage and Broccoli, etc.)	X
Chinese broccoli	X
Cucurbit Vegetables Crop Group 9: (Cucumbers, Melons (Muskmelons and Watermelons), Pumpkins, (Squash summer and winter)	X
Edamame (vegetable soybean)	X
Kohlrabi	X
Lima beans	X
Peas (Succulent)	X
Peppers	X
Rhubarb	X
Sweet Potatoes	X
Tuberous and Corm Vegetables Subgroup 1D (and other tuberous and corm vegetables, except sweet potatoes)	X
OIL PLANTS	
Rapeseed (canola varieties only)	X
Peppermint and Spearmint tops	X
HERBS Subgroup 19A	X

PRECAUTIONS

Off-site movement of spray drift or vapors of Command® 3ME Microencapsulated Herbicide can cause foliar whitening or yellowing of some plants.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product by air or through any type of irrigation system.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Prior to making applications, read and strictly follow all precautions and instructions in the APPLICATION DIRECTIONS, BUFFER & APPLICATION RESTRICTIONS and MANDATORY SPRAY DRIFT sections.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls; Chemical-resistant gloves; Barrier Laminate or Butyl Rubber, or Nitrile Rubber or Viton; Shoes plus socks.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

STORE ABOVE -4°F TO KEEP PRODUCT FROM FREEZING. If frozen, thaw before use. Observe recirculation directions under Mixing and Handling Instructions for Bulk/Mini-Bulk Containers.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put concentrate or dilute material into food or drink containers.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and spills): (800) 424-9300

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Plastic Non-refillable Containers: **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Sealed Containers: Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. **DO NOT** rinse container. **DO NOT** empty remaining formulated product. **DO NOT** break seals. Return intact to point of purchase. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat rinsing procedure two more times.

Mini-Bulk Containers: These containers are property of FMC Corporation and are returnable to FMC at FMC's discretion. These containers are provided for repackaging of Command 3ME* and must not be filled with any other product.

Bulk Drums: Command 3ME Bulk drums are returnable to FMC Corporation for reuse when the container is completely empty. Bulk drums containing product in excess of 1 gallon cannot be accepted for return.

Container Precautions

Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices. After filling and before transporting, check for leaks.

Container Restrictions

DO NOT refill or transport a damaged or leaking container.

* Any dealer wishing to repackage Command 3ME must comply with all Federal, State and local laws pertaining to bulk herbicide handling and possess a signed repackaging agreement from FMC Corporation.

WEED RESISTANCE MANAGEMENT

Command 3ME, which contains the active ingredient clomazone is a group 13 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches) with a burndown herbicide.
- To minimize resistance, apply full labeled rates of Command 3ME.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.
- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 13 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad-spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 13 herbicides.
- Avoid making more than two applications of Command 3ME and any other Group 13 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

MIXING AND HANDLING INSTRUCTIONS FOR BULK/MINI-BULK CONTAINERS

110 and 120 Gallon Compack Containers

Equipment Requirements:

Command 3ME Microencapsulated Herbicide is a microencapsulated herbicide and requires a diaphragm type pump in order to maintain product quality. **DO NOT use gear or piston-type pumps.** Bulk/Minibulk containers have been prefitted with a Scienco DD6 diaphragm type pump for recirculation and dispensing of product.

Dispensing Instructions:

When ready to dispense Command 3ME Microencapsulated Herbicide from the Compack, the applicator needs to recirculate the product in the container, if the product has settled or separated, for approximately 5 minutes or until the product is thoroughly turned over. The Scienco DD6 pump is equipped with recirculation capability. This allows for quick and efficient mixing of product which may have separated or settled in storage. To recirculate, press down the bypass pushrod lever to the locked position (slid under the motor) and turn on the motor. The discharge ball valve at the end of the hose must be closed before turning on the motor. Once the product is thoroughly recirculated the applicator may then begin the process of dispensing Command 3ME Microencapsulated Herbicide into the spray tank, loading or mixing system.

The dealer/applicator must wear proper clothing as listed on the Command 3ME Microencapsulated Herbicide label.

250 Gallon Bulkdrum III Containers

Equipment Requirements:

Command 3ME Microencapsulated Herbicide is a microencapsulated herbicide and requires a diaphragm type pump in order to maintain product quality. **DO NOT use gear or piston-type pumps.** The following pumps are suitable for moving Command 3ME Microencapsulated Herbicide from the 250 gallon Bulkdrum into spray tanks, mixing systems, or dedicated repackaging minibulk tanks*:

- Tuthill Fill-Rite Chemtraveller portable transfer pump
- Scienco Caddy-SS portable transfer pump

Scienco DD6 pump

Tuthill Fill-Rite Series 400 diaphragm pumps

Dispensing Instructions:

When ready to dispense Command 3ME Microencapsulated Herbicide from the Bulkdrum, the dealer/applicator must recirculate the product in the container, for at least 15 minutes. This can be done by hooking a portable pump such as listed above to the bottom bung, opening the valve and directing the outlet nozzle into the 6" top port (seal must be broken). Directing the nozzle stream into the corners will ensure more complete turning of the product volume. Once the product is thoroughly circulated, the dealer/applicator may then begin the process of dispensing Command 3ME Microencapsulated Herbicide into the dedicated repackaging minibulk container(s)*, or spray tank, loading or mixing system. Rinse the empty Bulkdrum container and transfer the rinsate directly to the mix or spray tank.

The dealer/applicator must wear proper clothing as listed on the Command 3ME Microencapsulated Herbicide label.

* Any dealer wishing to repack Command 3ME Microencapsulated Herbicide must comply with all Federal, State and local laws pertaining to bulk herbicide handling and possess a signed repackaging agreement from FMC Corporation.

RESTRICTIONS

- **DO NOT** use gear or piston-type pumps
- The dealer/applicator must wear proper clothing as listed on the Command 3ME Microencapsulated Herbicide label

APPLICATION DIRECTIONS IMPORTANT

FAILURE TO OBSERVE THE DIRECTIONS IN THIS SECTION OF THE LABEL MAY RESULT IN INJURY TO SENSITIVE PLANTS

- The microencapsulation of clomazone, the active ingredient in Command 3ME Microencapsulated Herbicide, is intended to minimize movement away from the site of application. Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Leave an adequate buffer zone between the area to be treated and desirable plants.
- **PRECAUTIONS:** Foliar contact with spray drift or vapors may cause foliar whitening or yellowing of sensitive plants. Symptoms are generally temporary in nature, but may persist on some plant species.

Application restrictions must be taken as follows:

- Observe all buffer restrictions.
- Before application, determine air movement and direction.

Refer to individual crop use directions for additional requirements.

BUFFER RESTRICTIONS

- **DO NOT** apply within 300 feet of downwind crops and desirable non-target plants.
- **DO NOT** apply within 1,200 feet of Towns and Housing Developments, Commercial Fruit, Nut, or Vegetable Production¹, Commercial Greenhouses or Nurseries.

¹ Except for vegetable crops registered for use on the Command 3ME herbicide label.

APPLICATION RESTRICTIONS

- **DO NOT** apply this product by air or through irrigation equipment.
- **DO NOT** apply Command 3ME Microencapsulated Herbicide to non-field areas including fence rows, waterways, ditches, and roadsides.
- When moving spray equipment to noncontiguous sites, **DO NOT** allow spray solution to spray or drip from tanks, hoses, fittings or spray nozzles and tips.
- **DO NOT** apply when weather conditions favor drift.

MANDATORY SPRAY DRIFT MANAGEMENT

GROUND BOOM APPLICATIONS

- User must only apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

ROTATIONAL CROPPING PRECAUTIONS

Under some conditions, temporary whitening or yellowing of leaves may occur on approved rotational crops where undesirable soil residues of Command 3ME exist.

Under abnormal conditions, carryover injury to rotational crops can occur. The following factors can contribute to increased risk of injury to rotational crops:

- 1) Over-application resulting from use of worn nozzles, excessive over-lapping of spray swaths, failing to shut off spray booms when turning (end row areas), or slowing or stopping sprayer.
- 2) Soil with pH less than or equal to 5.9.
- 3) Extreme dryness in the four months following application.
- 4) Choice of rotational crop hybrid.

Additional instructions to prevent rotational crop injury may be provided in the form of service bulletins for locations where risk of injury is significantly increased due to extremely dry conditions.

Refer to Rotational Crop Instructions and Replanting Instructions of specific crops for additional crop planting information.

SPRAYER CLEANUP

DO NOT drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or move into contact with their roots. **DO NOT** contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean-up instructions noted below to prevent spray tank residues from damaging other crops.

Sprayer equipment need to be thoroughly rinsed to remove residues of herbicide that might injure other subsequently sprayed crops. The steps below need to be followed for the thorough rinsing of spray equipment following applications of Command 3ME Microencapsulated Herbicide or tank mixes of Command 3ME Microencapsulated Herbicide with other labeled products.

- 1) Drain any remaining spray solution from tank, pump, hoses and boom and discard in an approved manner (See Note that follows).
- 2) Clean tank and fittings by:

- Thoroughly hosing down the inside walls of the spray tank with a quantity of water equal to 1/8 of the total tank capacity and operating the pump to circulate this solution through the sprayer system for 15 minutes.
 - Washing down the outside surfaces of equipment.
 - Removing nozzle tip and screen from end nozzle in each boom section and allowing several gallons of rinsate solution to flush completely through boom (collect rinsate while flushing).
- 3) Thoroughly drain remaining rinsate solution from tank, pump and hoses. Combine with boom flushing and dispose of all rinsates from this first rinsing in an approved manner (see Note that follows).
 - When switching from water dilutions to applications utilizing crop oil or liquid fertilizer as a carrier, a small volume of crop oil or liquid fertilizer need to be flushed through the tank, pump, hoses, and boom prior to the next use. Dispose of crop oil or liquid fertilizer rinsate in an approved manner (see Note for local, state and federal guidelines).
 - 4) Remove the remaining nozzle tips, and screens and the line filter and wash in a pail of warm soapy water, thoroughly rinse and replace.
 - 5) Hose down the inside walls of the spray tank a second time and circulate this solution using the same procedure as noted in #2 above.
 - 6) If the next use of the sprayer will be for applying a pre-emergent or preplant incorporated pesticide on any crop for which Command 3ME is registered, rinsate from this second rinsing may be utilized by diluting with the water for the next pesticide load;

HOWEVER

If the next use of the sprayer will be a post-emergence applied pesticide on any crop, drain rinsate solution from this second rinsing. Retain rinsate solution for use only with a soil incorporated pesticide to be applied on a crop for which Command 3ME Microencapsulated Herbicide is registered. Refill tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

NOTE: Dispose of excess spray mixture and/or *rinsate from first tank rinsing* by application to cropland as described on this label. If excess spray mixture and/or *rinsate from first rinsing* cannot be disposed of according to label instructions, dispose of in compliance with local, state and federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

MIXING INSTRUCTIONS

Care must be taken when mixing Command 3ME Microencapsulated Herbicide. Avoid mixing in areas adjacent to desirable plants.

Command 3ME Microencapsulated Herbicide Alone: Mix Command 3ME Microencapsulated Herbicide with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 3ME Microencapsulated Herbicide, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Fill spray tank one-fourth to one-third full with water; with agitator operating add the specified amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Mix thoroughly and fill tank one-half full continuing agitation. Add Command 3ME Microencapsulated Herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is advised, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing agents, follow specific product label instructions for order of addition to spray tank.

Fertilizer Spray Mixtures: Applications of Command 3ME Microencapsulated Herbicide alone, or with advised tank mixtures, in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Small quantities need to be tested for compatibility by the following procedure before mixing tank full quantities.

- 1) Put 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the table below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

Herbicide	Rate/Acre	Amount to Be Added Per Pint of Fluid Fertilizer*
Wettable Powders	1/2 pound	3/4 level teaspoon
or	1 pound	1 1/2 level teaspoons
or	2 pounds	3 level teaspoons
Dry Flowables (Dry Granules)	3 pounds	4 1/2 level teaspoons
Emulsifiable Concentrates	1 pint	1/2 level teaspoon
or	1 quart	1 level teaspoon
or	2 quarts	2 level teaspoons
Flowables	3 quarts	3 level teaspoons

* Based on a spray volume of 25 gpa. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below:
 - 1) **Command 3ME Microencapsulated Herbicide Alone:** Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.
 - 2) **Command 3ME Microencapsulated Herbicide plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Continue agitation during application.

PRODUCT USE RATE TO ACTIVE INGREDIENT CONVERSION CHART

Product Use Rate to Active Ingredient Conversion Chart	
Amount of COMMAND 3ME in Fluid Ounces per Acre	Amount of the Active Ingredient Clomazone in Pound(s) per Acre
3.2	0.075
4	0.094
4.7	0.11
6	0.14
6.4	0.15
7.3	0.17
8.1	0.19
8.5	0.2
10.7	0.25
11.1	0.26
12.8	0.3
14.1	0.33
14.9	0.35
16.2	0.38
17.1	0.4
20.1	0.47
21.3	0.5
24	0.56
26.9	0.63
28.2	0.66
32	0.75
34.1	0.8
35.8	0.84
40.1	0.94
42.7	1
48	1.12
53.3	1.25
64	1.5

APPLICATION INSTRUCTIONS

Ground Applications

Broadcast or Banded Applications: Apply Command 3ME Microencapsulated Herbicide alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. Use nozzles suitable for broadcast boom or banded application of herbicides. See "APPLICATION DIRECTIONS", "BUFFER & APPLICATION RESTRICTIONS and MANDATORY SPRAY DRIFT sections for specific instructions to reduce spray drift. For Command 3ME Microencapsulated Herbicide tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers need to be no finer than 50-mesh.

Command 3 ME Microencapsulated Herbicide may be used as a pre-emergent soil surface applied treatment from 30 days before planting to just prior to crop emergence. If field conditions indicate the need for additional seedbed preparation, the use of equipment which will move the herbicide no deeper than 1 1/2 - 2" is acceptable.

Banded Applications—Calculate the rates and volumes required by using the following formulas:


$$\frac{\text{Band width in inches}}{\text{Band width in inches}} \times \text{Broadcast Rate Per Acre} = \text{Band rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Volume Per Acre} = \text{Band volume per acre}$$

Cultivation After Planting

If cultivation is necessary because of soil crusting, compaction, or weed emergence, soil treated with Command 3ME Microencapsulated Herbicide may be shallow-cultivated (no deeper than 1 1/2 - 2") or rotary-hoed without reducing the weed control activity of Command 3ME Microencapsulated Herbicide. Avoid deep cultivation which may bring untreated soil to the surface and result in poor weed control.

**COMMAND GEOGRAPHICAL AREAS
For Rates, Weed Control, and Crop Rotation**



* Southern area includes Missouri bootheel which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

ROW CROPS

COTTONSEED SUBGROUP 20C: Cotton, Cottonseed; cultivars, hybrids, and/or varieties of these

PRODUCT INFORMATION

Command 3ME Microencapsulated Herbicide may be applied, as a pre-emergent banded or broadcast treatment in conventional tillage or conservation tillage systems for the control of annual grasses and broadleaf weeds in (cottonseed subgroup 20C) that includes cotton, cottonseed, cultivars, hybrids, and/or varieties.

APPLIED IN COMBINATIONS FOR COTTON RESTRICTIONS

DO NOT incorporate tank mixtures of Command 3ME and other products.

REQUIREMENTS FOR PLANTING TIME APPLICATIONS

Failure to apply either disulfoton or phorate insecticides with Command 3ME Microencapsulated Herbicide in accordance with in-furrow label use directions can result in crop phytotoxicity (bleaching) and/or stand reduction. Combinations of at planting systemic granular carbamate and organophosphate insecticides in conjunction with Command 3ME Microencapsulated Herbicide may result in injury to cotton. Crop injury may occur with higher Command 3ME Microencapsulated Herbicide rates on sandy soils. Diuron is not advised at planting when Command 3ME Microencapsulated Herbicide is used as plant injury may result.

Refer to the insecticide product labels for appropriate in-furrow application directions and maximum use rates. Monitor application equipment to insure accurate and uniform placement of the insecticide.

PRE-EMERGENT BANDED APPLICATIONS

For pre-emergent banded treatments, refer to the directions for banded applications in the APPLICATION INSTRUCTIONS section.

REPLANTING INSTRUCTIONS

If the initial planting of cotton fails to produce a uniform stand, cotton may be replanted in fields treated with Command 3ME Microencapsulated Herbicide alone. When tank mixing with a labeled product, refer to the replant instructions for that product. **DO NOT** replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP RESTRICTIONS on the Command 3ME Microencapsulated Herbicide label. Where a tank mix is used, refer to the product's labels for any additional replant instructions.

If replanting is required follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS noted above.

Command 3ME Microencapsulated Herbicide APPLIED ALONE

RATES AND WEEDS CONTROLLED¹

21.3 FL. OZ. (0.50 LB. A.I.) GRASSES	32 FL. OZ. (0.75 LB. A.I.) GRASSES	42.7 FL. OZ. (1.0 LB. A.I.) GRASSES	53.3 FL. OZ. (1.25 LB. A.I.) GRASSES
Suppression of some annual grasses	Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Foxtail —Giant —Green Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass Field Sandbur*	All grasses controlled at 32 fl oz <i>plus</i> : Cupgrass —South-western —Woolly Field Sandbur Bermudagrass* Red Rice* Itchgrass ³	All grasses controlled at 42.7 fl oz <i>plus</i> : Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*

21.3 ² FL. OZ. (0.50 LB. A.I.) BROADLEAVES	32 FL. OZ. (0.75 LB. A.I.) BROADLEAVES	42.7 FL. OZ. (1.0 LB. A.I.) BROADLEAVES	53.3 FL. OZ. (1.25 LB. A.I.) BROADLEAVES
Velvetleaf Spurred Anoda	Velvetleaf Spurred Anoda <i>plus</i> : Common Ragweed* Jimsonweed* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed Venice Mallow	All broadleaves controlled at 32 fluid ounces <i>plus</i> : Cocklebur Common Ragweed Dayflower Florida Beggarweed Florida Pusley Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia ⁴	All broadleaves controlled at 2 42.7 fluid ounces <i>plus</i> : Balloonvine* Black Nightshade* Curly Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth*

* Partially Controlled

1 Use higher label rates (a minimum of 42.7 fluid ounces per acre) for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, use a minimum application rate of 53.3 fluid ounces per acre for cocklebur or common ragweed control.

2 For heavy weed pressure or heavy soils use 26.9 fluid ounces per acre to control Velvet-leaf or Spurred Anoda.

3 For control of itchgrass, **DO NOT** incorporate Command 3 ME Microencapsulated Herbicide. For high weed pressure and heavy soils use 53.3 fluid ounces.

4 Use In AR/LA/MS/TX/OK only—for high pressure and heavy soils use 53.3 fluid ounces.

Command 3ME Microencapsulated Herbicide MAY BE APPLIED WITH OTHER PREEMERGENT HERBICIDES REGISTERED FOR THESE USES ON COTTON.

Command 3ME Microencapsulated Herbicide may be applied in combination with other herbicides registered for use in cotton to broaden the weed control spectrum. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed below. Refer to sections headed "Rotational Cropping Precautions."

Note: When using Command 3ME Microencapsulated Herbicide with other registered herbicides always refer to rotational restrictions and directions on the other product's label.

21.3 FL. OZ. (0.50 LB. A.I.) ANYTIME	32 to 42.7 FL. OZ. (0.75 to 1.0 LB. A.I.) ANYTIME	53.3 FL. OZ. (1.25 LB. A.I.) ANYTIME
Cabbage (transplanted) Cotton* Peas Peppers Pumpkins (processing) Rice Soybeans Squash Sweet Potato Tobacco Tuberous & Corm Veggies.	Cotton* Peppers Rice Soybeans Squash (winter)(32 fl. oz.) Sweet Potato Tobacco Tuberous & Corm Veggies.	Cotton* Soybeans Sweet Potato Tobacco Tuberous & Corm Veggies.

9 MONTHS	9 MONTHS	9 MONTHS
All crops listed above <i>plus:</i> Cabbage (direct seeded) Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Snap Beans Sorghum Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash (summer) Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Peppers Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash Sugar Beets Tomatoes (Transplanted)

12 MONTHS	12 MONTHS	12 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas.

* Refer to the section "Requirements for Planting Time Applications" and "Replanting Instructions."

RESTRICTIONS

- **DO NOT** apply Command 3ME within 65 days of harvest.
- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per acre per year.
- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per application.
- **DO NOT** allow livestock to graze on treated cotton forage or trash, or feed treated cotton forage or trash to livestock.

- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME treatment.
- **DO NOT** apply Command 3ME Microencapsulated Herbicide to cotton unless either disulfoton or phorate organophosphate insecticide is applied in-furrow with the seed at planting time at a minimum of 0.75 pound per acre of active ingredient.
- **DO NOT** reduce the application rate of the organophosphate insecticide when Command 3ME Microencapsulated Herbicide is applied as a banded treatment.
- **DO NOT** apply more than one application per year.

RICE

PRODUCT INFORMATION

Command 3ME Microencapsulated Herbicide may be utilized as a surface broadcast treatment prior to weed emergence, for the control of annual grass weeds in dry-seeded rice.

PRE-EMERGENT SURFACE BROADCAST APPLICATIONS

Command 3ME may be applied as a surface broadcast application 14 days prior to planting or up to 7 days after planting, but prior to weed emergence, using ground equipment in a minimum of 10 to 40 gallons of water per acre at the rate of 10.7 to 34.1 fl. oz. product (0.25 to 0.8 lb. active) per acre depending upon the soil texture. Refer to table below for specific rates and weeds controlled. For heavy soils use the higher specified rate, otherwise less than desirable weed control may result.

For pre-emergent broadcast treatments, refer to the directions for broadcast applications in the APPLICATION INSTRUCTIONS section.

EARLY POSTEMERGENCE APPLICATIONS

Command 3ME may be applied after planting as an early post-emergence treatment to rice through the five leaf stage to provide pre-emergence and residual control of grass weeds. Use ground equipment with nozzles that produce a course spray and a minimum of 10-40 gallons of water per acre. Refer to table below for specific rates and weeds controlled. For control of existing grass present at the time of application include a post-emergence herbicide registered for the control of grass species in rice. Consult post-emergence herbicide label for specific directions regarding use rates and stage of weeds and crop.

REPLANTING INSTRUCTIONS

If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with Command 3ME. When tank mixing with a labeled product, refer to the replant instructions for that product. When a tank mix is used, refer to the product's label for any additional rotational crop instructions.

Command 3ME Microencapsulated Herbicide Applied Alone - Single Application Pre-emergence or Post-emergence

RATES AND WEEDS CONTROLLED

Soil Texture	Broadcast Rates Per Acre
Coarse (light) Soils: (sand, loamy sand, sandy loam)	10.7 – 14.1 fl. oz. product (0.25 – 0.33 lb. ai)
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	17.1 – 21.3 fl. oz. product (0.4 - 0.5 lb. ai)
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	21.3 – 34.1 fl. oz. product (0.5 – 0.8 lb. ai)

*Select lower to higher rates based on lighter to heavier soils

GRASSES

Barnyardgrass (Watergrass)	(<i>Echinochloa crus-galli</i> , <i>E. colonum</i>)
Broadleaf signalgrass	
Crabgrass (Large and Smooth)	(<i>Urochloa platyphylla</i>)
Panicum	(<i>Digitaria spp.</i>)
—Common	(<i>Panicum spp.</i>)
—Fall	
—Texas	
Sprangletop	(<i>Leptochloa spp.</i>)

Partial weed control may result if levees are pulled after Command 3ME has been applied. Additional use of labeled post-emerge herbicide applications may be required.

SPLIT APPLICATIONS

Both pre-emergent surface broadcast applications and early post-emergence applications (out to the five leaf stage) of Command 3ME may be made during the same season to control the above weeds as long as the total yearly application does not exceed 34.1 fl oz/A product (0.8 lb. ai/A). Refer to the table below for specific rates.

Command 3ME Microencapsulated Herbicide Applied as Split Applications

Soil Texture	Pre-Emergent Application Broadcast Rates Per Acre	Post-Emergent Application Broadcast Rates Per Acre
Coarse (light) Soils: (sand, loamy sand, sandy loam)	10.7 fl. oz. product (0.25 lb. ai)	8.5 – 10.7 fl. oz. product (0.2 - 0.25 lb. ai)
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	12.8 – 14.9 fl. oz. product (0.3 – 0.35 lb. ai)	10.7 – 12.8 fl. oz. product (0.25 – 0.3 lb ai)
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	17.1 to 21.3 fl. oz. product (0.4 to 0.5 lb. ai)	12.8 – 17.1 fl. oz. product (0.3 to 0.4 lb. ai)

PRECAUTIONS

Application of Command 3ME Microencapsulated Herbicide to fields which have been precision leveled with deep cuts may result in rice crop injury including stand loss. Consult with rice specialists for soil amending practices which can reduce potential for herbicide injury in precision leveled fields.

ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed below. Refer to section headed "Rotational Cropping Precautions."

Note: When using Command 3ME with other registered herbicides always refer to rotational restrictions and directions on the other product's label.

10.7 – 34.1 fl. oz. (0.25 to 0.8 LB. A.I.)		
<u>ANYTIME</u>	<u>9 MONTHS</u>	<u>12 MONTHS</u>
Cotton*	All crops listed above	All crops
Peas	plus:	
Peppers	Cotton	
Pumpkins (processing)	Corn	
Rice	—Field	
Soybeans	—Pop	
Squash	—Seed	
Sweet Potato	—Sweet	
Tobacco	Cucurbits	
Tuberous and Corn Veg.	Dry Beans	
	Peanuts	
	Potatoes	
	Snap Beans	
	Sorghum	
	Sugar Beets	
	Tomatoes	
	(Transplanted)	

Cover crops, however, may be planted anytime but stand reductions may occur in some areas.

* Refer to the section "Requirements for Planting Time Applications" and "Replanting Instructions."

RESTRICTIONS

- **DO NOT** use this treatment in water-seeded rice.
- **DO NOT** apply Command 3ME on rice fields in which concurrent crayfish or catfish farming are included in the cultural practices.
- **DO NOT** use water containing Command 3ME residues from rice cultivation to irrigate food or feed crops which are not registered for use with Command 3ME.
- **DO NOT** apply more than 34.1 fl. oz. Command 3ME (0.8 lb. ai clomazone) per acre per application.
- **DO NOT** apply more than 34.1 fl. oz. Command 3ME (0.8 lb. ai clomazone) per acre per year.
- **DO NOT** apply more than one application as a pre-emerge application prior to rice emergence.
- **DO NOT** apply more than one application as a post-emergence application after rice emergence up to five leaf-rice.
- **DO NOT** retreat fields with a second pre-emergent application of Command 3ME.
- **DO NOT** make a post application within seven days when a pre-emerge application was used.
- **DO NOT** replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP INSTRUCTIONS on the Command 3ME label.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME treatment.

SOYBEANS

PRODUCT INFORMATION

Command 3ME Microencapsulated Herbicide may be utilized as a pre-emergent soil applied treatment for the control of annual grass and broadleaf weeds in soybeans. Command 3ME Microencapsulated Herbicide may be surface applied prior to planting up to 3 days after planting but prior to crop emergence.

Command 3ME Microencapsulated Herbicide may be tank mixed with other pre-emergent soybean herbicides to broaden the weed control spectrum compared to the products applied alone. It is the pesticide user's responsibility to ensure that all products are

registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Special attention must be paid to rotational cropping restrictions, potential carryover and crop injury warnings of each product's label when tank mixing.

Water or liquid fertilizer may be used as a carrier for Command 3ME Microencapsulated Herbicide when applied alone, or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

APPLIED IN COMBINATIONS FOR SOYBEANS RESTRICTIONS

DO NOT incorporate tank mixtures of Command 3ME and other products.

REPLANTING INSTRUCTIONS

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 3ME Microencapsulated Herbicide alone (or with specified tank mixtures). When tank mixing with a labeled product, refer to the soybean replant instructions for that product.

Soybeans - Northern Area—See Map

COMMAND® 3ME Microencapsulated Herbicide

APPLIED ALONE

RATES AND WEEDS CONTROLLED¹

21.3 FL. OZ. (0.50 LB. A.I.) GRASSES	32 FL. OZ. (0.75 LB A.I.) GRASSES	42.7 FL. OZ. (1.0 LB. A.I.) GRASSES
Suppression of some annual grasses	Barnyardgrass Crabgrass —Large —Smooth Field Sandbur Foxtail —Giant —Green —Robust —Yellow Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass*	All grasses controlled at 32 fluid ounces <i>plus:</i> Bermudagrass* Broadleaf Signalgrass Cupgrass —Southwestern —Woolly* Seedling Johnsongrass Shattercane* Wild Proso Millet*

21.3 FL. OZ. ² (0.50 LB. A.I.) BROADLEAVES	32 FL. OZ. (0.75 LB A.I.) BROADLEAVES	42.7 FL. OZ. (1.0 LB. A.I.) BROADLEAVES
Velvetleaf Spurred Anoda	Velvetleaf Spurred Anoda <i>plus:</i> Common Ragweed* Galinsoga Jimsonweed* Lambsquarters Prickly Sida Purslane Smartweed* —Ladysthumb* —Pennsylvania* Venice Mallow	All broadleaves controlled at 32 fluid ounces <i>plus:</i> Black Seeded Plantain Cocklebur* Common Ragweed Jimsonweed Kochia Smartweed —Ladysthumb —Pennsylvania Tropic Croton

* Partially Controlled

1 Use higher label rates (a minimum of 42.7 fluid ounces per acre) for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types.

2 For heavy weed pressure or heavy soils use 26.9 fluid ounces per acre to control Velvetleaf or Spurred Anoda.

ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed below. Refer to section headed "Rotational Cropping Precautions."

NOTE: When using Command 3ME Microencapsulated Herbicide with other registered herbicides always refer to rotational restrictions and directions on the other product's label.

21.3 FL. OZ. (0.50 LB. A.I.) ANYTIME	32 FL. OZ. (0.75 LB A.I.) ANYTIME	42.7 FL. OZ. (1.0 LB. A.I.) ANYTIME
Cabbage (transplanted) Cotton* Peas Peppers Pumpkins (processing) Rice Soybeans Squash Sweet Potato Tobacco Tuberous and Corm Veg.	Cotton* Peppers Rice Soybeans Squash (Winter) Sweet Potato Tobacco Tuberous and Corm Veg.	Cotton* Peppers Soybeans Sweet Potato Tobacco Tuberous and Corm Veg.

9 MONTHS	9 MONTHS	9 MONTHS
All crops listed above <i>plus:</i> Cabbage (Direct Seeded) Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Potatoes Snap Beans Sorghum Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash (Summer) Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash Sugar Beets Tomatoes (Transplanted)

12 MONTHS	12 MONTHS	12 MONTHS
All crops listed above <i>plus:</i> Tomatoes (all) Wheat	All crops listed above <i>plus:</i> Tomatoes (all) Wheat	All crops listed above <i>plus:</i> Seed Corn Sweet Corn Tomatoes (all) Wheat

16 MONTHS	16 MONTHS	16 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. **DO NOT**

* Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

RESTRICTIONS

- **DO NOT** allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment.
- **DO NOT** apply more than one application per year.
- **DO NOT** replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Instructions on this label.

- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per application.
- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per year.

Soybeans - Southern Area—See Map

COMMAND® 3ME Microencapsulated Herbicide APPLIED ALONE RATES AND WEEDS CONTROLLED¹

21.3 FL. OZ. (0.50 LB. A.I.) GRASSES	32 FL. OZ. (0.75 LB. A.I.) GRASSES	42.7 FL. OZ. (1.0 LB. A.I.) GRASSES	53.3 FL. OZ. (1.25 LB. A.I.) GRASSES
Suppression of some annual grasses	Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Foxtail —Giant —Green Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass Field Sandbur*	All grasses controlled at 32 fluid ounces <i>plus:</i> Cupgrass —South-western —Woolly Field Sandbur Bermudagrass* Red Rice* Itchgrass ³	All grasses controlled at 42.7 fluid ounces <i>plus:</i> Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*

21.3 FL. OZ. (0.50 LB. A.I.) BROADLEAVES	32 FL. OZ. (0.75 LB. A.I.) BROADLEAVES	42.7 FL. OZ. (1.0 LB. A.I.) BROADLEAVES	53.3 FL. OZ. (1.25 LB. A.I.) BROADLEAVES
Velvetleaf Spurred Anoda	Velvetleaf Spurred Anoda <i>plus:</i> Common Ragweed* Jimsonweed* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed Venice Mallow	All broadleaves controlled at 32 fluid ounces <i>plus:</i> Cocklebur Common Ragweed Dayflower Florida Beggarweed Florida Pusley Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia ⁴	All broadleaves controlled at 42.7 fluid ounces <i>plus:</i> Balloonvine* Black Nightshade* Curly Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth* Prostrate Spurge

* Partially Controlled

¹ Use higher label rates (a minimum of 42.7 fluid ounces per acre) for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, use a minimum application rate of 53.3 fluid ounces per acre for cocklebur or common ragweed control.

² For heavy weed pressure or heavy soils use 26.9 fluid ounces per acre to control Velvetleaf or Spurred Anoda.

³ For control of Itchgrass, **DO NOT** incorporate Command 3ME Microencapsulated Herbicide. For high weed pressure and heavy soils use 53.3 fluid ounces.

⁴ Use in AR/LA/MS/TX/OK only—for high pressure and heavy soils use 53.3 fluid ounces.

ROTATIONAL CROP RESTRICTIONS

Rotate to Crops as listed below. Refer to section headed "Rotational Cropping Precautions."

NOTE: When using Command 3ME Microencapsulated Herbicide with other registered herbicides always refer to rotational restrictions and directions on the other product's label.

21.3 FL. OZ. (0.50 LB. A.I.) ANYTIME	32 to 42.7 FL. OZ. (0.75 to 1.0 LB. A.I.) ANYTIME	53.3 FL. OZ. (1.25 LB. A.I.) ANYTIME
Cabbage (transplanted) Cotton* Peas Peppers Pumpkins (processing) Rice Soybeans Squash Sweet Potato Tobacco Tuberous & Corm Vegs.	Cotton* Peppers Rice Soybeans Squash (winter)(32 FL. OZ.) Sweet Potato Tobacco Tuberous & Corm Vegs.	Cotton* Soybeans Sweet Potato Tobacco Tuberous & Corm Vegs.

9 MONTHS	9 MONTHS	9 MONTHS
All crops listed above <i>plus:</i> Cabbage (direct seeded) Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Snap Beans Sorghum Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash (summer) Sugar Beets Tomatoes (Transplanted)	All crops listed above <i>plus:</i> Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Peppers Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash Sugar Beets Tomatoes (Transplanted)

12 MONTHS	12 MONTHS	12 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. **DO NOT**

* Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

RESTRICTIONS

- **DO NOT** allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME treatment.
- **DO NOT** apply more than one application per year.
- **DO NOT** replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Instructions on this label.
- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per application.
- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per year.

SUGARCANE

PRODUCT INFORMATION

Command 3ME Microencapsulated Herbicide may be utilized as a pre-emergent soil applied treatment for the control of annual grass and broadleaf weeds in sugarcane.

Command 3ME may be used at rates of 42.7 to 53.3 fluid ounces (1.0 to 1.25 lb. active ingredient) per acre on newly planted or ratoon sugarcane using ground equipment in a minimum of 10 to 40 gallons of water per acre. Command 3ME may be applied in late summer or early fall to newly planted sugarcane and to newly planted or ratoon sugarcane as a pre-emerge in spring through layby.

RESTRICTIONS

DO NOT replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP RESTRICTIONS on the Command 3ME label.

DO NOT apply more than one application per year.

DO NOT apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per application.

DO NOT apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per year.

PRE-EMERGENT BANDED APPLICATION

For pre-emergent-banded treatments, refer to the directions for banded application in the Application Instructions section.

REPLANTING INSTRUCTIONS

If the initial planting of newly planted cane fails to produce a uniform stand sugarcane may be replanted in fields treated with Command 3ME alone. When tank mixing with a labeled product, refer to the replant instructions for that product. Where a tank mix is used, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If replanting is required follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS noted above.

Command 3ME Microencapsulated Herbicide APPLIED ALONE

RATES AND WEEDS CONTROLLED¹

42.7 FL. OZ. (1.0 LB. A.I.) GRASSES	53.3 FL. OZ. (1.25 LB. A.I.) GRASSES
Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Cupgrass —Southwestern —Woolly Field Sandbur Foxtail —Giant —Green Goosegrass Itchgrass ² Panicum —Common —Fall —Texas Red Rice* Seedling Johnsongrass	All grasses controlled at 42.7 fluid ounces plus: Bermudagrass ¹ Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*

42.7 FL. OZ. (1.0 LB. A.I.) BROADLEAVES	53.3 FL. OZ. (1.25 LB. A.I.) BROADLEAVES
Velvetleaf Spurred Anoda Cocklebur Common Ragweed Dayflower Florida Beggarweed Jimsonweed Kochia* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redvine* Redweed Tropic Croton Venice Mallow Wild Poinsettia	All broadleaves controlled at 42.7 fluid ounces plus: Balloonvine* Black Nightshade* Curly Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth*

*Partially Controlled

1 For Suppression, Command 3ME at 53.3 fluid ounces (1.25 lb. ai/A) applied immediately after planted sugarcane in late summer and early fall has been effective in delaying Bermudagrass reestablishment.

2 For control of Itchgrass, **DO NOT** incorporate Command 3ME. For heavy weed population and heavy soils use 53.3 fluid ounces (1.25 lb. ai/A).

Command 3ME Microencapsulated Herbicide may be applied in combination with other herbicides registered for use in sugarcane, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed below, otherwise crop injury may occur. Refer to sections headed "Rotational Cropping Precautions" and "Product Information" for more detailed information.

Note: When using Command 3ME Microencapsulated Herbicide with other registered herbicides, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

42.7 FL. OZ. (1.0 LB. A.I.)	53.3 FL. OZ. (1.25 LB. A.I.)
<u>ANYTIME</u> Cotton* Peppers Soybeans Sugarcane Sweet Potato Tobacco Tuberous & Corm Veggies.	<u>ANYTIME</u> Cotton* Sugarcane Soybeans Sweet Potato Tobacco Tuberous & Corm Veggies.
<u>9 MONTHS</u> All crops listed above <i>plus</i> : Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash Sugar Beets Tomatoes (Transplanted)	<u>9 MONTHS</u> All crops listed above <i>plus</i> : Cabbage Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Peas Peppers Potatoes Pumpkins (processing) Rice Snap Beans Sorghum Squash Sugar Beets Tomatoes (Transplanted)
<u>12 MONTHS</u> All crops	<u>12 MONTHS</u> All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas.

* Refer to the section "Requirements for Planting Time Applications" and "Replanting Instructions" in the cotton directions for use section.

RESTRICTIONS

- **DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai./A) per application.
- **DO NOT** apply more than 53.3 fluid ounces (1.25 lb. ai/A) per acre of Command 3ME Microencapsulated Herbicide during one year.
- **DO NOT** apply more than one application per year.
- **DO NOT** apply Command 3ME Microencapsulated Herbicide within 120 days prior to harvest on sugarcane for milling.
- **DO NOT** apply aerially or through irrigation equipment.
- **DO NOT** allow livestock to graze on treated sugarcane forage or trash, or feed treated sugarcane forage or trash to livestock.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment.

TOBACCO

PRODUCT INFORMATION

Command 3ME Microencapsulated Herbicide may be utilized as a soil applied treatment prior to weed emergence, for the control of annual grass and broadleaf weeds in tobacco.

PRE-EMERGENCE (Prior to Transplant)

Make a single broadcast application in a minimum of 20 gallons of water per acre at the rate of 32 to 42.7 fluid ounces (0.75-1.0 lb. ai) per acre. For heavy weed pressure or heavy soils use the higher specified rate, otherwise less than desirable weed control may result.

Apply immediately before, or up to 30 days prior to, transplanting. Apply as a surface treatment prior to emergence of weeds. If weeds emerge before pre-emergence application can be made, cultivate before or immediately preceding treatment. If field conditions indicate the need for additional seedbed preparation, the use of equipment which will move the herbicide no deeper than 1 1/2 - 2" is acceptable.

PRE-EMERGENCE (Post Transplant)

Make a single broadcast application in a minimum of 20 gallons of water per acre at the rate of 32 to 42.7 fluid ounces (0.75-1.0 lb. active) per acre. For heavy weed pressure or heavy soils use the higher specified rate, otherwise less than desirable weed control may result. Apply no more than once per year using a maximum rate of 42.7 fluid ounces (1.0 lb. ai/A) per acre. Apply over-the-top of tobacco plants immediately, or up to 7 days after, transplanting but prior to weed emergence. If weeds emerge before application can be made, cultivate before herbicide treatment to remove existing weeds.

COMMAND 3ME MICROENCAPSULATED HERBICIDE COMBINATIONS FOR TOBACCO

Command 3ME Microencapsulated herbicide may be tank mixed with other herbicides registered for use on tobacco to broaden the weed control spectrum compared to the products applied alone. Command 3ME Microencapsulated Herbicide may also be tank mixed with other tobacco herbicides, insecticides, fungicides, and/or nematicides. When applying mixtures of Command 3ME Microencapsulated Herbicide plus other tobacco pesticides, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Water or liquid fertilizer may be used as a carrier for Command 3ME Microencapsulated Herbicide when applied alone, or when tank mixed with other tobacco pesticides unless label use directions/restrictions specifically state otherwise. Refer back to Mixing Instructions for tank mixing information.

TOBACCO REPLANTING INSTRUCTIONS

If initial transplanting fails to produce a uniform stand, tobacco may be replanted in a field treated with Command 3ME Microencapsulated Herbicide alone. When tank mixing with a labeled product, refer to the replant instructions for that product. Where a tank mixture was used refer to all product labels for any additional replant instructions.

Command 3ME Microencapsulated Herbicide APPLIED ALONE RATES AND WEEDS CONTROLLED

Weeds controlled (by rate) are as listed under Soybean for Northern and Southern areas.

TOBACCO ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed under Soybean Rotational Crop Restrictions.

Cover crops may be planted anytime but stand reductions may occur in some areas. **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME treatment.

TOBACCO RESTRICTIONS

- **DO NOT** apply this more than one application per year.
- **DO NOT** use on Tobacco Seedling Beds.
- **DO NOT** replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Restrictions on the Command 3ME Microencapsulated Herbicide label.
- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per application.
- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per year.
- **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment.

VEGETABLE CROPS

Include:

- Stalk and Stem Vegetable Subgroup 22A, except kohlrabi: Agave; aloe vera; asparagus; bamboo, shoots; celtuce; fennel, Florence, fresh leaves and stalk; fern, edible, fiddlehead; kale, sea; palm hearts; prickly pear, pads; prickly pear, Texas, pads; cultivars, varieties, and hybrids of these commodities
- Beans (Dry and Succulent, Except Lima): *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, blackeyed pea and cowpea)
- Brassica head and stem vegetable group 5-16: Broccoli, Brussels sprouts, Cabbage, Chinese Cabbage, Cauliflower, & Cultivars, varieties, and hybrids of these commodities
- Chinese Broccoli
- Cucurbit Vegetables Crop Group 9:
 - o Sub Group 9A: Citron melon; muskmelon; watermelon
 - o Sub Group 9B: Chayote (fruit); Chinese waxgourd; cucumber; gherkin; gourd, edible; *Momordica* spp.; pumpkin; summer squash; winter squash
- Vegetable Soybean (Edamame)
- Kohlrabi
- Lima Beans
- Peas (Succulent)
- Peppers

- Rhubarb
- Sweet Potatoes
- Tuberous and Corm Vegetables Subgroup 1D (except potato): Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen; ginger; leren; sweet potato; tanager; turmeric; yam bean; yam, true

PRODUCT INFORMATION

Before applying Command 3ME on vegetable crops, users, growers and/or applicators must review the VEGETABLE DIRECTIONS found in the TERMS OF SALE OR USE section at the end of this label.

Command 3ME Microencapsulated Herbicide may be utilized as a soil applied treatment prior to weed emergence, for suppression or control of labeled annual grass and broadleaf weeds in the vegetable crops listed above.

Refer to Weeds Controlled table in the Vegetable Crops section for specific weeds.

Make a single herbicide application in a minimum of 10 gallons of water per acre at the rate(s) specified in the individual crop listings hereafter. Apply as a pre-emergent soil applied treatment prior to seeding or transplanting, or after seeding but prior to crop emergence. Place seed, or roots of the transplants, below the chemical barrier when planting.

Command 3ME Microencapsulated Herbicide may be tank mixed with other herbicides registered for use on vegetables to provide a broader spectrum of controlled weeds. When applying mixtures of Command 3ME Microencapsulated Herbicide plus other vegetable herbicides, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Water or liquid fertilizer may be used as a carrier for Command 3ME Microencapsulated Herbicide when applied alone, or when tank mixed with other vegetable herbicides unless use directions on those labels specifically restrict use or provide other limitations.

DIRECTIONS

Place seeds or roots of the transplants below the chemical barrier when planting.

Under drought conditions where rainfall has not sufficiently activated and moved Command 3ME Microencapsulated Herbicide through the soil profile, certain irrigation systems may allow the crop to grow to maturity and not activate Command 3ME Microencapsulated Herbicide. Crop symptomology may occur late in the season under heavy rainfall situations which reactivate Command 3ME Microencapsulated Herbicide that was on the soil surface.

Certain crop varieties may have potential for injury or loss due to use of this herbicide. Consult your agricultural experiment station, county agent, or other qualified crop advisors for information pertaining to crop varieties grown in your area.

PRECAUTIONS: Temporary whitening and/or yellowing of the treated crop may occur.

VEGETABLE ROTATIONAL CROP RESTRICTIONS

Rotate to crops as listed below. Refer to sections headed "Rotational Cropping Precautions" and "Product Information."

NOTE: When using Command 3 ME Microencapsulated Herbicide with other herbicides registered for vegetables always refer to rotational restrictions and directions on the other product's label.

6.4 to 10.7 fluid ounces (0.15 to 0.25 LB. A.I.)	16.2 fluid ounces (0.38 LB. A.I.)	21.3 fluid ounces (0.5 LB. A.I.)
<u>ANYTIME</u> Snap Beans (succulent) Cabbage (all) Cotton* Cucurbits Peas Peppers Rice Soybeans Sweet Potatoes Tobacco Tuberous &Corm Veggies.	<u>ANYTIME</u> Cabbage (transplant) Cotton* Cucurbits Peas Peppers Rice Soybeans Sweet Potatoes Tobacco Tuberous &Corm Veggies.	<u>ANYTIME</u> Cabbage (transplant) Cotton* Peas Peppers Rice Soybeans Sweet Potatoes Tobacco Tuberous &Corm Veggies.
2 MONTHS Lima beans		
<u>9 MONTHS</u> All crops listed above <i>plus:</i> Beans (dry) Corn —Field —Pop —Seed —Sweet Cotton	<u>9 MONTHS</u> All crops listed above <i>plus:</i> Beans (succulent & dry) Cabbage (direct seeded) Corn —Field —Pop —Seed	<u>9 MONTHS</u> All crops listed above <i>plus:</i> Beans (succulent & dry) Cabbage (direct seeded) Corn —Field —Pop —Seed

Peanuts Potatoes Sorghum Sugar Beets Tomatoes (Transplanted) <u>12 MONTHS</u> All crops listed above <i>plus:</i> Tomatoes (direct seeded) Wheat <u>16 MONTHS</u> All crops	—Sweet Cotton Peanuts Potatoes Sorghum Sugar Beets Tomatoes (Transplanted) <u>12 MONTHS</u> All crops listed above <i>plus:</i> Tomatoes (direct seeded) Wheat <u>16 MONTHS</u> All crops	—Sweet Cotton Cucurbits Peanuts Potatoes Sorghum Sugar Beets Tomatoes (Transplanted) <u>12 MONTHS</u> All crops listed above <i>plus:</i> Tomatoes (direct seeded) Wheat <u>16 MONTHS</u> All crops
---	---	--

VEGETABLE ROTATIONAL CROP RESTRICTIONS (CONT'D)

32 fluid ounces (0.75 LB. A.I.)	42.7 to 53.3 fluid ounces (1.0 to 1.25 LB. A.I.)	64 fluid ounces (1.5 LB. A.I.)
<u>ANYTIME</u> Cotton* Peppers Rice Soybeans Squash (winter) Sweet Potatoes Tobacco Tuberous & Corn Veggies. <u>9 MONTHS</u> All crops listed above <i>plus:</i> Those listed under 9 months at 21.3 fluid ounces <i>plus:</i> Cabbage (transplant) Peas Rice <u>12 MONTHS</u> All crops listed above <i>plus:</i> Tomatoes (direct seeded) Wheat <u>16 MONTHS</u> All crops	<u>ANYTIME</u> Cotton* Peppers (42.7 fl oz) Soybeans Sweet Potatoes Tobacco Tuberous & Corn Veggies. <u>9 MONTHS</u> All crops listed above <i>plus:</i> Those listed under 9 months at 21.3 and 32 fluid ounces <i>except:</i> Seed Corn Sweet Corn <u>12 MONTHS</u> All crops listed above <i>plus:</i> Seed Corn Sweet Corn Tomatoes (direct seeded) Wheat <u>16 MONTHS</u> All crops	<u>ANYTIME</u> Sweet Potatoes <u>9 MONTHS</u> Those listed under 9 months at 21.3 and 32 fluid ounces <i>except:</i> Seed Corn Sweet Corn <i>plus:</i> Cotton Peppers Soybeans Sweet Potatoes Tobacco Tuberous & Corn Veggies. <u>12 MONTHS</u> All crops listed above <i>plus:</i> Tomatoes (direct seeded) Seed Corn Sweet Corn

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment. **DO NOT** allow livestock to graze on treated vines or feed treated vines or vine trash to livestock.

*Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

SOIL TEXTURES*:

Coarse (light) soils:	sand, loamy sand, sandy loam
Medium soils	loam, silt, silt loam, sandy clay, sandy clay loam
Fine (heavy) soils:	silty clay, clay loam, silty clay loam, clay

*Select lower to higher rates within the range noted in the individual vegetable crop listings based on lighter to heavier soil types within a textural group.

WEEDS CONTROLLED

RATE/ACRE 6.4 to 16.2 fluid ounce (0.15 - 0.38 LB. A.I.)	RATE/ACRE 16.2 to 21.3 fluid ounces (0.38 - 0.5 LB. A.I.)	RATE/ACRE 21.3 to 64 fluid ounces (0.5 - 1.5 LB. A.I.)
<u>GRASSES</u> Barnyardgrass* Crabgrass* —Large —Smooth Foxtail* —Giant —Green —Robust Goosegrass* Panicum* —Common —Fall <u>BROADLEAVES</u> Galinsoga spp.* Lambsquarters* Spurred Anoda* Velvetleaf*	<u>GRASSES</u> Barnyardgrass* Crabgrass* —Large —Smooth Field Sandbur* Foxtail —Giant —Green —Robust Goosegrass Panicum —Common —Fall Seedling Johnsongrass* <u>BROADLEAVES</u> Galinsoga spp.* Lambsquarters* Prickly Sida* Spurred Anoda Velvetleaf Venice Mallow*	<u>GRASSES</u> Barnyardgrass Crabgrass —Large —Smooth Field Sandbur Foxtail —Giant —Green —Robust —Yellow Goosegrass Panicum —Common —Fall Seedling Johnsongrass** <u>BROADLEAVES</u> Velvetleaf Spurred Anoda Common Ragweed** Galinsoga Jimsonweed** Lambsquarters Prickly Sida Purslane Smartweed** Venice Mallow

*Suppression/Partially Controlled

**Controlled at 1-1.5 Lb. ai/A

VEGETABLE REPLANTING INSTRUCTIONS

If initial seeding or transplants fail to produce the desired stand, vegetables may be replanted in a field treated with Command 3ME Microencapsulated Herbicide alone. **DO NOT** Where a tank mixture is used, refer to the product(s) labels for any additional replant instructions.

RESTRICTIONS

DO NOT retreat field with a second application of Command 3ME Microencapsulated Herbicide.

DO NOT replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Restrictions on the Command 3ME Microencapsulated Herbicide label.

Stalk and Stem Vegetable Subgroup 22A, except Kohlrabi, includes (agave; aloe vera; asparagus; bamboo, shoots; celtuce; fennel, florence, fresh leaves and stalk; fern, edible, fiddlehead; kale, sea; palm hearts; prickly pear, pads; prickly pear, Texas, pads; cultivars, varieties, and hybrids of these commodities)

Command 3ME Microencapsulated Herbicide may be applied as a soil applied treatment prior to weed emergence for suppression or control of annual grass and broadleaf weeds in the stalk and stem subcrop group 22A (not including kohlrabi, included (agave; aloe vera; asparagus; bamboo, shoots; celtnce; fennel, florence, fresh leaves and stalk; fern, edible, fiddlehead; kale, sea; palm hearts; prickly pear, pads; prickly pear, Texas, pads; cultivars, varieties, and hybrids of these commodities). For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

Apply 42.7 fluid ounces of Command 3ME Microencapsulated Herbicide (1.0 lb. ai clomazone) in a minimum of 10 gallons of water per treated acre as a broadcast application to the soil prior to spear emergence. If spears have emerged, make an application after a clean harvest. Cover exposed plants with soil prior to application. Make application at least 14 days prior to harvest.

NOTE: Prior to use, refer above to the Directions sections for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per application.
- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per year.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** make more than one application per year.

BEANS (DRY AND SUCCULENT, EXCEPT LIMA): *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, blackeyed pea and cowpea)

For the control or suppression of annual grasses and broadleaf weeds in beans, apply broadcast at 6.4 to 10.7 fluid ounces per acre (0.15 – 0.25 lb. ai/A) after seeding and prior to emergence. Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 10.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.25 lb. ai/A) per application.
- **DO NOT** apply more than 10.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.25 lb. ai/A) per year.
- **DO NOT** make more than one application to beans (dry and succulent, except lima) per year.
- **DO NOT** apply within 45 days of harvest for beans.

BRASSICA (HEAD AND STEM VEGETABLE GROUP 5-16): Broccoli; Brussels sprouts; Cabbage; Chinese Cabbage (napa); Cauliflower; cultivars, varieties, and hybrids of these commodities

For the control or suppression of annual grasses and broadleaf weeds in head and stem brassica. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

Direct Seeded Cabbage - apply broadcast at 10.7 fluid ounces per acre (0.25 lb. ai/A) after planting prior to emergence.

Transplant Cabbage - apply broadcast at 10.7 to 21.3 fluid ounces per acre (0.25 – 0.5 lb. ai/A) prior to transplanting. Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above.

Transplant Broccoli – apply broadcast at 10.7 fluid ounces per acre (0.25 lb. ai/A) within 48 hours of transplanting broccoli.

All Transplanted Brassica other than Cabbage and Broccoli – Apply broadcast at 10.7 fluid ounces per acre (0.25 lb. ai/A) within 72 hours of transplanting.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** make more than one application to brassica (head and stem vegetable group 5-16) per year.
- **DO NOT** apply within 45 days of harvest for cabbage.

- Direct seeded cabbage and transplant broccoli:
 - **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per application for direct seeded cabbage and transplant broccoli.
 - **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per year for direct seeded cabbage and transplant broccoli.
- Transplant cabbage:
 - **DO NOT** apply more than 21.3 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.5 lb. ai/A) per application for transplant cabbage.
 - **DO NOT** apply more than 21.3 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.5 lb. ai/A) per year for transplant cabbage.
- All transplanted brassica other than cabbage or broccoli:
 - **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per application.
 - **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per year.

CHINESE BROCCOLI (GAI LAN)

For the control or suppression of annual grasses and broadleaf weeds in Chinese broccoli, broadcast apply 10.7 fluid ounces (0.25 lb. ai) per acre Command 3ME Microencapsulated Herbicide within 48 hours of transplanting and prior to weed emergence. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

Transplant Chinese Broccoli – apply broadcast at 10.7 fluid ounces per acre (0.25 lb. ai/A) within 48 hours of transplanting Chinese broccoli.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per application.
- **DO NOT** apply more than 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per year.
- **DO NOT** make more than one application to Chinese broccoli per year.
- **DO NOT** apply within 45 days of harvest of Chinese broccoli.

CUCURBIT VEGETABLES CROP GROUP 9:

- Sub Group 9A: Citron melon; muskmelon; watermelon
- Sub Group 9B: Chayote (fruit); Chinese waxgourd; cucumber; gherkin; gourd, edible; *Momordica* spp.; pumpkin; summer squash; winter squash

SUB GROUP 9A: MELONS (CITRON MELON; MUSKMELONS AND WATERMELONS)

Muskmelon: hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon)

For the control or suppression of annual grasses and broadleaf weeds in sub group 9A melons: cantaloupes, citron melons; muskmelons and watermelons, apply broadcast at 6.4 to 10.7 fluid ounces (0.15 - 0.25 lb. ai) per acre. Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

For direct seeded melons: Make one broadcast application after seeding but prior to melon emergence.

For transplanted melons: Make one broadcast application before transplanting and ensure to place the roots of the transplant below the chemical barrier when planting.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 10.7 fluid ounces Command 3ME Microencapsulated Herbicide (0.25 lb. ai clomazone) per acre per year.

- **DO NOT** apply more than 10.7 fluid ounces Command 3ME Microencapsulated Herbicide (0.25 lb. ai clomazone) per acre per application.
- **DO NOT** make more than one application to melons per year.

SUB GROUP 9B: CHAYOTE (FRUIT); CHINESE WAXGOURD; CUCUMBER; GHERKIN; GOURD, EDIBLE; *MOMORDICA* SPP.; PUMPKIN; SUMMER SQUASH; WINTER SQUASH

CUCUMBERS

For the control or suppression of annual grasses and broadleaf weeds in cucumber, apply broadcast at 6.4 to 16.2 fluid ounces (0.15 – 0.38 lb. ai/A) per acre in a minimum spray volume of 10 gallons of water per treated acre. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above. Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above.

For direct seeded cucumber: Make one broadcast application after seeding but prior to cucumber emergence.

For transplanted cucumber: Make one broadcast application before transplanting, and ensure to place the roots of the transplant below the chemical barrier when planting.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 16.2 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.38 lb. ai/A) per application.
- **DO NOT** apply more than 16.2 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.38 lb. ai/A) per year.
- **DO NOT** make more than one application to cucumber per year.
- **DO NOT** apply within 30 days of harvest of cucumbers.

SUMMER SQUASH

Squash, summer (*Cucurbita pepo* var. *meloepo*) (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini).

For the control or suppression of annual grasses and broadleaf weeds in summer squash (refer to Weeds Controlled table in the Vegetable Crops Section).

For summer squash, apply broadcast, prior to seeding or transplanting, at 10.7 to 21.3 fluid ounces per acre (0.25 – 0.5 lb. ai/A). Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above.

OR

Apply as a pre-emergent (to weeds) soil surface treatment before, or just prior to transplanting, to row middles between plastic covered beds. Use rates as noted above.

NOTE: Prior to use, refer to the Directions section for vegetable crops above.

RESTRICTIONS

- **DO NOT** make more than one application to squash per year.
- **DO NOT** apply to soil that will be covered with plastic mulch.
- Summer squash (*Cucurbita pepo* var. *meloepo*) (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini):
- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per application for summer squash.
- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per year for summer squash.
- **DO NOT** apply within 30 days of harvesting summer squash.

WINTER SQUASH and PROCESSING PUMPKINS

Squash, winter (*Cucurbita maxima*; *C. moschata*) (includes butternut squash, calabaza, hubbard squash); (*C. mixta*; *C. pepo*) (includes acorn squash, spaghetti squash):

For the control or suppression of annual grasses and broadleaf weeds in winter squash and processing pumpkins (refer to Weeds Controlled table in the Vegetable Crops Section).

For winter squash and processing pumpkins, apply broadcast, prior to seeding or transplanting, at 10.7 to 32 fluid ounces per acre (0.25 – 0.75 lb. ai/A). Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above.

OR

Apply as a pre-emergent (to weeds) soil surface treatment before, or just prior to transplanting, to row middles between plastic covered beds. Use rates as noted above.

NOTE: Prior to use, refer to the Directions section for vegetable crops above.

RESTRICTIONS

- **DO NOT** make more than one application to squash or processing pumpkins per year.
- **DO NOT** apply to soil that will be covered with plastic mulch.
- **DO NOT** use on the following cultivars or varieties as unacceptable whitening of the fruit may occur: NK530, NK580, Turks Turban, Golden Delicious, all Banana types, and all other Cucurbita maxima types that have a pink or burnt orange coloration at harvest.
- **DO NOT** use on Jack-o-lantern pumpkins.
- Winter squash (*Cucurbita maxima*; *C. moschata*) (includes butternut squash, calabaza, hubbard squash); (*C. mixta*; *C. pepo*) (includes acorn squash, spaghetti squash):
- **DO NOT** apply more than 32 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.75 lb. ai/A) per application for winter squash and processing pumpkins.
- **DO NOT** apply more than 32 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.75 lb. ai/A) per year for winter squash and processing pumpkins.
- **DO NOT** apply within 45 days of harvesting winter squash and processing pumpkins.

EDAMAME (Vegetable Soybean)

Command 3ME Microencapsulated Herbicide may be applied as a soil applied treatment prior to weed emergence and after seeding for suppression or control of annual grass and broadleaf weeds in edamame. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

Apply 21.3 fluid ounces of Command 3ME (0.5 lb. ai clomazone) in a minimum of 10 gallons of water per treated acre as a broadcast application to the soil prior to weed emergence and after seeding.

RESTRICTIONS

- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per application.
- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per year.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** make more than one application to edamame per year.

KOHLRABI

For the control or suppression of annual grasses and broadleaf weeds in kohlrabi, apply Command 3ME Microencapsulated Herbicide broadcast. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

Direct Seeded kohlrabi - apply broadcast at 10.7 fluid ounces per acre (0.25 lb. ai/A) after seeding but prior to emergence.

Transplant kohlrabi - apply broadcast at 10.7 to 21.3 fluid ounces per acre (0.25 to 0.5 lb. ai/A) prior to transplanting. Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- Direct seeded kohlrabi:
 - **DO NOT** apply more than 10.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per application for direct seeded kohlrabi.
 - **DO NOT** apply more than 10.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide per acre (0.25 lb. ai/A) per year for direct seeded kohlrabi.
- Transplant kohlrabi:
 - **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per acre per application for transplant kohlrabi.
 - **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide per acre (0.5 lb. ai/A) per year for transplant kohlrabi.
- **DO NOT** make more than one application to kohlrabi per year.
- **DO NOT** apply within 45 days of harvest for kohlrabi.

LIMA BEANS

For the suppression of annual grasses and broadleaf weeds in lima beans, apply Command 3ME Microencapsulated Herbicide broadcast at 4 to 6 fluid ounces per acre as a soil applied treatment prior to weed emergence and after seeding. This use direction is only for suppression of early season grass and certain broadleaf weeds to provide additional time for follow up cultivation and application of other labeled post emergent herbicides. Observe a 60-day rotational period when planting to lima beans.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 6 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.14 lb. ai/A) per application.
- **DO NOT** apply more than 10.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.25 lb. ai/A) per year.
- **DO NOT** make more than two applications to lima beans per year.
- Allow 14 days between re-treatment applications.

PEAS (SUCCULENT)

pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea); and pigeon pea.

pea (*Pisum* spp.) (includes English pea, garden pea, green pea); and pigeon pea.

For the control or suppression of annual grasses and broadleaf weeds in succulent pea, apply broadcast at 21.3 fluid ounces per acre (0.5 lb. ai/A) after seeding and prior to weed emergence. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per application.
- **DO NOT** apply more than 21.3 fluid ounces of Command 3ME (0.5 lb. ai) per acre per year.
- **DO NOT** make more than one application to succulent peas per year.

PEPPERS (Bell and NonBell)

For the control or suppression of annual grasses and broadleaf weeds in Bell peppers and NonBell peppers, apply broadcast at 10.7 to 42.7 fluid ounces per acre (0.25 – 1 lb. ai/A). Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT use on banana peppers.**
- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per year.
- **DO NOT** apply more than 42.7 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.0 lb. ai/A) per application.
- **DO NOT** make more than one application to peppers per year.

RHUBARB

For the control or suppression of annual grasses and broadleaf weeds in rhubarb. Apply broadcast at 64 fluid ounces per acre (1.5 lb. ai/A) to dormant rhubarb prior to leaf emergence. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

NOTE: Prior to use, refer above to the Directions section for vegetable crops

RESTRICTIONS

- **DO NOT** apply more than 64 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.5 lb. ai/A) per application.
- **DO NOT** apply more than 64 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.5 lb. ai/A) per year.
- **DO NOT** make more than one application to rhubarb per year.

SWEET POTATOES

For the control of annual grasses and broadleaf weeds in sweet potatoes, apply broadcast at 21.3 to 64 fluid ounces per acre (0.5 – 1.5 lb. ai/A). Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

OR

Apply as a single application post transplant to the crop and prior to weed emergence at a maximum of 64 fluid ounces per acre (1.5 lb. ai/A).

NOTE: Prior to use, refer above to the Directions section for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 64 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.5 lb. ai/A) per application.
- **DO NOT** apply more than 64 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.5 lb. ai/A) per year.
- **DO NOT** make more than one application to sweet potatoes per year.
- **DO NOT** apply within 95 days of harvest (or 125 days if more than 53.3 fluid ounces per acre (1.25 lb. ai/A) was applied) for sweet potatoes.

CROP SUBGROUP 1D. TUBEROUS AND CORM VEGETABLES (EXCEPT POTATO) & (ARRACACHA, CASSAVA, TANIER, YAMS) (FOR SWEET POTATO REFER TO DIRECTIONS ABOVE)

For the control of annual grasses and broadleaf weeds in arracacha, cassava, tanier and yams, apply broadcast at 32 to 53.3 fluid ounces per acre (0.75 – 1.25 lb. ai/A). Use the lower specified rate on coarse soils and the higher specified rate on fine soils. Refer to the soil texture chart found above. For a list of weeds controlled, refer to the Weeds Controlled table in the Vegetable Crops section above.

RESTRICTIONS

- DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per application.
- DO NOT** apply more than 53.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (1.25 lb. ai/A) per year.
- DO NOT** make more than one application to tuberous and corm vegetables per year.
- DO NOT** apply within 125 days of harvest.

OIL CROPS

Rapeseed (canola varieties only)

Command 3ME Microencapsulated Herbicide may be applied as a soil applied broadcast treatment prior to seeding and weed emergence for control or suppression of annual grass and broadleaf weeds in winter or spring canola and rapeseed, cultivars, varieties, and/or hybrids of these.

Command 3ME Microencapsulated Herbicide is to be used as part of a system approach for the control of cleavers (Galium aparine) in all herbicide tolerant canola varieties. Application of a post-emergence herbicide that is effective on cleavers, e.g. a herbicide to which the canola variety is tolerant, is required for season-long control. When used on non-herbicide tolerant canola varieties, Command 3ME Microencapsulated Herbicide will provide some early season weed suppression only.

Apply 3.4 - 4.5 fluid ounces of Command 3ME Microencapsulated Herbicide (0.08 – 0.11 lb. ai clomazone) in a minimum spray volume of 10 gallons of water per treated acre.

Application of less than 4.5 fl. oz. will suppress cleavers and can be used if cleaver populations are light.

Command 3ME Microencapsulated Herbicide may be tank mixed with other labeled herbicides such as Aim EC herbicide and glyphosate.

PRECAUTION: Temporary whitening and/or yellowing of the crop may occur when emerging from the treated soil. Susceptible weed species may emerge from the soil but are devoid of pigmentation and plant death occurs in a short period of time.

NOTE: Prior to use, refer above to the Directions sections for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 4.5 fluid ounces Command 3ME Microencapsulated Herbicide (0.11 lb. ai clomazone) per acre per application.
- **DO NOT** make more than one application to canola and rapeseed per year.
- **DO NOT** apply more than 4.5 fluid ounces Command 3ME Microencapsulated Herbicide (0.11 lb. ai clomazone) per acre per year.
- **DO NOT** apply within 90 days of harvest of canola and rapeseed cultivars, varieties, and/or hybrids of these.
- **DO NOT** apply to sandy soils.

PEPPERMINT and SPEARMINT TOPS

Before applying Command 3ME on peppermint and spearmint, users, growers and/or applicators must review the **VEGETABLE DIRECTIONS** found in the **TERMS OF SALE OR USE** section at the end of this label.

Command 3ME Microencapsulated Herbicide may be utilized as a pre-emergent treatment for the control of annual grass and broadleaf weeds in peppermint and spearmint. Command 3ME Microencapsulated Herbicide may be applied at the rate of 21.3 fluid ounces per acre (0.5 lb. ai/A) to the soil surface prior to the emergence of weeds and the initiation of new growth of peppermint and spearmint plants.

Weeds Controlled

GRASSES	BROADLEAVES
Barnyardgrass*	Galinsoga spp.*
Crabgrass	Lambsquarters*
—Large*	Spurred anoda*
—Smooth*	Velvetleaf
Foxtail	Venice mallow*
—Giant*	
—Green*	
—Robust*	
Goosegrass*	
Panicum	
—Common*	
—Fall*	

*Suppression/Partially Controlled

RESTRICTIONS

- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per application.
- **DO NOT** apply more than 21.3 fluid ounces per acre Command 3ME Microencapsulated Herbicide (0.5 lb. ai/A) per year.
- **DO NOT** make more than one application to peppermint and spearmint per year.
- **DO NOT** apply within 84 days of harvest.

Rotational Crop Restrictions

Rotate to crops as listed below. Refer to sections headed “Rotational Cropping Precautions” and “Product Information.”

NOTE: When using Command 3ME Microencapsulated Herbicide with other herbicides registered for peppermint and spearmint, always refer to rotational restrictions and directions on the other product’s label.

ANYTIME	9 MONTHS	12 MONTHS
Cabbage (transplant)	All crops listed under ANYTIME <i>plus:</i>	All crops listed under 9 MONTHS <i>plus:</i>
Cotton*	Beans (succulent & dry)	Tomatoes (direct seeded)
Peas	Cabbage (direct seeded)	Wheat
Peppers	Corn	
Rice	—Field	<u>16 MONTHS</u>
Soybeans	—Pop	All crops
Squash	—Seed	
Sweet Potatoes	—Sweet	
Tobacco	Cotton	
Tuberous & Corm Veggies.	Curcubits	
	Peanuts	
	Potatoes	
	Sorghum	
	Sugar Beets	
	Tomatoes (Transplanted)	

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. **DO NOT** graze or harvest for food or feed cover crops planted less than 9 months after Command 3ME Microencapsulated Herbicide treatment.

* Refer to the section “Requirements for Planting Time Applications” and “Replanting Instructions” in the use directions for cotton.

HERB SUBGROUP 19A

(Angelica; balm; basil; borage; burnet; camomile; catnip; chervil (dried); chive; chive, Chinese, clary; coriander (leaf); costmary; culantro (leaf); curry (leaf); dillweed; horehound; hyssop; lavender; lemongrass; lovage (leaf); marigold; marjoram (*Origanum* spp.); nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage; savory, summer and winter; sweet bay; tansy; tarragon; thyme; wintergreen; woodruff; and wormwood)

Before applying Command 3ME on crops in Herb Subgroup 19A: angelica; balm; basil; borage; burnet; camomile; catnip; chervil (dried); chive; chive, Chinese, clary; coriander/cilantro (leaf); costmary; culantro (leaf); curry (leaf); dillweed; horehound; hyssop; lavender; lemongrass; lovage (leaf); marigold; marjoram (*Origanum* spp.); nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage; savory, summer and winter; sweet bay; tansy; tarragon; thyme; wintergreen; woodruff; and wormwood, **users, growers and/or applicators must review the VEGETABLE DIRECTIONS found in the TERMS OF SALE OR USE section at the end of this label.**

Command 3ME Microencapsulated Herbicide may be applied as a soil applied broadcast treatment after seeding and prior to crop and weed emergence for control or suppression of annual grass and broadleaf weeds in crops in Herb Subgroup 19A: angelica; balm; basil; borage; burnet; camomile; catnip; chervil (dried); chive; chive, Chinese, clary; coriander/cilantro (leaf); costmary; culantro (leaf); curry (leaf); dillweed; horehound; hyssop; lavender; lemongrass; lovage (leaf); marigold; marjoram (*Origanum* spp.); nasturtium; parsley (dried); pennyroyal; rosemary; rue; wintergreen; woodruff; and wormwood. For a list of weeds controlled or suppressed, refer to the Weeds Controlled table in the Vegetable Crops section on the label.

Apply 10.7 fluid ounces of Command 3ME Microencapsulated Herbicide (0.25 lb. ai clomazone) in a minimum spray volume of 10 gallons of water per treated acre. Make application at least 50 days prior to harvest.

NOTE: Prior to use, refer above to the Directions sections for vegetable crops.

RESTRICTIONS

- **DO NOT** apply more than 10.7 fluid ounces Command 3ME Microencapsulated Herbicide (0.25 lb. ai clomazone) per acre per application.
- **DO NOT** make more than one application to crops in Herb Subgroup 19A: angelica; balm; basil; borage; burnet; camomile; catnip; chervil (dried); chive; chive, Chinese, clary; coriander/cilantro (leaf); costmary; culantro (leaf); curry (leaf); dillweed; horehound; hyssop; lavender; lemongrass; lovage (leaf); marigold; marjoram (*Origanum* spp.); nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage; savory, summer and winter; sweet bay; tansy; tarragon; thyme; wintergreen; woodruff; and wormwood per year.
- **DO NOT** apply more than 10.7 fluid ounces Command 3ME Microencapsulated Herbicide (0.25 lb. ai clomazone) per acre per year.
- **DO NOT** apply within 50 days of harvest.

Dealers Must Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: To the extent consistent with applicable law, FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, except as so warranted the product is sold as is. To the extent consistent with applicable law, buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specified and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's instructions for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: To the extent consistent with applicable law, buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Vegetable Directions: FMC Corporation intends that this product be distributed only to end users and/or growers (and/or applicators acting on behalf of growers) who agree to the terms and conditions below, and understand the directions regarding use Command 3ME Microencapsulated Herbicide in pre-emergence ground treatments on beans (dry and succulent, except lima), Brassica head and stem vegetable group 5-16, Chinese broccoli, Cucurbit Vegetables Crop Group 9, edamame (vegetable soybean), kohlrabi, HERBS Subgroup 19A, lima beans, peas (Succulent), peppermint tops, peppers, pumpkins, Rapeseed (canola varieties only), rhubarb, spearmint tops, Stalk and Stem Vegetable Subgroup 22A, Summer and Winter squash, sweet potatoes, Tuberous and Corm Vegetables Subgroup 1D.

If such directions are unacceptable, FMC requests the return of Command 3ME Microencapsulated Herbicide at once unopened. FMC considers the user to have read and understood such directions upon use of this product.

This product when used as a pre-emergence ground treatment on beans (dry and succulent, except lima), Brassica head and stem vegetable group 5-16, Chinese broccoli, Cucurbit Vegetables Crop Group 9, edamame (vegetable soybean), kohlrabi, HERBS Subgroup 19A, lima beans, peas (Succulent), peppermint tops, peppers, pumpkins, Rapeseed (canola varieties only), rhubarb, spearmint tops, Stalk and Stem Vegetable Subgroup 22A, Summer and Winter squash, sweet potatoes, Tuberous and Corm Vegetables Subgroup 1D may lead to crop injury, loss, or damage. FMC advises that the user and/or grower test this product in order to determine its suitability for such intended use. FMC makes this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use this product must be made by each individual user and/or grower on the basis of possible crop injury from Command 3ME Microencapsulated Herbicide, the severity of weed infestations, the cost of alternative weed controls, and other factors.

COMMAND and FMC—Trademarks of FMC Corporation
©2021 FMC Corporation. All rights reserved
Label Code D-4477 062521
Replaces Label Code D-4121 010719