



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

September 16, 2010

Christopher Davis Registration Manager 1735 Market Street Philadelphia, PA 19103

Subject:

Notification - Adding Trademarked Tag Line Logo

EPA Registration Number: 279-3168 Date of Submission: August 16, 2010

Dear Mr. Davis:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration (PRN) 98-10 dated August 16, 2010, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact BeWanda Alexander at (703) 305-7460 or e-mail <u>alexander.bewanda@epa.gov</u>.

Sincerely,

Richard Gebken Product Manager 10

Insecticide Branch

Registration Division (7505P)

Branda Weyandor for

Enclosures:

Copy of Label Stamped "Notification"

Please read instructions	on reverse before comp	Form Approve	, <u>MB No. 2070-0060</u>	. Approvel expires 2-28-95
\$EPA	United States Environmental Protection Agency Washington, DC 20460		Registration Amendment Other	OPP Identifier Number
	Application for Pestici	de - Section	1	

, , , , , , , , , , , , , , , , , , ,	✓ Other				
Application for	Pesticide - Section I				
. Company/Product Number 279-3168	2. EPA Product Manager Gebken 3. Proposed Classification				
R. Company/Product (Name) Talstar-PL-Granular-Insecticide	PM# 13				
. Name and Address of Applicant (Include ZIP Code)	6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3)				
FMC Corporation 1735 Market Street Philadelphia, PA 19103 Check if this is a new address	(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 279-3168 Product Name Talstar PL Granular Insecticide				
	ction - II				
labeling or the confidential statement of formula for this product. I understar EPA. I further understand that if this notification is not consistent with the te	PA regulations at 40 CFR 152.46, and no other changes have been made to the not that it is a violation of U.S.C. Sec.1001 to willfully make any false statement to times of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of				
FIFRA and I may be subject to enforcement and penalties under sections 12	2 and 14 of FIFRA. Ction - III				
. Material This Product Will Be Packaged In:					
Yes Yes No. per If "Yes" No. per If "Yes"	Yes No No No. per age wgt container 2. Type of Container Metal Plastic Glass Paper Other (Specify)				
. Location of Net Contents Information 4. Size(s) Retail Conte	siner 5. Location of Label Directions				
Label Container					
	tion - IV				
. Contact Point	Talleptions No. (include Area Code) ration Manager				
Certification I certify that the statements I have made on this form and all attact I acknowledge that any knowlingly false or misleading statement in both under applicable law. 3. Title Registra					
. Typed Name 5. Date Christopher Davis	8/16/10				

FMC Agricultural Products

FMC Corporation 1735 Market Street Philadelphia, PA 19103

215.299.6000 Phone 215.299.6468 Fax

www.fmc.com

August 16, 2010

Mr. Richard Gebken, PM-13 United States Environmental Protection Agency Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202

Dear Mr. Gebken:

Subject:

Notification of other revisions per PR notice 98-10 (II.N)

Talstar PL Granular Insecticide (EPA Reg. Number: 279-3168)

FMC is notifying the Agency that we have revised Talstar PL Granular Insecticide's logo to include our trademarked tag line. Enclosed is 1 copy of the label for you records.

Please do not hesitate to call me if you have any other questions. My phone number is (215) 299-6334.

Sincerely,


Christopher Davis Registration Manager

Enclosures:

This letter

EPA Form 8570-1

1 Copy of Talstar PL Granular Insecticide label



Talstar PL Granular

Insecticide

NOTIFICATION

SEP 1-6 2010



NOTIFICATION SEP 1 6 2010

FIRST AID

For use to control Ants (including Imported Fire Ants), Mole Crickets and other Insect pests on lawns in landscaped areas and perimeters around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

EPA Reg. No. 279-3168	EPA Est. 279-IL-1
Active Ingredient:	By Wt.
Bifenthrin*	
Other Ingredients	
	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum. U.S. Patent No. 4,238,505

KEEP OUT OF REACH OF CHILDREN **CAUTION**

See other panels for additional precautionary information

See back panel pamphlet for additional precautionary information and complete directions for use.

Net Contents



08-13-10

FMC Corporation Agricultural Products Group 1735 Market Street Philadelphia PA 19103

If on skin or clothing

- · Take off contaminated clothing.
- · Rinse skin immediately with plenty of water for 15- 20 minutes.
- · Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.

NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested. milk, cream and other digestible fats and oils may increase absorption and so should be avoided.

For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362).

[Insert photo of house with beautiful lawn]

PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

Caution

Harmful if absorbed through skin. Avoid contact with skin, eyes or

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale of other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around 'private homes, apartment buildings, condominiums, non-agricultural outbuildings, noncommercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre bet application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas creathletic fields).

Do not broadcast apply more than 0.4 lbs. A.I. (200 lbs. Talstar PL Granular) per Acre per year.

Not for use on golf courses and farms, nurseries, in commercial greenhouses or on grass grown for seed.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Paper and plastic bags: Non-refillable. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, or by burning. Do not burn unless allowed by state or local authorities. If burned, stay out of smoke.

LAWN AND PERIMETER APPLICATIONS

Lawn Application: Broadcast Talstar® PL granular insecticide with suitable application equipment to ensure uniform coverage over the treatment area.

APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, Talstar PL may be applied at up to 0.4 lb Al/A to control each of the pests listed in this Table. However, for residential lawn use, do not apply more than 0.2 lbs. A.I. per Acre per application.

Pest	Active ingredient lbs. per acre	Application Rate Talstar PL		
Armyworms¹ Cutworms¹ Sod Webworm¹	0.1 lbs. Al/A	50 lbs. per acre	1.15 lbs. per 1000 sq. ft.	
Annual Bluegrass Weevil (Hyperodes) (Adult) ² Billbugs (Adult) ³ Black Turfgrass Ataenius (Adult) ⁴ Leafhoppers Mealybugs	0.1 - 0.2	50 - 100 lbs.	1.15 - 2.3 lbs.	
	lbs. Al/A	per acre	per 1000 sq. ft.	
Ants Centipedes Chinch Bugs ⁵ European Crane Files (larvae) ⁶ Fleas (Larvae) ⁷ Fleas (Adult) Imported Fire Ants (Adult) ⁸ Imported Fire Ants (Mounds) ⁸ Millipedes Mole Cricket (Adult) ⁸ Mole Cricket (Nymph) ¹⁰ Pillbugs Scorpions ¹¹ Sowbugs Ticks ¹²	0.2 - 0.4	100 - 200 lbs.	2.3 - 4.6 lbs.	
	lbs. Al/A	per acre	per 1000 sq. ft.	

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Talstar PL Granular if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Comments

¹Armyworms, Cutworms, and Sod Webworms: To ensure optimum control, irrigate the treated area with up to 0.1 inches of water immediately after application to activate (release from the granule) the insecticide.

²Annual Bluegrass Weevil (*Hyperodes*) adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood is. in. full bloom... Consult your State-Cooperative Extension Service for more specific information regarding application timing.

³Billbug adults: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

⁴Black Turfgrass Ataenius adults: Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).

⁵Chinch Bugs: Chinch bugs infest the base of grass plants and are often found in the thatch layer. Irrigate the treated area with up to 0.25 inches of water immediately after application to activate (release from the granule) the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher applications rates may be required to control populations that contain both nymphs and adults during the summer.

⁶European Crane Flies: Treatments should be made to control early to midseason larvae (approximately August – February) as they feed on plant crowns. Treatments made to late-season larvae (approximately March - April) may only provide suppression. Consult your local extensions agent for specific recommendations for your area.

⁷Flea larvae: Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release from the granule) the insecticide.

⁸Imported Fire ants: Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound treatments that will control existing colonies. If the soil is not moist, then it is important to irrigate before application. Broadcast treatments should apply 0.2 to 0.4 lb Al/A. Mounds should be treated by applying ½ cup of Talstar PL Granular per mound and then drenching the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. Treat three feet out around the mound. For best results, apply in cool weather (65 - 80° F) or in early morning or late evening hours. This treatment will kill the queen within 24 hours of application. Application of 125 lbs./acre of Talstar PL granular insecticide will provide six months residual activity for control of foraging Imported Fire Ants and newly mated Fly-In Queens.

⁹Mole Cricket adults: Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

¹⁰Mole Cricket nymphs: Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young hymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application sets, and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then its important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

¹¹Scorpions: To ensure optimum control, treat the Eudring perimeter at dusk to prevent outdoor scorpions from entering the building. Scorpions reside hidden in cracks and voids during the day and are active at night at temperatures above 77°F/25°C. Resting areas can be identified at night using a blacklight (UV bulb) as scorpions will fluoresce. These areas should also be treated. Prior to treatment, to increase treatment efficacy, remove trash, debris, or firewood that scorpions may use as resting sites.

12Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted fever): Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Do not allow public use of treated areas during application.

Deer ticks (Ixodes sp.) have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in mid to late-apring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These-ticks commonly-congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from midspring to early fall to control American dog tick larvae, nymphs and adults.

Perimeter Application: Apply to ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Applications: Apply to soil areas, ornamental gardens, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas

Applic. Rate:		Pounds of Talstar PL Granular for Perimeter Applications Linear Distance (Feet) Around Structure							
Pounds Per 1,000 sq. ft.	Band Width	100	150	200	250	300	400	500	1,000
2.3	5 feet	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10 feet	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5 feet	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10 feet	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5 feet	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10 feet	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5 feet	2.6	3.8	5.0	6.0	7.2	9.6		23.4
4.6	10 feet	6.0	8.4	10.6	13.0	15.2	19.8		47.4

Note: The amounts of Talstar PL Granular listed in the above table are based on the total area being treated. This area is determined by multiplying the linear distance by the band width and then adding to that value the area of a circle whose radius is equal to the band width (to account for the area treated at the four corners of a rectangular structure).

Fire ant mound treatment: Apply 1/2 cup of Talstar® PL Granular Insecticide per mound and then drench the mound with 1 to 2 gallons of water. The mound should be drenched with sufficient force to break its apex and allow the insecticide to flow into the ant tunnels. Treat three feet out around the mound. For best results, apply in cool weather (65 to 80°F) or in early morning or late evening hours. Treat new mounds as they appear. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS

(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

Broadcast Spot Applications: For the control of actively foraging ants (including Imported Fire Ants) broadcast uniformly 2.3 to 4.6 lbs. Talstar PL granular insecticide per 1000 sq. ft. (0.2 to 0.4 lbs. a.i./Acre). See comments section under **Lawn Applications** for additional rate recommendations.

Pad Mounted Electrical Sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 1.0 to 1.5 Tablespoons (0.06 to 0.09 lbs.) of Talstar PL granular insecticide per 4 sq. ft. of mounting pad. Where existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application. Food utensils such as tablespoons must not be used for food purposes after use in measuring pesticides.

Note: Electrical transformers appear to be more attractive to ants during the cooler months. For best results treat in the Fall or early Spring.

Insert spreader setting table here

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control or FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Talstar and FMC – Registered trucemarks of FMC Corporation. Verge - trademark of Oil-Dri Corporation of America