

Code 1719




0.05 LAWN GRANULAR INSECTICIDE

Lawn Insect Control

Treats 5,000 sq. ft.

ACCEPTED
AUG 13 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 279-3172



KEEP OUT OF REACH OF CHILDREN
CAUTION

See other panels for additional precautionary information.

Controls major home lawn insect problems.

- Kills insects that cause your lawn to thin and turn brown.
• Kills fleas and ticks before they get on your pets.


For Residential Lawn Use Only

EPA Reg. No. 279-3172

EPA Est. 279-

Table with 2 columns: Active Ingredient, Inert Ingredients, and By Wt. (0.05%, 99.95%, 100.00%)

*Cis isomers 97% minimum, trans isomers 3% maximum. U.S. Patent No. 4,238,505



FMC Corporation
Agricultural Products Group
Philadelphia PA 19103

Net Weight 11.5 pounds

PRECAUTIONARY STATEMENTS
Hazards to Humans (and Domestic Animals)
Caution
Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.
Environmental Hazards
This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

How and when to apply

Do apply when insect activity or damage is first observed in the lawn (lawn thins and turns brown or moths fly from the grass when you mow).

Do water lawn after application to wash granules from the lawn foliage.

Don't treat just the browned areas of lawn. To prevent further damage, treat the entire lawn.

Don't use any other lawn control product for at least one week after application.

In New York State, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Talstar 0.05 Lawn Granular Insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

For fire ant control, sprinkle granules over fire ant mound at the rate of 2 tablespoons* per 3 square feet (1.5 ft. X 2 ft. area). For best results, 1) apply in cool weather, 65° to 80°F, or in early morning or late evening hours, 2) avoid disturbing ants, 3) treat new mounds as they appear and, 4) thoroughly water the treated area immediately after application.

For grub control, apply Scotts® GrubEx® anytime from April to mid-August. A single application provides season-long grub control.

*Note to EPA reviewer: For product with a different bulk density, this will change.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. For rigid containers: After partial use replace lids and close tightly. For bags: After partial use close tightly.

Pesticide Container Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Spreader Tips


For best results, use a Scotts spreader

	Spreader Settings covers 5,000 sq. ft.	Directions
Scotts Drop Spreaders	X	Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.
Scotts EasyGreen® Rotary	XX	Slide shut-off lever to the right; application automatically starts and stops when you do. For complete coverage make each pass about 30 inches from the previous one so some overlapping occurs.
Scotts SpeedyGreen® Rotary	X	Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs. Avoid applying this product on flower, vegetable and shrub plantings.

Additional Spreader Settings

- EZ Drop, Republic (#530)X
- EZ Broadcast, Republic (#540)X
- Red Devil Drop (#FH-22)X
- Red Devil Rotary (#SB-50)X
- Quaker Drop (Model # 10850)X
- Quaker Rotary (Model #10301)X

Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.



What to expect

Talstar Lawn Insect Control kills these active insects: ants (fire ants*, argentine, carpenter, odorous, pyramid, pavement, Pharoah (sugar), red harvester), armyworms, billbugs, centipedes, chiggers, chinch bugs, crickets, cutworms, earwigs, fleas, grasshoppers, millipedes, scorpions, sod webworms (lawn moth), sow bugs, spiders, ticks (including brown dog ticks).

*aids in control

Questions?

[PhoneArt]

Call Scotts' toll-free Consumer Hotline: 1-800-543-TURF

[Computer Art]

Visit Scotts "Lawn Success Made Easy" website at www.scottslawn.com.

Scotts No-Quibble Guarantee™

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

Scotts®...the lawn people
Marysville, Ohio 43041
896-594

©1997, The Scotts Company. World rights reserved.

Talstar® 0.05 Lawn Granular Insecticide is a high-performance product specially designed to be used in very low concentrations and is:

- Virtually Odor-free
- Fast Acting
- Easy-to-Use
- Broad Spectrum
- Kills Insects, Spiders and Ticks
- Not harmful to grasses
- Treats up to 5,000 sq.ft.
- Soil penetrating formula
- Kill [insects/pests] above and/& below the [lawn]surface
- Kills insects in/& under the lawn
- Guaranteed kill or your money back

Talstar is designed for use as a ready-to-use granular insecticide to control various insects and spiders on lawns. Please read the following "Directions for Use" carefully and fully before applying this product.

Surface Feeding Lawn Insects

Use Talstar for the control of ants, including Carpenter, Red Harvester, Pavement, Odorous Pyramid, Argentine, Pharoah (Sugar) and Fire Ants*, Foraging Fire Ants, Armyworms, Centipedes, Chiggers, Chinch Bugs, Crickets, Cutworms, Earwigs, Grasshoppers, Fleas, Millipedes, Scorpions, Sod Webworms (Lawn Moth), Sow Bugs, Spiders, Ticks (including Brown Dog Ticks) and Weevils (including Hyperodes). Uniformly apply 2.3 pounds over a 1,000 sq. ft. area of lawn using any hand or power operated gravity or rotary type spreader designed for the application of granular materials. A 11.5 pound bag will treat a 5,000 sq. ft. area of lawn. Best results are obtained if the treated area is thoroughly watered immediately after application. Treat when insects first appear and retreat as necessary.

*Aids in control.

Mole Crickets

Uniformly apply 4.6 pounds over a 1,000 sq. ft. area of lawn. For best results: treat in late spring, and water the treated area thoroughly immediately after application.

For Fire Ant Mounds

To control Fire ants, uniformly apply 2 tablespoons or 6 teaspoons** of Talstar to the surface of each mound. For best results, 1) apply in cool weather, 65 to 80 °F, or in early morning or late evening hours, 2) avoid disturbing ants, 3) treat new mounds as they appear and 4) thoroughly water the treated area immediately after application.

**Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring pesticides.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

In no event shall FMC or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.