

279-3194

03/01/2005

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

MAR 1 2005

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Callista O . Chukwunenye
FMC Corporation
1735 Market St.
Philadelphia, PA 19103

Dear Dr. Chukwunenye:

SUBJECT: Label Amendments - Revised Master Labels
Aim
EPA Registration No. 279-3194
Aim EC
EPA Registration No. 279-3241
Aim EW
EPA Registration No. 279-3242
Your Application Dated December 21, 2004

The labelings referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide act, as amended is acceptable with the following provisions:

1. Under the "Wild Rice" section of the labelings, in the fourth paragraph, the phrase "24 hours" was deleted from the last sentence. Revise the sentence to read as follows: "If water is lowered for application, it may be reflooded to normal depths 24 hours after the application."
2. Under the "Potato" section of the labelings, in the "Harvest Aid Desiccation Application" section, delete the phrase "5 to 7 days" and replace it (as noted in the latest accepted label of November 8, 2004) with "7 to 14" days after the first application.
3. Under the "Grass" section of the labelings, add the following statement: "Do not make more than three applications per season."

These amended labelings supersedes all previously accepted ones. Please submit one (1) copy of the final printed labelings before you release the products for shipment.

Sincerely yours,


L

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3/21



ACCEPTED
with COMMENTS
in EPA Letter Dated
MAR 1 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
279-3194

**For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA
FOR SALE OR USE IN CALIFORNIA, USE
SHARK Herbicide**

EPA Reg. No. 279-3194 EPA Est. 279-

Active Ingredient: **By Wt.**
Carfentrazone-ethyl: Ethyl α ,2-dichloro-5-[4-
(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-
triazol-1-yl]-4-fluorobenzenepropanoate 40.0%
Inert Ingredients: 60.0%
..... 100.0%

This product contains 40% of active ingredient per pound of product.

U.S. Patent No. 5,125,958

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician: Carfentrazone-ethyl is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. This product contains a granular material (sand) that may cause mechanical irritation to the eyes. Treatment is otherwise controlled removal of exposure followed by symptomatic and supportive care.

See other panels for additional precautionary information.

ACTIVE INGREDIENT MADE IN CHINA, FORMULATED AND PACKAGED IN USA.



FMC Corporation
Agricultural Products Group
Philadelphia, PA 19103
Aim(Cal)_3_12-20-04

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals) Caution

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should:
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment wash waters.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

TABLE OF CONTENTS

| Section | Page |
|--|------|
| Active ingredient | 1 |
| First Aid Instructions | 1 |
| Precautionary Statements | 1 |
| Environmental Hazards | 1 |
| Physical/Chemical Hazards | 1 |
| Directions for Use | 2 |
| Agricultural Use Requirements | 2 |
| Storage and Disposal | 2 |
| General Information | 2 |
| Application Information | 3 |
| Spray Drift Management | 3 |
| Allowable Use Information | 4 |
| Preharvest Intervals | 4 |
| Crop Rotation Restrictions | 4 |
| Fallow Systems | 5 |
| Preplant Burndown | 5 |
| Hooded Sprayer Applications | 5 |
| Harvest Aid Applications | 7 |
| Corn: Field, Seed, Popcorn, Silage, Sweet Corn | 7 |
| Cotton | 9 |
| Berries | 10 |
| Sorghum | 11 |
| Rice | 12 |
| Wild Rice | 13 |
| Soybeans | 13 |
| Small Grains | 15 |
| Tree Fruits and Tree Nuts | 15 |
| Grape | 16 |
| Tobacco | 17 |
| Potato | 18 |
| Grasses | 18 |
| Hops | 19 |
| Warranty Statement | 19 |

4/24

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Not for use or storage in or around the house.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put formulated or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

Container Disposal

Plastic containers: Triple rinse (or equivalent). Then offer for approved pesticide container recycling program, or puncture and dispose of in an approved waste disposal facility. Provided on site incineration is allowed by state and local authorities, containers may be burned, stay out of smoke.

GENERAL INFORMATION

Aim Herbicide is a water dispersible granule formulation. Aim Herbicide is to be mixed with water and applied to labeled crops for selective postemergence control of broadleaf weeds. Weed control is optimized when the product is applied to actively growing weeds up to 4 inches in height. Aim Herbicide is a contact herbicide.

Aim Herbicide is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation, and in subsequent days necrosis and death of the plant occur. Due to environmental conditions and with certain spray tank additives, some herbicidal symptoms may appear on the crop.

Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the

activity of Aim Herbicide. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms may be reduced as weeds hardened off by drought are less susceptible to Aim Herbicide.

Tank Mixtures

Aim Herbicide may be tankmixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide except for specific recommendations on this label. Tank mixtures of Aim Herbicide with EC formulations of other crop protection products, crop oil concentrates, methylated seed oils, silicone based adjuvants, 28% nitrogen or ammonium sulfate may increase crop response.

Adjuvant Use Requirements

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

Mixing and Loading Instructions:

Fill the spray tank 3/4 full with clean water. Make sure the agitation system is operating while adding products. Prepare a slurry of Aim Herbicide in a clean container using clean water. Slowly add the Aim Herbicide/water slurry. Carefully rinse the slurry container adding the rinsate to the spray tank. Complete filling the spray tank to the desired level. The spray tank agitation should be sufficient to ensure uniform spray mixture during application and must continue until the spray tank has been emptied. When tankmixing with other products, Aim Herbicide should be mixed first in the spray tank. After the Aim Herbicide is thoroughly mixed, add the other products as specified on their label. Ensure the compatibility of other products with Aim Herbicide before mixing them together in the spray tank. Avoid the overnight storage of Aim Herbicide spray mixtures. Premixing Aim Herbicide spray solutions in nurse tanks is not recommended.

Maintain continuous and adequate spray solution agitation until all the spray solution has been used.

Do not use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range as appropriate.

Spray Equipment Clean-Out

Many new pesticides are very active at low rates, especially to sensitive crops. Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying Aim Herbicide and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with Aim Herbicide as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.

2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.

3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.

4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tip) separately in an ammonia solution.

5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

Do not apply sprayer cleaning solutions or rinsate to sensitive crops.

5/2

Do not store the sprayer overnight or for any extended period of time with Aim Herbicide spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of Aim Herbicide remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

APPLICATION INFORMATION

GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in chart below, are required where local indigenous endangered plant species are found.

| Buffers to Indigenous Endangered Plant Species | | |
|--|--------------------------------|---------------------------------|
| AIM USE RATE (lbs. ai per acre) | Low Spray Boom Buffer (ft.) | High Spray Boom Buffer (ft.) |
| 0.024 | 20 | 33 |
| 0.031 | 26 | 46 |

Conventional Boom and Nozzle Sprayers

Use a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 18 inches above the crop. Operate the sprayer to avoid the application of high herbicide rates directly over the rows and/or into the whorl of treated crop plants.

Directed Sprayers

Aim Herbicide may be applied with drop nozzles or other spray equipment capable of directing the spray to the target weeds and away from sensitive plant parts. Aim Herbicide may be applied up to the maximum rate for the target crop for the control of larger weed sizes or weeds not controlled with lower use rates. Use appropriate rates of adjuvants such as nonionic surfactants, crop oil concentrates or methylated seed oils.

Hooded Sprayers

Hooded sprayers may also be used to apply Aim Herbicide. Refer to the Hooded Sprayer Section on page 5 for specific adjustment and operation instructions.

AERIAL APPLICATION

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Higher aerial spray volumes are required for harvest aid and defoliation treatments. Higher spray volumes are required when there is a dense weed population or crop canopy.

SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications of dry materials.

Where states have more stringent regulations, they must be observed.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The optimum drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift when applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Spray Droplet Size

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - For aerial application, orient nozzles so that the spray is released parallel to the airstream which results in larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. For aerial applications, solid stream nozzles oriented straight back produce the largest droplets and potentially the least drift.

Boom Length - For some aerial use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height - Making applications at the lowest height that is safe reduces exposure of spray droplets to evaporation and wind movement. Aerial applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety.

Swath Adjustment - Swath adjustment distance must increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind - Drift potential is lowest between winds speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications shall be avoided below 3 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator shall be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity - When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions - Applications shall not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the following morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas - The pesticide shall only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species and non-target crops).

ALLOWABLE AIM HERBICIDE USE INFORMATION

Refer to the crop section of this label for specific product use directions.

| Maximum Allowable Aim Herbicide Use Per Acre Per Season for crops or crop grouping | | |
|--|---|--------------------------------------|
| Total Allowed Aim Herbicide Use Per Season * | | |
| Crop/Crop Group/Crop Subgroup | Aim Herbicide (dry oz./acre) Per Season | Maximum Rate (lb ai/acre) Per Season |
| Vegetable, root (Subgroups 1A and 1B) | 4.02 | 0.096 |
| Vegetable, bulb (Group 3) | | |
| Vegetable, leafy (Group 4) | | |
| Vegetable, brassica (Group 5) | | |
| Vegetable, legume (Group 6) | | |
| Vegetable, fruiting; Okra (Group 8) | | |
| Bushberry (Subgroup 13A) | | |
| Herbs and Spices (Group 19) | | |
| Tropical Fruits | | |
| Rapeseed | | |
| Mustard seed | | |
| Flax seed | | |
| Sunflower seed | | |
| Safflower seed | | |
| Crambe seed | | |
| Borage seed | | |
| Strawberry | | |
| Horseradish | | |
| Sugarcane | | |
| Peanut | | |
| Crop/Crop Group/Crop Subgroup | Aim Herbicide (dry oz./acre) Per Season | Maximum Rate (lb ai/acre) Per Season |
| Vegetable, tuberous and corm (Subgroups 1C and 1D) | 7.5 | 0.181 |
| Citrus fruit (Group 10) | 5 | 0.124 |
| Pome fruit (Group 11) | 5 | 0.124 |
| Stone fruit (Group 12) | 5 | 0.124 |
| Caneberry (Subgroup 13B) | 16 | 0.4 |
| Tree Nut, Pistachio (Group 14) | 5 | 0.124 |
| Grass (Group 17) | 3.7 | 0.093 |
| Tropical Tree Fruit | 5 | 0.124 |
| Small Grains | 1.2 | 0.031 |
| Sorghum (preplant and in-season) | 0.6 | 0.016 |
| Sorghum (harvest aid) | 0.6 | 0.016 |
| Corn | 1.24 | 0.031 |
| Rice | 5.5 | 0.138 |
| Rice ** | 12 | 0.3 |
| Rice, harvest aid only | 1 | 0.025 |
| Cotton | 5 | 0.124 |
| Cotton, harvest aid only | 2 | 0.05 |
| Soybeans (preplant and in-season and harvest aid) | 0.92 | 0.023 |
| Hops | 1.2 | 0.031 |
| Grape | 5 | 0.124 |
| Tobacco | 2 | 0.05 |
| Potato | 7.3 | 0.181 |
| Wild Rice ** | 12 | 0.3 |

*The total allowable usage includes all applications made to the field per calendar year. This includes fallow treatments, burndown treatments and all in-season treatments, including harvest aid.

** In California Only

6/21

PREHARVEST INTERVALS

Refer to the crop section of this label for specific product use directions.

| Preharvest Intervals (PHI) or Maximum Growth Stage for Aim Herbicide Applications | |
|---|---|
| Crop/Crop Group/Crop Subgroup | PHI (Days Before Harvest) or Growth Stage |
| Vegetable, root (Subgroups 1A and 1B) | 0 |
| Vegetable, bulb (Group 3) | 0 |
| Vegetable, leafy (Group 4) | 0 |
| Vegetable, brassica (Group 5) | 0 |
| Vegetable, legume (Group 6) | 0 |
| Vegetable, fruiting; Okra (Group 8) | 0 |
| Bushberry (Subgroup 13A) | 0 |
| Herbs and Spices (Group 19) | 0 |
| Tropical Fruits | 0 |
| Rapeseed | 0 |
| Mustard seed | 0 |
| Flax seed | 0 |
| Sunflower seed | 0 |
| Safflower seed | 0 |
| Crambe seed | 0 |
| Borage seed | 0 |
| Strawberry | 0 |
| Horseradish | 0 |
| Sugarcane | 0 |
| Peanut | 0 |
| Crop/Crop Group/Crop Subgroup | PHI (Days Before Harvest) or Growth Stage |
| Vegetable, tuberous and corm (Subgroups 1C and 1D) | 7 |
| Citrus fruit (Group 10) | 3 |
| Pome fruit (Group 11) | 3 |
| Stone fruit (Group 12) | 3 |
| Caneberry (Subgroup 13B) | 15 |
| Tree Nut, Pistachio (Group 14) | 3 |
| Grass (Group 17) | 0 |
| Tropical Tree Fruit | 3 |
| Small Grains | Jointing Stage |
| Small Grains (harvest aid) | 3 |
| Sorghum (preplant and in-season) | 6 Leaf Collars |
| Sorghum (harvest aid) | 3 |
| Corn | 14 Leaf Collars |
| Rice (preplant and in-season) | 4 |
| Rice (preplant and in-season)** | 60 |
| Rice (harvest aid) | 3 |
| Cotton (preplant and in-season) | 7 |
| Cotton (harvest aid) | 7 |
| Soybeans (preplant and in-season) | V10 |
| Soybean (harvest aid) | 3 |
| Hops | 0 |
| Grape | 3 |
| Tobacco | 6 |
| Potato | 7 |
| Wild Rice** | 60 |

** In California Only

CROP ROTATIONAL RESTRICTIONS

Following an application of Aim Herbicide, a treated field may be rotated to a registered crop at any time. All other crops may be planted after 12 months.

For Aerial Application of Aim Herbicide In California Only:

(Refer to individual crop sections to see if Aim Herbicide application is permitted by air)

For applications near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable vegetation or annual crops:

-Do not apply within 100 feet of all desirable vegetation or crops.

-If wind up to 10 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet of the desirable vegetation or crops.

-Do not apply when winds are in excess of 10 mph or when inversion conditions exist.

FALLOW SYSTEMS

Apply Aim Herbicide by ground or air alone or with other herbicides in the fallow period prior to planting or the emergence of any crop listed on this label to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good weed control.

Apply Aim Herbicide at up to 1.2 dry ozs. (up to 0.031 pound active ingredient) per acre in fallow systems. A nonionic surfactant or crop oil concentrate is required. Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate at 1.5 to 2.0 pints per acre.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate or paraquat. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is added to the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

For all products used in tank mixes, refer to the specific product labels for all restrictions on tankmixing and observe all label precautions, instructions and rotational cropping restrictions.

PREPLANT BURNDOWN

(For all Labeled Crops)

Apply Aim Herbicide alone or with other herbicides or liquid fertilizers as a burn-down treatment prior to planting or emergence of labeled crops to control or suppress weeds. Aim may be used as a burndown treatment for previous crops prior to new plantings. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good control. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with burndown herbicides such as glyphosate, glufosinate, paraquat, 2,4-D, dicamba or Distinct.

Aim Herbicide Plus Glyphosate or Glufosinate

Apply Aim Herbicide at 0.3 to 0.7 dry oz. (0.008 to 0.016 pound active ingredient) per acre in combination with glyphosate or glufosinate products at their labeled rates for increased speed of activity and improved control of weeds such as those listed below.

When applied as directed, Aim Herbicide plus glyphosate or glufosinate will provide:
Increased speed of activity and improved control of listed weeds.

| | |
|------------------------|-----------------------|
| Buttercup, smallflower | Chickweed |
| Dandelion, common | Henbit |
| Kochia | Lambsquarters, common |
| Marestail | Morningglory, spp. |
| Pennycress, field | Shepardspurse |
| Smartweed, PA | Tansymustard |
| Thistle, Russian | |

Aim Herbicide Plus 2,4-D or Dicamba or Distinct

Apply Aim Herbicide at 0.3 to 0.7 dry oz. (0.008 to 0.016 pound active ingredient) per acre in combination with

2,4-D or dicamba or Distinct at the recommended rates for increased speed of activity and improved control of weeds such as those listed below.

When applied as directed, Aim Herbicide plus 2,4-D or Dicamba or Distinct will provide:

Increased speed of activity and improved control of listed weeds.

| | |
|-----------------------|--------------------|
| Buckwheat, wild | Buttercup |
| Henbit | Kochia |
| Lambsquarters, common | Lettuce, prickly |
| Marestail | Morningglory, spp. |
| Pennycress, field | Shepardspurse |
| Smartweed, PA | Tansymustard |
| Thistle, Russian | |

Aim Herbicide Plus Glyphosate or Glufosinate Plus 2,4-D or Dicamba or Distinct

Apply Aim Herbicide at 0.3 to 0.7 dry oz. (0.008 to 0.016 pound active ingredient) per acre in combination with glyphosate or glufosinate plus 2,4-D, or dicamba or Distinct at the labeled use rates for increased speed of activity and improved control of weeds. The three-way combination is recommended for situations with dense weed pressure and difficult to control weeds, including various weeds that may be resistant to glyphosate or phenoxy type herbicides.

Users must follow the most restrictive labeling regarding plant back restrictions, rotational guidelines, methods of application, and surfactant requirements of the tank mixture components.

When tank mixing with fertilizer solutions be sure to prepare an Aim Herbicide premixture of Aim Herbicide and clean water.

For other specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

HOODED SPRAYER APPLICATIONS

(For all labeled Crops)

Aim Herbicide may be applied to the following crops using hooded sprayers in accordance with specific use information in the following

Directions for Use section:

Beans (Snap, Dried, Lima), Blueberries, Borage, Broccoli, Cabbage, Caneberries (Blackberry, Raspberry), Canola, Carrots, Celery, Corn, Cotton, Crambe, Flaxseed, Grain Sorghum, Grapes, Head Lettuce, Mustard greens, Onions, Okra, Peanuts, Field Peas, Cow Peas, Radish, Rice, Soybeans, Spinach, Strawberries, Sugarbeets, Sugarcane, Sunflowers, Triticale, Teosonite, Tropical Fruits, Wheat, Barley, Oats, Tobacco

Other crops included in the following Crop Groups:

Vegetable, root and tuber (Group 1) such as Beets, Carrots, Horseradish, Radish, Turnips, Ginger, Sugar beets, Yams, Sweet potatoes, Potato

Vegetable, leaves of root and tuber (Group 2) such as Beets, Radish, Sugar beets, Turnip tops, Chicory

Vegetable, bulb (Group 3) such as Chive, Garlic, Onions, Leeks, Scallions, Dry bulb onions, Shallots

Vegetable, leafy (Group 4) such as Celery, Cress, Endive, Lettuce, Parsley, Purslane, Rhubarb, Spinach, Radicchio, Swiss chard

Vegetable, brassica (head, stem and leafy) (Group 5) such as Broccoli, Brussels sprouts, Cabbage, Collards, Kale, Kohlrabi, Greens, Mustard

Vegetable, legume (Group 6) such as Edible peas, Cowpeas, Succulent shelled peas, Dwarf peas, Snap beans, Wax beans, Lima beans, Endamame

8/21

Vegetable, storage of the tuber (Group 7) such as Field pea, Cowpea, Endamame, Lima, Lentil, Beans, Guar, Catina;

Vegetable, fruiting (Group 8) such as Eggplant, Bell pepper, Pimento pepper, Sweet pepper, Tomatillo, Tomato, Pepino;

Vegetable, cucurbit (Group 9) such as Cucumber, Cantaloupe, Summer squash, Winter squash, Pumpkin, Watermelon;

Berries (Group 13) such as Blackberries, Blueberries, Dewberries, Elderberries, Gooseberries, Raspberries, Currant;

Grasses (Group 17) such as Centipede, Bahiagrass, Bermudagrass, Bluegrass, Orchardgrass, Ryegrass, Fescue;

Herbs and Spices (Group 19) such as Chive, Clove, Dill, Cinnamon, Fennel, Ginger, Horseradish, Nutmeg, Parsley;

(For additional information regarding crops within a group, refer to the EPA Website: <http://www.epa.gov/fedrgstr/EPA-PEST/1995/May/Day-17/pr-266.html>)

Directions for Use:

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the above listed crops. This treatment may be made to crops grown in rows, and includes crops grown in rows where mulch or plastic barriers are used as a weed control tool in the drill or plant line. Aim Herbicide may be applied at use rates up to 1.2 dry ozs. (0.031 pound active ingredient) per broadcast acre per application in a minimum of 10 gallons per acre of finished spray. Always refer to the Maximum Allowable Aim Herbicide chart on page 4 of this label for additional use information. Aim Herbicide may be tankmixed with other pesticides registered for crops utilizing this treatment pattern.

Hooded sprayers must be designed, adjusted and operated in such a manner to totally enclose the spray pattern and to prevent any spray deposition to green stems, leaf tissue, flowers or fruit of the crop.

Sprayers shall not be operated at more than five (5) miles per hour in order to minimize vertical movement of the sprayer during application, including the bouncing or raising of the equipment. Use extreme care in applying to fields where the soil surface is uneven, has deep furrows, drains or other contours that would disturb the adjustment and positioning of the spray equipment and/or the spray pattern. Applications must not be made when wind conditions may disturb the spray patterns and result in spray deposition to sensitive plants or plant parts.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

Use a quality spray adjuvant such as crop oil concentrate (COC) or nonionic surfactant (NIS) at the recommended rates.

When used as directed, Aim Herbicide will provide: Control of the listed weeds up to four (4) inches in height or as specified.

| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) per acre |
|---|---|
| Lambsquarters, common (up to 3 inches tall) | 0.3 dry oz. (0.008 pound active ingredient) per acre |
| Morningglory, ivyleaf (up to 3 leaves) | |
| Morningglory, pitted (up to 3 leaves) | |
| Nightshade, Eastern black | |
| Pigweed, redroot | |
| Velvetleaf | |
| Waterhemp (up to 2 inches tall) | |

| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) per acre |
|---|---|
| All the weeds controlled at 0.3 dry oz. (0.008 pound active) per acre plus the weeds listed below: | 0.5 dry oz. (0.013 pound active ingredient) per acre |
| Cheeseweed | |
| Filaree, redstem | |
| Flixweed | |
| Lambsquarters, common | |
| Mallow, common | |
| Morningglory, entireleaf | |
| Morningglory, ivyleaf | |
| Morningglory, pitted | |
| Morningglory, scarlet | |
| Nightshade, hairy | |
| Pennycress, field | |
| Pigweed, prostrate | |
| Pigweed, smooth | |
| Pigweed, tumble | |
| Purslane, common | |
| Sesbania, hemp | |
| Smartweed, PA (seedling) | |
| Tansymustard | |
| Waterhemp | |
| Weeds Controlled | Use Rate dry oz. (pound active ingredient) per acre |
| All the weeds controlled at 0.5 dry oz. (0.013 pound active) per acre plus the weeds listed below: | 0.7 dry oz. (0.016 pound active ingredient) per acre |
| Amaranth, spiny | |
| Anoda, spurred | |
| Bedstraw, catchweed | |
| Buffalobur | |
| Carpetweed | |
| Cocklebur | |
| Copperleaf, hophornbeam | |
| Cotton, GMO Varieties | |
| Cotton, volunteer | |
| Dayflower | |
| Eclipta | |
| Fiddleneck, coast | |
| Groundcherry, smooth (seedling) | |
| Groundcherry, Wright's | |
| Jimsonweed | |
| Kochia | |
| Rocket, London | |
| Morningglories | |
| Nightshade, American black | |
| Nightshade, black | |
| Shepardspurse | |
| Spiderwort, tropical | |
| Thistle, Russian | |
| Wallflower, bushy | |
| Weeds Controlled | Use Rate dry oz. (pound active ingredient) per acre |
| All the weeds controlled at 0.7 dry ozs. (0.016 pound active) per acre plus the weeds listed below: | 0.7 dry oz. (0.025 pound active ingredient) per acre |
| Amaranth, Palmer | |
| Burclover | |
| Spurry, corn | |
| Filaree, broadleaf | |
| Filaree, white | |
| Lettuce, prickly | |
| Mallow, Venice (up to 2 inches tall) | |
| Meadowfoam | |
| Mustard spp. | |
| Redmaids | |

Precautions:

Crop injury will occur when spray is allowed to come in contact with the leaves, green stem tissue, flowers or fruit of the crop.

Restrictions:

Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) during the preplant timing and no more than 2.6 dry ozs. (0.064 pound active ingredient) in-season as a row middle application. Do not apply more than 3.8 dry ozs. (0.096 pound active ingredient) per crop season.

HARVEST AID TREATMENT

Aim Herbicide may be applied to the soybeans and the grain/forage crops (barley, millet, oats, rice, sorghum, triticale, wheat) to defoliate and/or desiccate troublesome broadleaf weeds such as morningglories, pigweeds, velvetleaf and others that may be present at harvest. Aim Herbicide may be used alone or as a tank mixture with other harvest aids.

Applications shall be made when the crop is mature and the grain has begun to dry down, or according to Extension Service recommendations in the use area. Apply Aim Herbicide as a broadcast spray at rates not to exceed the amount as listed in the **MAXIMUM ALLOWABLE AIM HERBICIDE USE TABLE** shown on page 4 of this label. If treatments of Aim Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volume sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

Use a crop oil concentrate (COC) at the rate of 1.0% v/v (1 gallon of COC per 100 gallons of spray solution) or other suitable adjuvant at recommended rates.

Do not apply within 3 days of harvest.

Coverage is essential for satisfactory performance. Repeat application if necessary.

If applied as a tank mixture, refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

CORN

Field Corn, Seed Corn, Popcorn, Corn Silage, and Sweet Corn for Processing and Fresh Market

Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from prior to planting up to 14-leaf collar growth stage.

Applications to corn greater than V8 stage should be made using directed applications to improve weed coverage within the crop canopy and to minimize spray interception by the crop leaves. Do not apply when conditions favor drift or when wind is above 10 mph.

For optimum performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across.

Coverage is essential for good control.

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Under dry conditions, the use of a crop oil concentrate may improve weed control. The use of a crop oil concentrate may increase leaf speckling on the treated corn leaves.

To control weeds not listed on this label, Aim Herbicide may be tank mixed with other herbicides registered for use in corn. When tank mixing Aim Herbicide with other products, be sure Aim Herbicide is added to the spray tank water first and thoroughly mixed. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION sections.

Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions, and rotational cropping restrictions. Adjust sprayers to position spray tips no lower than 18 inches above the crop. Operate the sprayer to avoid the application

of high herbicide rates directly over the rows and/or into the whorl of the corn plant. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

Aim Herbicide Use Rates:

Use Aim Herbicide at 0.3 to 0.7 dry oz. (0.008 to 0.016 pound active ingredient) per acre. Use higher rates when weeds are under stress or are larger.

Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre.

Application Precaution:

The application of Aim Herbicide to corn may result in temporary crop response such as speckling or necrosis of the leaves. Yields will not be affected. Do not make applications when air temperatures are abnormally cool or humidity is high or if the corn foliage is wet from dew, rainfall or irrigation. Users should be aware of these inherent risks and accept these risks prior to application of AimEC.

For additional information regarding potential crop response, refer to the General Information section of the AimEC label.

When used as directed, Aim Herbicide will provide:

Control of the listed weeds up to four (4) inches in height, or as specified.

| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) per acre |
|--|---|
| Lambsquarters, common (up to 3 inches tall) | 0.3 dry oz. (0.008 pound active ingredient) per acre |
| Morningglory, ivyleaf (up to 3 leaves) | |
| Morningglory, pitted (up to 3 leaves) | |
| Nightshade, Eastern black | |
| Pigweed, redroot | |
| Velvetleaf | |
| Waterhemp (up to 2 inches tall) | |
| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) per acre |
| All the weeds controlled at 0.3 dry oz. (0.008 pound active) per acre plus the weeds listed below: | 0.5 dry oz. (0.013 pound active) per acre |
| Cheeseweed | |
| Filaree, redstem | |
| Flixweed | |
| Lambsquarters, common | |
| Mallow, common | |
| Morningglory, spp. | |
| Nightshade, hairy | |
| Pennycress, field | |
| Pigweed, prostrate | |
| Pigweed, smooth | |
| Purslane, common | |
| Sesbania, hemp | |
| Smartweed, PA (seedling) | |
| Tansymustard | |
| Waterhemp | |
| Velvetleaf (up to 24 inches tall) (up to 36 inches for drop nozzle sprayers) | |
| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) Per acre |
| All the weeds controlled at 0.5 dry oz. (0.013 pound active) per acre plus the weeds listed below: | 0.7 dry oz. (0.016 pound active ingredient) per acre |
| Amaranth, spiny | |
| Anoda, spurred | |
| Bedstraw, catchweed | |
| Carpetweed | |
| Cocklebur | |

9/21

| |
|---------------------------------|
| Copperleaf, hophornbeam |
| Cotton, GMO varieties |
| Cotton, volunteer |
| Dayflower |
| Eclipta |
| Fiddleneck, coast |
| Groundcherry, smooth (seedling) |
| Groundcherry, Wright's |
| Jimsonweed |
| Kochia |
| Rocket, London |
| Morningglories, spp. |
| Nightshade, American black |
| Nightshade, black |
| Shepardspurse |
| Spiderwort, tropical |
| Thistle, Russian |
| Wallflower, bushy |

Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) of Aim Herbicide per acre per season including fallow/preplant burndown and labeled crop applications.

Tank Mixtures

Aim Herbicide may be tankmixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide. When tankmixing Aim Herbicide with other products, be sure Aim Herbicide is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with 2,4-D (amine), Accente®, Accent Gold®, Atrazine, Banvel®, Basise®, Basis Gold®, Beacone®, Callisto, Clarity™, Distinct®, Equip®, Exceed®, Hornets®, Liberty®, Lightning®, Marksman®, Northstar™, Option®, Permit®, Poast®, glyphosate products, Scorpion® III, Sencore®, Shotgun®, Spirit™, Steadfast, Sterling®, and Touchdown®.

When tankmixing Aim Herbicide with Accent, Accent Gold, Atrazine, Basis Gold, Liberty, Poast®, glyphosate products for use on GMO corn, and Shotgun use adjuvants recommended on the tank mix partner label. These may include nonionic surfactant, crop oil concentrate, 28% nitrogen, ammonium sulfate or combinations of these.

Leaf speckling can occur when Aim Herbicide is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information. Bromoxynil mixtures and Basagran mixtures may cause significant crop response when in contact with crop foliage.

Aim Herbicide Plus Atrazine

Aim Herbicide may be tankmixed at a rate of 0.3 dry oz. (0.008 pound active ingredient) per acre with Atrazine 4L (16 fluid ounces per acre) or Atrazine 90DF (9 ounces per acre) to control the following weeds:

When used as directed, Aim Herbicide will provide:
Control of listed weeds up to 4 inches tall.

| | |
|---|-------------------|
| Amaranth, Palmer (not triazine resistant) | |
| Amaranth, spiny | Pigweed, redroot |
| Buckwheat, wild | Pigweed, smooth |
| Buffalobur | Pigweed |
| Carpetweed | Potato, volunteer |
| Cocklebur | Purslane, common |
| Copperleaf, hophornbeam | Sesbania, hemp |
| Croton, woolly | Smartweed, annual |
| Devilscaw | Thistle, Russian |
| Eveningprimrose, cutleaf | Velvetleaf |
| Jimsonweed | Mallow, Venice |
| Kochia++ | Anoda, spurred |
| Lambsquarters, common | Waterhemp, common |
| Morningglory, spp. | Waterhemp, tall |
| Nightshade, Eastern black | |
| Nightshade, hairy | |

++ Kochia control up to 2 inches tall with Aim Herbicide + Atrazine + COC only.

Refer to the Atrazine labels for additional weed listings and for higher use rates.

Aim Herbicide Plus Dicamba

Aim Herbicide at 0.3 dry oz. (0.008 pound active ingredient) per acre plus 0.25% v/v nonionic surfactant (2 pints per 100 gallons) can be tankmixed with dicamba herbicides (8 fluid ounces per acre) for control of general broadleaf weeds including the following:

When used as directed, Aim Herbicide will provide:
Control of listed weeds up to 4 inches tall.

| | |
|--------------------|-----------------------------|
| Buckwheat, wild | Pigweed, triazine resistant |
| Cocklebur, common | Potato, volunteer |
| Jimsonweed | Ragweed, common |
| Kochia ++ | Ragweed, giant |
| Lambsquarters | Smartweed, PA (seedling) |
| Morningglory, spp. | Sunflower, common |
| Nightshade, black | Thistle, Russian |
| Pigweed, redroot | Velvetleaf |
| Pigweed, smooth | Waterhemp, common |
| | Waterhemp, tall |

++ Kochia control up to 2 inches tall can be obtained with Aim Herbicide plus atrazine plus COC only. Refer to the dicamba labels for additional weed listings and for higher use rates. Refer to the Tank Mixture Section for information on potential leaf injury.

Aim Herbicide Plus Atrazine Plus Dicamba or 2,4-D

For the control of additional or certain larger weeds up to 6 inches tall Atrazine may be added to the tank mixtures of Aim Herbicide plus dicamba. Aim Herbicide plus 2,4-D (amine) or Aim Herbicide plus dicamba.

Add 2,4-D (amine) to the tank mix at 0.125 to 0.25 pound active ingredient per acre or dicamba at 3 to 4 fluid ounces per acre. Higher rates of atrazine, dicamba herbicides can be used, but do not exceed the recommended label use rates allowed by these labels. Add a 0.25% v/v nonionic surfactant (2 pints per 100 gallons) to the tank mixture. Under very dry soil moisture conditions, the use of crop oil concentrate at 1% v/v (1 gallon per 100 gallon spray solution) may improve weed control. The use of crop oil concentrate may increase leaf speckling. Refer to the Tank Mixture section for information on potential leaf injury.

For control of the following weeds up to 6 inches in height, or as specified, add dicamba at 3 to 4 ounces per acre to Aim Herbicide tank mixes with atrazine or to Aim Herbicide tank mixes with other products that allow the use of dicamba on their labels.

| |
|---|
| Amaranth, Palmer (up to 4 inches) |
| Amaranth, Spiny (up to 4 inches) |
| Cocklebur, common |
| Lambsquarters, common |
| Kochia (up to 4 inches tall) |
| Morningglory spp. |
| Nightshade, Eastern black |
| Nightshade, hairy |
| Pigweed, redroot |
| Pigweed, smooth |
| Ragweed, common |
| Ragweed, giant (up to 4 inches tall) |
| Smartweeds, annual (seedling) |
| Sunflower, common (up to 4 inches tall) |
| Velvetleaf (up to 24 inches) |
| Waterhemp, common |
| Waterhemp, tall |

Special Corn Use Applications

For Directed Applications

Aim Herbicide may be applied with drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl of the corn plant. Aim Herbicide may be used up to the maximum of 1.2 dry ozs. (0.031 pound active) per acre. Rates above 0.5 fl. oz. can be used to aid in control of larger weeds as listed under, "Control of Weeds". Be aware that weeds growing in and under dense canopies may not receive adequate spray coverage necessitating the use of higher spray volumes for acceptable control. Use appropriate rates of adjuvants such as non-ionic surfactant, crop oil concentrate or methylated seed oil.

W/21

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

Seed Corn Production

For seed production fields, apply Aim Herbicide using drop nozzles or other equipment to make a directed spray treatment. Avoid directing spray solution into the whorl.

Seed corn inbreds have generally shown good tolerance to Aim Herbicide, however, all inbreds have not been tested. Broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

Sweet Corn Production

Aim Herbicide may be applied to sweet corn, however, the user assumes all responsibility for herbicide tolerance with such use. All hybrids/varieties have not been tested for sensitivity to Aim Herbicide nor does FMC Corporation have access to all seed company or food processor data. Broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

Therefore, any crop response arising from the use of Aim Herbicide on sweet corn is the responsibility of the user. Use Aim Herbicide only under the recommendation of the Seed Company, food processor, or State Agricultural Extension Service.

COTTON

TIMING AND METHOD OF APPLICATION

Removal of Failed Cotton Stands

Apply Aim Herbicide at the rate of up to 1.0 dry oz. (up to 0.025 pound active ingredient) per acre broadcast as a foliar spray over the top of the remaining cotton plants with sufficient spray volume to provide coverage of the cotton plant, particularly the terminal area. **Coverage is essential for good control.**

Use a crop oil concentrate at 1% v/v (1 gallon per 100 gallons of spray solution).

Do not apply when conditions favoring drift exist or wind is above 10 mph.

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

Post-directed and Lay-by Application

Aim Herbicide is a contact herbicide for postemergence directed sprayer or hooded/shielded sprayer applications for the control of broadleaf weeds in cotton. Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

Applications of Aim Herbicide or Aim Herbicide tank mixes must be made with directed sprayers or hooded sprayers to prevent contact of spray solution with the cotton plant. Do not allow spray solution to contact cotton foliage or green stem tissue. Directed spray equipment must position nozzles a minimum 3 to 4 inches above the soil, with nozzles directed beneath the crop canopy. Aim Herbicide or Aim Herbicide tank mix applications shall be made to cotton that is a minimum of 6 inches in height. Applications to cotton at 5 to 6 nodes or less must be made with hooded or shielded sprayer equipment to completely avoid contact with cotton plants. Lay-by applications of Aim Herbicide or Aim Herbicide tank mixtures at later growth stages of cotton may be made when cotton plants have

achieved a height of 12 inches or more with sufficient bark development and height differential between crop bottom leaves and the soil. Spray solution shall be directed at the base of cotton plants for minimal contact with green stem tissue or foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size.

Do not apply when conditions favoring drift exist or wind is above 10 mph.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

Use a crop oil concentrate at 1% v/v (1 gallon per 100 gallons of spray solution).

Use Rates and Weeds Controlled

Apply Aim Herbicide as a post-directed treatment using a directed sprayer a hooded sprayer or lay-by sprayer delivering a minimum finished spray volume of 10 gallons per acre. Do not apply more than 2 dry oz. (0.05 lb.ai) Aim Herbicide per season by post-directed and lay-by applications.

When applied at 0.5 dry oz. (0.013 pound active ingredient) per acre, Aim Herbicide applied alone will provide:

Control of listed weeds.

| | |
|-------------------------|-----------------------|
| Amaranthus spp. | Purslane, common |
| Sesbania, hemp | Spurge, prostrate |
| | Velvetleaf |
| Lambsquarters | Mallow, Venice |
| Nightshade spp. | Cotton, volunteer |
| Smartweed, Pennsylvania | Cotton, GMO Varieties |

When applied at 0.7 dry oz. (0.016 pound active ingredient) per acre, Aim Herbicide applied alone will provide:

Control of listed weeds.

| All weeds controlled at 0.5 dry oz. plus: | |
|---|--------------------------|
| Anoda, spurred | Morningglory, entireleaf |
| Carpetweed | Morningglory, ivyleaf |
| Cheeseweed | Morningglory, pitted |
| Cocklebur, common | Morningglory, scarlet |
| Fiddleneck, coast | Nettle, stinging |
| Groundcherry, Wright | Sage, lanceleaf |
| Kochia | Shepherdspurse |
| Rocket, London | Spiderwort, tropical |

When applied at 1.0 dry oz. (0.025 pound active ingredient) per acre, Aim Herbicide applied alone will provide:

Control of listed weeds.

| All weeds controlled at 0.7 dry oz. plus: | |
|---|--|
| Ragweed, common | |
| | |

For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with other herbicides such as glyphosate products, Staple, Buctril, Caparol, Cotoran (or other products containing fluometuron), Karmex, MSMA, or other herbicides registered for cotton post-directed and/or lay-by applications. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions and rotational cropping restrictions.

Harvest Aid Application

Aim Herbicide may be applied as a harvest aid to defoliate and desiccate cotton and troublesome weeds that may be present at harvest. It may be used alone or as a tank mixture with other cotton harvest aids.

Use a quality spray adjuvant, such as nonionic surfactant (NIS) or crop oil concentrate (COC) at the recommended rates. NIS is the recommended adjuvant during warmer periods with COC being the better choice for applications during cooler periods.

Make application when 60 to 70 percent of the bolls are open, or according to the State Agricultural Extension Service recommendations in the use area.

Apply Aim Herbicide as a broadcast spray at a rate of up to 1.0 dry ozs. per acre (up to 0.025 lb ai per acre) in spray volume sufficient to provide complete coverage of cotton foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. **Coverage is essential for defoliation.** Repeat application if necessary to remove remaining foliage or control regrowth. Do not apply more than 2.0 dry ozs. (0.05 pound active ingredient) per acre total as a harvest aid. Dense cotton canopy, large plant size, and environmental conditions not conducive to complete plant coverage may reduce initial application performance and increase the need for a second application.

Aim Herbicide may be applied as a tank mix or as a sequential application tankmixed with Dropp, Def, Finish, Prep, Folex, Harvade, Ginstar, CottonQuik, or other registered cotton harvest aid products.

Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions and rotational cropping restrictions.

Restrictions

Do not apply within 7 days of harvest.
Do not apply more than 5.0 dry ozs. (0.124 pound active ingredient) per acre total for preplant and in-season weed control.
Do not apply more than 2.0 dry ozs. (0.05 pound active ingredient) per acre total as a harvest aid.

BERRIES

BUSHBERRY

Such as Blueberry (highbush and lowbush), Currant, Elderberry, Gooseberry, Huckleberry

TIMING AND METHOD OF APPLICATION

Dormant Applications

Aim Herbicide may be applied broadcast to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.

Post-directed Applications For Broadleaf Weed Control

Aim Herbicide may be applied for postemergence weed control of certain susceptible broadleaf weeds at a minimum of 20 gallons finished spray per broadcast acre when used alone or in combination with other herbicides. Apply Aim Herbicide at 0.7 to 1.2 dry ozs. ((0.016 to 0.031 pound active ingredient) per acre for control of susceptible broadleaf weeds. Use the lower rate for control of small seedling weeds at the 2 to 3-leaf stage; use higher rates for control of larger weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in only partial control.

Aim Herbicide may be tankmixed with other registered herbicides that have preemergence or postemergence activity. Any preemergence activity must rely on activity from other herbicides as directed on their labels. Contact herbicides may be tank mixed with Aim Herbicide to obtain a broader spectrum of weeds controlled. If Aim Herbicide is used in a tank mixture, refer to the other product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

Coverage is essential for good control. Use a spray volume adequate to obtain thorough coverage with the minimum being 20 gallons of finished spray per acre. Apply only with ground equipment. Applications may be made with boom equipment, shielded or hooded sprayers, hand-held and high-volume wands or orchard guns. Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a nonionic surfactant (NIS) having at least 80 percent active ingredient at 0.25 % v/v, 2 pints NIS per 100 gallons of spray volume or a quality crop oil concentrate (COC) at recommended rates.

If Aim Herbicide is used in a tank mixture, refer to the other product labels for all restrictions on tankmixing and observe all label precautions, instructions and rotational cropping restrictions.

Band Treatment Applications

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

When applied at up to 1.2 dry ozs. (0.031 pound active ingredient) per acre, Aim Herbicide will provide:
Control of the following weeds.

| | |
|-----------------------|---------------------------|
| Amaranth, Palmer | Morningglory, pitted |
| Burclover | Nettle, burning |
| Cheeseweed | Nettle, stinging |
| Cocklebur, common | Nightshade, black |
| Fiddleneck, coast | Nightshade, Eastern black |
| Filaree, spp. | Nightshade, hairy |
| | Pigweed, redroot |
| | Pigweed, smooth |
| | Lettuce, prickly |
| Lambsquarters, common | Redmaids |
| Rocket, London | Shepherdspurse |
| Mallow, common | Sowthistle |
| Morningglory, ivyleaf | Velvetleaf |

Precautions

Extreme caution must be taken during applications when desirable fruit or foliage is present in order to avoid fruit spotting or leaf necrosis. Do not allow Aim Herbicide spray mist to come in contact with desirable fruit or foliage. On seedling or newly transplanted bushes, do not allow spray to contact green bark of trunk area. Other herbicides may be more injurious to young bushes than Aim Herbicide.

Restrictions

Do not apply within 1 day of harvest.
Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) during the dormant stage, and 2.6 dry ozs. (0.064 pound active ingredient) in-season as a row middle application.
Do not apply more than 3.8 dry ozs. (0.096 pound active ingredient) per crop season.

CANE BERRY

Such as cultivars or hybrids of Blackberry, Boysenberry, Black Raspberry, Red Raspberry

TIMING AND METHOD OF APPLICATION

Post-Directed Application For Primocane and Weed Control

Aim Herbicide is a contact herbicide for directed application for the control of primocanes. Apply when primocanes are approximately 6 inches in height as a directed application of 6.4 dry ozs. (0.1 pound active ingredient) per acre in a minimum of 20 gallons of finished spray per broadcast acre at intervals of 14 to 21 days. Direct the spray to the bottom 18 inches of the canes and also to contact the soil out to 24 inches from each side of the plant row for the control of primocanes and broadleaf weeds.

Band Treatment Applications

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

For weed control apply Aim Herbicide according to the table below using a minimum finished spray volume of 20 gallons per acre. For optimum performance, make applications to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across.

Coverage is essential for good control.

Use a crop oil concentrate at 1% v/v (1 gallon per 100 gallons of spray solution), or a methylated seed oil or organosilicone surfactant at recommended rates.

Do not apply when conditions favor drift or when wind is above 10 mph.

**When applied at 0.5 dry oz. (0.013 pound active ingredient) per acre, Aim Herbicide applied alone will provide:
Control of listed weeds.**

| | |
|-----------------|-------------------------|
| Amaranthus spp. | Purslane, common |
| Bindweed, field | Smartweed, PA (seeding) |
| Hemp Sesbania | Spurge, prostrate |
| Lambsquarters | Velvetleaf |
| Nightshade spp. | |

**When applied at 0.7 dry ozs. (0.016 pound active ingredient) per acre, Aim Herbicide applied alone will provide:
Control of listed weeds.**

| All weeds controlled at 0.5 dry oz. plus: | |
|---|--------------------------|
| Anoda, spurred | Morningglory, entireleaf |
| Carpetweed | Morningglory, ivyleaf |
| Cocklebur, common | Morningglory, pitted |
| Groundcherry, Wright | Morningglory, scarlet |
| Kochia | Sage, lanceleaf |

**When applied at 1.0 dry oz. (0.025 pound active ingredient) per acre, Aim Herbicide applied alone will provide:
Control of listed weeds.**

| All weeds controlled at 0.7 dry oz. plus: |
|---|
| Nightshade, silverleaf (Suppression) |

**When applied at up 1.2 dry ounces (0.031 pound active ingredient) per acre, Aim Herbicide will provide:
Control of the following weeds.**

| | |
|-----------------------|---------------------------|
| | Pigweed, smooth |
| | Prickly lettuce |
| | Morningglory, pitted |
| Amaranth, Palmer | Pigweed, redroot |
| Burclover | Nettle, burning |
| Cheeseweed | Nettle, stinging |
| Cocklebur, common | Nightshade, black |
| Fiddleneck, coast | Nightshade, Eastern black |
| Filaree, spp. | Nightshade, hairy |
| Lambsquarters, common | Redmaids |
| Mallow, common | Sowthistle |
| Morningglory, ivyleaf | Velvetleaf |
| Rocket, London | Shepherdspurse |

For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with other herbicides registered for use in caneberries. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

Restrictions

Do not apply when conditions favor drift or when wind is above 10 mph.

Do not apply more than 16 dry ozs. per acre per season (0.4 pound active ingredient) per acre per season.

Do not make application less than 14 days apart.

Do not apply within 15 days of harvest

SORGHUM (Grain and Forage)

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to sorghum in all tillage systems from prior to planting up through the 6-leaf growth stage.

13/21
Do not apply when conditions favoring drift exist or wind is above 10 mph. For optimum performance, make applications to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.** Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Postemergence broadcast applications of Aim Herbicide with crop oil concentrate are not recommended as increased crop response may occur.

To control weeds not listed on this label, Aim Herbicide may be tankmixed with other herbicides registered for use in grain sorghum. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions, and rotational cropping restrictions. Sprayers shall be adjusted and operated to avoid the application of excessive herbicide rates directly over the row and/or into the whorl of the sorghum plant.

Broadcast applications of Aim Herbicide to sorghum with wet foliage or application during periods of adverse environmental conditions such as cool, cloudy, wet, or high humidity may cause increased crop response. For additional information on crop response, refer to the General Information section of the Aim Herbicide label.

Use Rates

Use Aim Herbicide at 0.5 to 1 fl. oz. (0.008 to 0.016 pound active ingredient) per acre. Use higher rates when weeds are under stress or are larger. Aim Herbicide use rates of 0.6 to 1 fl ozs may only be made with directed spray equipment or hooded sprayers.

Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre.

**When used as directed, Aim Herbicide will provide:
Control of the listed weeds up to four (4) inches tall unless otherwise specified.**

| Weeds Controlled | Aim Herbicide Use Rate dry oz. (pound active ingredient) per acre |
|---|---|
| Lambsquarters, common (up to 3 inches tall) Morningglory, ivyleaf (up to 3 leaves) Morningglory, pitted (up to 3 leaves) Nightshade, Eastern black Pigweed, redroot Velvetleaf (up to 18 inches) Waterhemp (up to 2 inches tall) | 0.3 dry oz. (0.008 pound active ingredient) per acre |
| All the weeds controlled at 0.3 dry oz. (0.008 pound active) per acre plus the weeds listed below: Cheeseweed Filaree, redstem Flixweed Lambsquarters, common Mallow, common Morningglory, spp. Nightshade, hairy Pennygrass, field Pigweed, prostrate Pigweed, smooth Purslane, common Sesbania, hemp Smartweed, PA (seeding) Tansymustard Waterhemp (common) Waterhemp (tall) Velvetleaf (up to 24 inches) | 0.5 dry oz. (0.013 pound active ingredient) per acre |

14/21

Tank Mixtures

Aim Herbicide may be tankmixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide except for specific recommendations on this label. When tank mixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with 2,4-D (amine), Atrazine, Banvel®, Clarity™, Laddok®, Paramount, Peak®, Permite, Starane® and Sterling®.

Leaf speckling can occur when Aim Herbicide is used with certain formulations of crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information.

For Directed Applications

Drop nozzles are recommended if applications are to be made under any of these conditions such as cool, cloudy, wet, or high humidity to limit the amount of product deposited onto sorghum leaves and/or into the sorghum whorl. Aim Herbicide may be used up to the maximum of 0.7 dry oz. (0.016 pound active ingredient) per acre using drop nozzles for control of larger weed sizes for those weeds listed under "Control of Weeds".

When applying Aim Herbicide postemergence to sorghum grown for seed, the use of drop nozzles is recommended to direct spray from uppermost crop leaves and the sorghum whorl.

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

Precautions

Drop nozzles should be used to minimize spray solution contact with crop foliage when the Aim Herbicide use rate is higher than 0.3 dry oz. (0.008 pound active ingredient) per acre.

Restrictions

Do not apply more than 0.6 dry oz. (0.016 pound active ingredient) per acre per season including fallow, preplant burndown and labeled crop applications.

RICE

(For Rice Grown in the Southern United States only)

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Apply to rice in all tillage systems from 30 days before planting up to 3 days before harvest. Aim Herbicide may be applied with either ground or aerial spray equipment. Do not apply when conditions favor drift.

To control weeds not listed on this label, Aim Herbicide may be tankmixed with other herbicides registered for use on rice. When tankmixing Aim Herbicide with other products, be sure Aim Herbicide is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions, and rotational cropping restrictions.

Postemergence Pre-flood Applications to Dry Seeded Rice

Apply Aim Herbicide at 1.0 to 2.0 dry ozs. (0.025 to 0.05 pound active ingredient) per acre. Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons per acre or by air at a minimum finished spray volume of 3 gallons per acre. For optimum results, Aim Herbicide should be applied to weeds up to 4 inches tall and rosettes less than 3 inches across. Use a quality nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For

more active treatments, use a Crop Oil Concentrate (COC) at 0.5 to 1.0% v/v (one half to one gallon per 100 gallons). Apply when the rice is at the 2 leaf stage or larger, but prior to flooding. Some leaf speckling may occur.

**When used as directed Aim Herbicide will provide:
Control of listed weeds up to 4 inches tall.**

| | |
|-------------------------|--------------------------|
| Cocklebur, common | Morningglory, spp. |
| Copperleaf, hophornbeam | Sesbania, hemp |
| Dayflower, spreading | Smartweed, PA (seedling) |
| Groundcherry, cutleaf | Water hyssop |
| Jointvetch, Indian | Purslane, common |
| Jointvetch, northern | Redweed |
| | Pigweed spp. |

Suppression of listed weeds.

| | |
|---------------|-----------------|
| Alligatorweed | Flatsedge, rice |
| Ducksalad | Redstem |
| Eclipta | Texasweed |

Tank Mixtures

For control of weeds listed as suppressed or not listed on this label, Aim Herbicide may be applied following a preemergence grass herbicide or may also be tankmixed with other rice herbicides for broad spectrum weed control. Tank mix applications should be used when rice is well established and in the appropriate stage of growth for treatment with Aim Herbicide and the tank mix partner. For optimum results, weed species should also be in the proper stage of growth as specified on the Aim Herbicide and tank mix partner label. Read and follow all manufacturers' label recommendations for the companion herbicide except for specific recommendations on this label. Do not add a surfactant or crop oil concentrate when tankmixing herbicides formulated as emulsifiable concentrates. Use a nonionic surfactant (NIS) at 0.25% by volume with tank mix partners formulated as dry or liquid flowables.

When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, Aim Herbicide may be applied before, after, or with an application of propanil tank mixed with other herbicides, registered for use on rice. Observe all applicable directions, restrictions and precautions on the partner herbicide labels.

Post Flood Applications to Exposed Weeds

Aim Herbicide may be applied to rice and weeds after the establishment of the permanent flood and when 80% of the foliage of the weeds are exposed. Apply Aim Herbicide at 1.6 to 6.4 dry ozs. per acre (0.025 to 0.10 pound active ingredient) per acre to actively growing weeds. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 1.0% v/v (one gallon per 100 gallons. Apply when the rice is at the 2-leaf stage or later but before internode elongation. Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre. For optimum results, applications should be made to small rather than large weeds. Do not apply to rice after internode elongation. If water level has been lowered to allow this treatment, it should be returned to normal levels 24 hours following treatment. Users of Aim Herbicide must hold the water on the rice fields for 30 days following treatment.

**When used as directed, Aim Herbicide will provide:
Control of listed weeds.**

| | |
|----------------------|-------------------|
| Arrowhead, annual | Morningglory spp. |
| Jointvetch, Indian | Sesbania, hemp |
| Jointvetch, northern | |

Suppression of listed weeds up to 4 inches.

| | |
|----------------------|-----------------|
| Alligatorweed | Ducksalad |
| Ammannia, purple | Flatsedge, rice |
| Dayflower, spreading | Texasweed |

Restrictions

Do not apply more than 6.0 dry ozs. (0.138 pound active ingredient) of Aim Herbicide per acre per season including fallow/preplant burndown and other labeled crop applications.

RICE

(For Rice Grown in California Only)

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre.

Do not apply Aim Herbicide by air.

Do not apply within 1/2 mile of sensitive crops. Do not apply when conditions favoring drift exist. Do not apply more than 12 dry ozs. (0.3 pound active ingredient) per acre per season including fallow, preplant, burndown, and labeled crop applications. Do not apply within 60 days of harvest.

Users of Aim Herbicide must hold the water on the rice fields for 30 days when applications are made to flooded fields.

To control weeds not listed on this label, Aim Herbicide may be tank mixed with other herbicides registered for use on rice. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions.

Early Postseeding Applications to Submerged Weeds

Apply Aim Herbicide at 8 dry ozs. (0.2 pounds active ingredient) per acre. Evenly distribute the spray solution over the flooded rice. The floodwater must be 3 to 6 inches deep. Apply at the 2 to 4 leaf stage of rice. Earlier applications may cause unacceptable crop response. Rice must be well rooted and actively growing at the time of application. Hold the floodwater static for at least five days after application of Aim Herbicide.

When used as directed Aim Herbicide will provide:

Control of listed weeds at the 2 leaf stage or less.

Bulrush, Ricefield
Arrowhead, California
Ammannia, purple
Ammannia, redstem
Umbrellaplant, smallflower

Tank Mixtures

Aim Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide except for specific recommendations on this label. When tank mixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first.

Aim Herbicide may be applied before, after, or with an application of Londax®, Ordram® and Bolero® herbicides. Observe all applicable directions, restrictions (including water holding requirements) and precautions on the Londax, Ordram and Bolero labels.

Foliar Applications to Emerged Weeds Above the Water Surface

Apply Aim Herbicide to weeds at 4 dry ozs. (0.10 pound active ingredient) per acre to the foliage of exposed weeds. At least 80% of the weed foliage must be exposed before spraying with Aim Herbicide. For optimum results, apply to actively growing weeds 20 to 45 days postseeding or the earliest practical opportunity to spray. Weed control is enhanced with greater weed exposure. If the field was drained at application, relood twenty-four hours after application to the normal flood depth.

When used as directed Aim Herbicide will provide:

Control or suppression of the following weeds.

Bulrush, Ricefield
Arrowhead, California
Ammannia, purple (suppression only)
Ammannia, redstem (suppression only)

Umbrellaplant, smallflower (suppression only)

Crop Response

Some temporary leaf speckling may occur shortly after application.

Tank Mixes

Aim Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Aim Herbicide may be tank mixed with propanil-containing herbicides, Londax®, Bolero®, or Whip® herbicides. Not all combinations of Aim Herbicide and other formulated herbicides have been tested. In general, the EC formulations, nonionic and silicone based surfactants and crop oil concentrates, when mixed with Aim Herbicide will increase leaf speckling on the rice leaves. These tank mixtures should be tested on a small portion of the field to ensure crop safety prior to general use.

WILD RICE

(For Wild Rice Grown in California Only)

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre.

Do not apply by air.

Do not apply within 1/2 mile of sensitive crops. Do not apply when conditions favoring drift exist. Do not apply when winds exceed 10 mph. Do not apply more than 12.0 dry ozs. (0.3 pound active ingredient) per acre per season, including fallow/preplant, burndown, and labeled crop applications. Do not apply within 60 days of harvest.

Users of Aim Herbicide must hold the water on the rice fields for 30 days when applications are made to flooded fields.

Apply Aim Herbicide to weeds at the rate of 4.0 to 8.0 dry ozs. (0.1 to 0.2 pound active ingredient) per acre to the foliage of exposed weeds above the water surface. Make applications after the floating leaf stage through tillering. The water in paddies may be lowered if practical. Smaller weeds with more leaf area exposed will give better control. If water is lowered for application, it may be re-flooded to normal depths after the application.

When used as directed Aim Herbicide will provide:

Control or suppression of the following weeds.

Bulrush, ricefield
Arrowhead, California
Waterplantain, common (Suppression only)
Burrweed, giant (Suppression only)
Ammannia, purple (Suppression only)
Ammannia, redstem (Suppression only)
Umbrellaplant, smallflower (Suppression only)

Crop Response

Some temporary leaf speckling may occur following application.

Tank Mixtures

Aim Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Not all combinations of Aim Herbicide and other formulated herbicides and adjuvants have been tested. In general, EC formulations, nonionic and silicone based surfactants, and crop oil concentrates, will increase leaf speckling on the wild rice leaves. These tank mixes should be tested on a small portion of the field to ensure crop safety prior to general use.

SOYBEANS

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to soybeans in all tillage systems from prior to planting up to emergence. Do not apply Aim Herbicide during a period from emergence to V2. After plants

have reached V3, applications can be made up to V10. Do not apply when conditions favoring drift exist.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. Use the higher rates when treating more mature weeds or dense vegetative growth. **Coverage is essential for good control.**

For additional information on crop response refer to the General Information section of the Aim Herbicide label.

Broadcast Postemergence Applications

Apply Aim Herbicide at 0.15 dry oz. (0.004 pound active ingredient) per acre for the control of velvetleaf. Do not apply Aim Herbicide to soybeans with maturities less than Group 2.0. For soybeans of maturity Group 2.1 to 3.4, Aim Herbicide may be used at rates up to 0.25 fl. oz. per acre. Use caution when making applications when making these treatments.

For later maturing soybeans than Group 3.5, Aim Herbicide may be applied at rates up to 0.3 dry oz. (0.008 pound active ingredient) per acre. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints NIS per 100 gallons of spray solution) having at least 80% active ingredient.

Broadcast Application Precaution

The application of Aim Herbicide to soybeans may result in crop response. Soybeans may show some burn, speckling or necrosis of crop leaves. Soybeans quickly outgrow initial herbicide effects and yields are not affected. Do not make applications during conditions of abnormal cool, high humidity or if foliage is wet from dew, rainfall or irrigation. Users should be aware of these potential effects prior to making applications. If the user is not willing to accept these risks, applications should not be made.

For additional information on crop response, refer to the General Information section of this label.

Tank Mixtures

Aim Herbicide may be tankmixed with other herbicides to control weeds not listed on this label. **Do not use with diphenylether herbicides.** Read and follow all manufacturer's label directions for the mixture herbicide except for specific recommendations on this label. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is added in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with glyphosate or glufosinate products for use on GMO soybeans. Leaf injury can occur when Aim Herbicide is used with certain formulations of crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information.

**When used as directed Aim Herbicide at 0.15 dry oz. (0.004 pound active ingredient) per acre will provide:
Control of listed weeds up to 4 inches tall.**

| | |
|------------|--|
| Velvetleaf | |
|------------|--|

**When used as directed, Aim Herbicide at 0.3 dry oz. (0.008 pound active ingredient) per acre will provide:
Control of weeds up to 4 inches tall, or as specified.**

| | |
|-----------------------|---|
| Nightshade, black | Morningglory, ivyleaf (up to 3 true leaves) |
| Lambsquarters, common | Morningglory, Pitted (up to 3 true leaves) |
| Pigweed, redroot | Waterhemp, spp. (up to 3 inches tall) |

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications of this label for additional specific use directions.

For Directed Applications

Use Aim Herbicide at 0.3 to 0.9 dry oz. (0.008 to 0.023 pound active ingredient) per acre. Applications shall be made by ground equipment using a finished volume of 10 to 20 gallons of spray per acre. When soybeans are grown under very dry soil moisture

conditions, a high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v) or 2 to 4 gallons per 100 gallon spray solution) may be used in addition to the nonionic surfactant. Apply as a post-directed treatment with spray directed toward the base of the plant and avoid contact with soybean foliage. The use of spray shields may reduce spray contact with soybean foliage. Aim Herbicide contact with soybean foliage can result in significant crop response.

**When used as directed Aim Herbicide at the rate of 0.3 dry oz. (0.008 pound active ingredient) per acre, will provide:
Control of the listed weeds up to four (4) inches in height, or as specified.**

| | |
|---|---------------------------------|
| Lambsquarters, common (up to 3 inches tall) | Pigweed, redroot |
| Morningglory, ivyleaf (up to 3 leaves) | Velvetleaf |
| Morningglory, pitted (up to 3 leaves) | Waterhemp (up to 2 inches tall) |
| Nightshade, Eastern black (up to 4 inches tall) | |

**When used as directed Aim Herbicide, at the rate of 0.5 dry oz. (0.013 pound active ingredient) per acre, will provide:
Control of the listed weeds up to four (4) inches in height, or as specified.**

| | |
|--|--------------------------|
| All the weeds controlled at 0.3 dry oz. (0.008 pound active ingredient) per acre plus the listed weeds: | |
| Bindweed, field (Above ground plant parts only) | Pennycress, field |
| Cheeseweed | Pigweed, smooth |
| Filaree, redstem | |
| Flixweed | Pigweed, prostrate |
| Lambsquarters, common | Purslane, common |
| Mallow, common | Sesbania, hemp |
| Morningglory spp. | Smartweed, PA (seedling) |
| Nightshade, hairy | Tansymustard |
| Waterhemp, tall | Waterhemp, common |

**When used as directed Aim Herbicide, at the rate of 0.7 dry oz. (0.016 pound active ingredient) per acre, will provide:
Control of the listed weeds up to four (4) inches in height, or as specified.**

| | |
|--|---------------------------------|
| All the weeds controlled at 0.5 dry oz. (0.013 pound active ingredient) per acre plus the listed weeds: | |
| Amaranth, spiny | Groundcherry, Wright's |
| Anoda, spurred | Groundcherry, smooth (seedling) |
| Bedstraw, catchweed | Jimsonweed |
| Buffalobur | Kochia |
| Carpetweed | Rocket, London |
| Cocklebur | Morningglories |
| Copperleaf, hophornbeam | Nightshade, black |
| Cotton, volunteer | Nightshade, American black |
| Cotton, GMO Varieties | Spiderwort, tropical |
| Dayflower | Shepardspurse |
| Eclipta | Thistle, Russian |
| Fiddleneck, coast | Wallflower, bushy |

**When used as directed Aim Herbicide, at the rate of 0.9 dry oz. (0.023 pound active ingredient) per acre, will provide:
Control of the listed weeds up to four (4) inches in height.**

| | |
|--|--------------------------------------|
| All the weeds controlled at 0.7 dry oz. (0.016 pound active ingredient) per acre plus the listed weeds: | |
| Ammannia, purple | Lettuce, prickly |
| Buckwheat, wild | Mallow, Venice (up to 2 inches tall) |
| Buffalobur | Meadowfoam |
| Burclover | Mustard spp. |
| Filaree, broadleaf | Redmaids |
| Filaree, white | Spurry, corn |

Restrictions

Do not apply more than 0.9 dry oz. (0.023 pound active ingredient) per season.
Do not feed treated soybean forage or soybean hay to livestock.
Do not use with diphenylether herbicides.
Do not apply when conditions favoring drift exist.

Do not apply when crop foliage is wet from dew, rainfall or irrigation.

SMALL GRAINS

Such as Barley, Grain and Forage Millets, Oats, Rye, Teosinte, Triticale, and Wheat

TIMING AND METHOD OF APPLICATION

Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to small grains in all tillage systems from prior to planting up to jointing. For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. For dense weed pressure, use the higher recommended rate plus tank mix combinations. **Coverage is essential for good control.**

Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. A high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v or 2 to 4 gallons per 100 gallon spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre may be used in addition to the nonionic surfactant.

To control weeds not listed on this label, Aim Herbicide may be tankmixed with other registered herbicides.

When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions, and rotational cropping restrictions. Aim Herbicide may be applied by ground or air. **Coverage is essential for good control.** Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre. Applications made by air shall utilize a minimum finished spray volume of 3 gallons per acre. Up to half of the spray volume (by air or ground) may be liquid nitrogen fertilizer.

When applied at 0.3 to 0.7 dry oz. (0.008 to 0.016 pound active ingredient) per acre Aim Herbicide will provide:

Control of listed weeds up to 4 inches tall, or as specified.

| | |
|--------------------------------|-------------------|
| Bedstraw, catchweed | Nightshade, black |
| Fiddleneck, coast | Cheeseweed |
| Flixweed | Pennycress, field |
| Lambsquarters (up to 3 inches) | Pigweed, redroot |
| Rocket, London | Velvetleaf |
| Mustard, tansy | Mallow, common |

Suppression of listed weeds up to 4 inches tall.

| | |
|-------------------|------------------|
| Bindweed, field | Mustards |
| Filaree, redstem | Shepherdspurse |
| Nightshade, hairy | Thistle, Canada |
| Kochia | Thistle, Russian |
| Lettuce, prickly | Buckwheat, wild |

When applied at 0.9 to 1.2 dry oz (0.023 to 0.031 pound active ingredient) per acre Aim Herbicide will provide:

Control of the listed weeds up to 4 inches tall.

| | |
|--|--------------------|
| All of the weeds controlled at 0.5 to 1.0 dry oz. (0.008 to 0.016 pound active ingredient) per acre, plus the following weeds: | |
| Bittercress | Nightshade, hairy |
| Buckwheat, wild | Pigweeds |
| Filaree, redstem | Shepherdspurse |
| Kochia | Sowthistle, annual |
| Lambsquarters | Thistle, Russian |
| Mustard, tumble | Wallflower, bushy |

Tank Mixtures With Other Herbicides

To control additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with other labeled herbicides. Refer to the other product label for specific instructions and restrictions, including the

adjuvant recommendations. Tank mixtures with other EC or Ester formulations may increase leaf speckling. Do not use Aim Herbicide with crop oil concentrates (COC), methylated seed oils (MSO) or silicone based adjuvants.

With 2,4-D (amine or ester) or MCPA (amine or ester)

Aim Herbicide may be tank mixed at a rate of 0.3 to 0.7 dry oz. (0.008-0.016 pound active ingredient) per acre with 2,4-D (amine or ester) or MCPA (amine or ester) for use on small grains. For optimum results add 2,4-D (amine or ester) to the tank at 0.25 lb. acid equivalent per acre or MCPA (amine or ester) at 0.375 lb acid equivalent per acre. Higher rates of these herbicides can be used, but do not exceed the recommended label use rates allowed by these labels. Add nitrogen fertilizer (2 to 4% v/v) 2 to 4 gallons per 100 gallons or ammonium sulfate 4 lbs. per acre) to the tank mixture.

When applied as directed, Aim Herbicide in tank mixtures with 2,4-D (amine or ester) or MCPA (amine or ester) herbicides will provide:

Control of listed weeds up to 4 inches tall.

| | |
|-----------------------|---------------------------|
| Bedstraw, catchweed | |
| Buckwheat, wild | Pennycress, field** |
| Cocklebur | Pepperweed, greenflower** |
| Croton, woolly | Pigweed, prostrate |
| Fiddleneck | Pigweed, redroot |
| Filaree, redstem | Pigweed, smooth |
| Flixweed** | Pigweed, tumble |
| Gromwell, common | Primrose, cutleaf |
| Groundsel, common | Amaranthus, spp. |
| Knotweed, prostrate* | Radish, wild |
| Kochia | Ragweed, common |
| Lambsquarters, common | Ragweed, giant |
| Lettuce, miners | Sowthistle, annual |
| Lettuce, prickly | Speedwell, ivyleaf |
| Rocket, London** | Sunflower, wild |
| Mustard, blue*** | Tarweed, coast |
| Mustard, tansy** | Thistle, Russian |
| Mustard, tumble** | Wallflower, bushy |
| Mustard, wild** | Waterhemp, tall |
| Nightshade, black | |

*For Knotweed control, use Aim Herbicide + 2,4-D (amine or ester) only.

**These weeds can be treated from the rosette through bolting growth stages.

***Apply to rosette growth stage (before bolting) of blue mustard.

Restrictions

Do not apply when conditions favoring drift exist.

Do not harvest for forage within 7 days of application.

Do not apply more than 1.2 dry ozs. of Aim Herbicide (0.031 pound active ingredient) per acre per season including fallow or preplant burndown and labeled crop applications.

TREE FRUIT, TREE NUT and OTHER CROPS

Citrus Fruits such as: Calamondin, Citrus Citron, Chironja, Tangelo, Tangor, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (sour), Orange (Sweet), Pummelo, Satsuma Mandarin

Pome Fruits such as: Apple, Crabapple, Loquat, MayHaw, Pear, Pear (Oriental), and Quince

Stone Fruits such as: Apricot, Cherry (Sweet), Cherry (Tart), Nectarine, Peach, Plum, Plum (Chickasaw), Plum (Damson), Plum (Japanese), Prune, and Plumcot

Tree Nuts such as: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio Walnut (Black and English)

Tropical Fruits such as Papaya, Avocado, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star apple, Guava, Feijoa, Jaticaba, Wax jambu, Starfruit, Passionfruit, Acerola, Lychee,

Longan, Spanish lime, Rambutan, Pulasan, Sugar apple, Atemoya, Custard apple, Cherimoya, Llama, Soursop, and Biriba.

Other Crops such as: Kiwifruit, Pomegranate, Fig, Olive, Date, Persimmon, Banana, Cacao, Tea, Indian Mulberry, Vanilla, Coconut, Palm Heart, Coffee and Guayule.

TIMING AND METHOD OF APPLICATION

Weed Control

Apply Aim Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply Aim Herbicide up to 1.2 dry ozs. (up to 0.031 pound active ingredient) per acre. Aim Herbicide alone or tank mixtures may be used for general weed control, in middles (between rows of trees), and in strips (in row of trees). Aim Herbicide may be applied at any time during the season. Aim Herbicide may be mixed with other herbicides that have preemergence or postemergence activity. Any preemergence activity must rely on activity from other herbicides as directed on their labels. Contact herbicides such as glyphosate and paraquat may be tankmixed with Aim Herbicide for broader spectrum weed control.

Chemical Mowing

Aim Herbicide may be used alone or in tank mixtures with other herbicides in chemical mowing practices for orchard vegetation management.

Sucker Management

Aim Herbicide may be used in the management of undesirable sucker growth from the base of the trunks or root sprouts. Apply Aim Herbicide at 1.2 dry ozs. (0.031 pound active ingredient) per acre. Suckers and other undesirable growth must be treated when the tissue is young and not mature and hardened off. Care must be taken not to allow spray mist to contact desirable fruit or foliage or green bark (see Precautions).

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the **Hooded Sprayer Applications** section of this label for additional specific use directions.

Equipment and Application

Coverage is essential for good control. Use a spray volume adequate to get thorough coverage, but use a minimum of 20 gallons of finished spray per acre. Apply only with ground equipment. Applications may be made with boom equipment, hooded sprayers, shielded sprayers, hand-held and high volume wands or orchard guns. Always add Aim Herbicide to the spray tank first. See "Mixing and Loading Instructions" under GENERAL INFORMATION.

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate at 1% v/v (one gallon COC per 100 gallons). Aim Herbicide may also be applied with recommended rates of MSO or silicone adjuvants.

Postemergent Weed Control of Broadleaf Weeds: Apply Aim Herbicide up to 1.2 dry ozs. (up to 0.031 pound active ingredient) per acre for control of susceptible broadleaf weeds. The lower rate is for small seedling weeds at the 2 to 3-leaf stage; higher rates are needed for larger weeds up to the 6-leaf stage. Applications to weeds beyond the six-leaf stage may result in only partial control.

When used as directed, Aim Herbicide will provide:

Control of the listed weeds.

| | |
|-----------------------|---------------------------|
| Amaranth, Palmer | Momingglory, pitted |
| Burclover | Nettle, burning |
| Cheeseweed | Nettle, stinging |
| Cocklebur, common | Nightshade, black |
| Fiddleneck, coast | Nightshade, Eastern black |
| Filaree, broadleaf | Nightshade, hairy |
| Filaree, redstem | Pigweed, redroot |
| Filaree, whitestem | Pigweed, smooth |
| Dayflower | Lettuce, prickly |
| Lambsquarters, common | Redmaids |

| | |
|-----------------------|----------------|
| Rocket, London | Shepherdspurse |
| Mallow, common | Sowthistle |
| Morningglory, ivyleaf | Velvetleaf |
| Balsamapple | |

Precautions

Extreme caution must be used during applications when desirable fruit and/or foliage are present in order to avoid fruit spotting and/or leaf necrosis. Do not allow spray mist of Aim Herbicide to come in contact with desirable fruit and/or foliage. On seedling or newly transplanted trees do not allow spray to contact green bark of trunk area. Other herbicides may be more injurious to young trees than Aim Herbicide. When tank mixtures are used, the precautions and restrictions on the labels of all tankmixed herbicides must be followed.

Restrictions

Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) per acre per application and 7.9 dry ozs. (0.124 pound active ingredient) per acre per season, including preplant site preparation.

Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) per acre in a single application for other crops (Tropical Fruits, Pistachio, Kiwifruit, Pomegranate, Fig, Olive, Date, Persimmon, Banana, Cacao, Tea, Indian Mulberry, Vanilla, Coconut, Palm Heart, Coffee and Guayule).

Do not make applications less than 14 days apart.

Allow a minimum of three days between last application and harvest.

If Aim Herbicide is used in a tank mixture, observe the other product's label for restrictions, precautions, and rotational cropping instructions.

GRAPE

Raisin, Table, Juice and Wine

TIMING AND METHOD OF APPLICATION

Weed Control:

Aim Herbicide may be applied alone or as a tank mixture with other herbicides as a postemergence directed treatment or as a hooded spray to control emerged and actively growing weeds. Apply Aim Herbicide at up to 2.0 dry ozs. (0.031 pound active ingredient) per acre. Applications may be made to middles (between rows of plants) and in strips (in row of plants). Aim Herbicide may be applied at any time during the season (see precautions). Aim Herbicide may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any pre-emergence activity must rely on activity from other herbicides as directed on their labels. Herbicides such as glyphosate may be tank mixed with Aim Herbicide for broader spectrum weed control. If Aim Herbicide is used in a tank mixture, observe the other product's label for restrictions, precautions and rotational cropping instructions.

Sucker Management: Aim Herbicide may be used to aid in the management of undesirable sucker growth from the base of vine trunks or root sprouts. Apply Aim Herbicide at 1.2 dry ozs. (0.031 pound active ingredient) per acre. Suckers and other undesirable growth must be treated when the tissue is young and not mature and hardened off. Care must be taken not to allow spray mist to contact desirable fruit or foliage or on to green bark (see precautions). Aim Herbicide may be applied with other sucker control herbicides.

Hooded Sprayer Applications

Aim Herbicide may be applied with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the **Hooded Sprayer Applications** section of this label for additional specific use directions.

Equipment and Application: Coverage is essential for good control. Use a spray volume adequate to obtain thorough coverage and use a minimum of 10 gallons of finished spray per acre. Apply only with ground equipment. Applications may be made with hooded sprayers, boom equipment, shielded sprayers, hand-held and high-volume wands or orchard guns. **Always add Aim**

Herbicide to the spray tank first. See "Mixing and Loading Instructions" under GENERAL INFORMATION.

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one gallon COC per 100 gallons).

Postemergent Weed Control of Broadleaf Weeds

Apply Aim Herbicide at 0.7 to 1.2 dry ozs. (0.016 to 0.031 pound active ingredient) per acre for the control of susceptible broadleaf weeds. Lower rates may be used to control small seedling weeds at the 2 to 3 leaf stage. Higher rates are needed for larger weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in only partial control.

When applied at up to 1.2 dry ozs. (0.031 pound active ingredient) per acre Aim Herbicide will provide:
Control of listed weeds.

| | |
|-----------------------|---------------------------------|
| Pigweed, redroot | |
| Pigweed, smooth | Lettuce, prickly |
| Amaranth, Palmer | Morningglory, pitted |
| Burclover | Nettle, burning |
| Cheeseweed | Nettle, stinging (foliage only) |
| Cocklebur, common | Nightshade, black |
| Fiddleneck, coast | Nightshade, Eastern black |
| Filaree spp. | Nightshade, hairy |
| Lambsquarters, common | Redmaids |
| Rocket, London | Shepherdspurse |
| Mallow, common | Sowthistle |
| Morningglory, ivyleaf | Velvetleaf |

Precautions: Extreme caution must be used during applications when desirable fruit or foliage is present in order to avoid fruit spotting or leaf necrosis. Do not allow Aim Herbicide spray mist to come in contact with desirable fruit or foliage. Do not use on seedling or newly transplanted vines do not allow spray to contact green bark of trunk area. Other herbicides may be more injurious to vines than Aim Herbicide.

Restrictions

Do not apply more than 1.2 dry ozs. (0.031 pound active ingredient) per acre per application (including preplant site preparation treatments).
Do not apply more than 5.0 dry ozs. (0.124 pound active ingredient) per acre per season.
Do not make application less than 14 days apart.
Allow a minimum of three days between last application and harvest.

TOBACCO

Apply Aim Herbicide alone or as a tank mixture with other registered herbicides to emerged and actively growing weeds at use rates up to 1 dry oz. (0.024 pounds active ingredient) per acre. For optimum performance, make applications to weeds up to 4 inches tall and rosettes less than 3 inches across. Use higher rates when treating more mature weeds or dense vegetative growth. Coverage is essential for good control. Use adequate spray volume to achieve thorough coverage, but a minimum of 10 gallons of finished spray per acre is required. Use a quality crop oil concentrate (COC) at 1% v/v (1 gallon of COC per 100 gallons of spray solution).

Aim Herbicide may be tankmixed with other herbicides registered for use on tobacco to provide additional weed control. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. Refer to the other product label for restriction on tankmixing and observe all label precautions, instructions and rotational cropping restrictions.

For additional information refer to the general information section of the Aim Herbicide label.

TIMING AND METHOD OF APPLICATION

Pre-transplant burndown

Aim Herbicide is a contact herbicide for pre-transplant burndown control of broadleaf weeds in tobacco. Apply Aim Herbicide as a broadcast application alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Aim Herbicide may be applied up to one (1) day prior to transplanting

Shielded spray or Hooded spray

Aim Herbicide may be applied using shielded sprayers or hooded sprayers to emerged and actively growing broadleaf weeds in tobacco from transplanting until layby. Shielded spray or hooded spray applications of Aim Herbicide or Aim Herbicide tank mixes should utilize application equipment that must prevent contact of spray solution with the tobacco plant. Do not allow spray solution to contact tobacco foliage or green stem tissue. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

Directed spray after first priming (Flue Cured Tobacco Only)

Aim Herbicide may be applied as a directed spray application after the first priming in only flue cured tobacco only for the control of emerged and actively growing broadleaf weeds. Directed spray equipment should position nozzles a minimum of 3 to 4 inches above the soil, with nozzles directed underneath the crop canopy. Spray solution should be directed at the base of tobacco plants for minimal contact with foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size. Do not apply when conditions favor drift or wind is above 10 mph.

When applied at 0.5 dry oz. (0.013 pound active ingredient) per acre Aim Herbicide alone will provide:
Control of listed weeds up to 4 inches tall.

| |
|-------------------------------|
| Amaranthus spp. |
| Bindweed, bindweed (burndown) |
| Sesbania, hemp |
| Lambsquarters |
| Nightshade, spp. |
| Purslane, common |
| Smartweed, PA (seeding) |
| Velvetleaf |

When applied at 0.7 dry oz. (0.016 pound active ingredient) per acre Aim Herbicide alone will provide:
Control of listed weeds up to 4 inches tall.

| |
|---|
| All weeds controlled at 0.5 dry ounce plus: |
| Anoda, spurred |
| Carpetweed |
| Cocklebur, common |
| Cotton, volunteer |
| Cotton, GMO Varieties |
| Groundcherry, Wright |
| Kochia |
| Morningglory, spp. |
| Sage, lanceleaf |
| Spiderwort, tropical |

When applied at 0.9 dry oz. (0.023 pound active ingredient) per acre Aim Herbicide alone will provide:
Control of listed weeds.

| |
|---|
| All weeds controlled at 0.7 dry oz. plus: |
| Dayflower, spreading |
| Ragweed, common |

For control of additional broadleaf weeds and grasses, Aim Herbicide may be tankmixed with other herbicides registered for use in tobacco at the appropriate timing. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions and rotational cropping restrictions.

Restrictions

Do not apply within 6 days of Harvest.
Do not apply more than 2.0 dry ozs. (0.048 pounds active ingredient) per acre per season

POTATO

TIMING AND METHOD OF APPLICATION

Aim Herbicide may be used alone or in a tank mix combination with other herbicides and insecticides as a fallow systems treatment, as a preplant burndown treatment and/or as a harvest aid to desiccate potatoes and those susceptible weeds that may be present.

Fallow Systems

See the **Fallow Systems** section for directions for application.

Preplant Burndown

See the **Preplant Burndown** section for directions for application.

Harvest Aid Desiccation Application

Apply Aim Herbicide as a broadcast spray at a rate of 2.0 to 3.8 dry ozs. (0.05 to 0.09 pound active ingredient) per acre in spray volume sufficient to provide complete coverage of potato foliage. Aim Herbicide may be used alone or as a tank mixture with other potato harvest aids as a desiccant prior to harvest. Aim Herbicide can be applied foliarly to potatoes in the later stages of senescence and will provide adequate desiccation of potato foliage and vines. Aim Herbicide will also desiccate late season susceptible broadleaf weeds to aid in tuber harvest. Adequate desiccation is generally achieved within 14 days after the initial treatment is applied. If the potato crop is in the active vegetative growth stage when desiccation is initiated, two applications may be required to provide desiccation of leaf and stem tissue. Dense potato canopy, large plant size, and environmental conditions not conducive to product absorption or activity will reduce initial application efficacy and increase the need for a second application. If a second application is necessary, apply at 5 to 7 days after the first application. Thorough coverage of the potato plant to be desiccated is essential. Use a sufficient volume of water to obtain thorough coverage of the potato leaves and vines. For optimum results, apply Aim Herbicide when the potato crop is in the early stages of natural senescence.

Ground Application

Apply Aim Herbicide in at least 20 gallons of water per acre using 80-degree or 110-degree flat-fan nozzles. Select a spray pressure between 30 to 60 pounds per square inch (psi) measured at the nozzle to obtain a droplet size of approximately 300 microns. Vary the spray volume and spray pressure as indicated by the density of the potato canopy and vines to assure thorough spray coverage. Increase the spray volume and pressure if the potato canopy is dense or under cool, cloudy or dry conditions. Increased spray volumes will enhance performance. If Turbo TeeJet® nozzles are used, a spray pressure of 60 psi or more will be required to obtain thorough coverage. Do not apply when winds are gusty or prone to cause herbicide drift from desired target, particularly when high spray pressures are utilized.

Aerial Application

Apply Aim Herbicide with aerial equipment using 5 to 10 gallons of water per acre, using higher volumes when potato canopies and vines are dense. Apply at a height of 10 feet or less above the potato canopy using drift reduction nozzles. Adjust the nozzles to provide a uniform pattern and a droplet size of 350 to 450 microns. Do not apply aerially when atmospheric conditions are conducive to spray drift and do not apply when wind could drift to surrounding vegetation.

Adjuvant

Aim Herbicide must be applied with a high quality spray adjuvant such as methylated seed oil, COC or an organosilicone at the labeled rates. Adjuvant rates should increase as spray volumes exceed 20 gallons per acre.

Tank mixes

Aim Herbicide may be applied as a tank mix or as a sequential application with other potato desiccants. Refer to the other product's label for restrictions on tankmixing, and observe all label precautions, instructions and rotational cropping restrictions.

Restrictions

Do not apply more than 7.2 dry ozs. of Aim Herbicide (0.18 pound active ingredient) per acre per crop season as a desiccant.
Do not apply when conditions favoring drift exist or wind is above 10 mph.
Do not apply within 7 days of harvest.

GRASS

Such as Forage, Fodder, Hay, Seed and Sod

Aim Herbicide may be applied alone or in combination with other registered pesticides for the control of weeds in rangeland, pastures, hay, grasses grown for hay or silage and grass seed production. Aim Herbicide may be applied at use rates up to 1.2 dry ozs. (0.031 pound active ingredient) per broadcast acre. For optimum results, weeds should be treated when small. Applications shall be made with ground equipment delivering a minimum of 10 gallons of finished spray per acre and adjusted to provide optimum coverage of the target weeds.

Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. A high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v or 2 to 4 gallons per 100 gallon spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre may be used in addition to the nonionic surfactant.

When Aim Herbicide is applied alone, grazing and hay operations may proceed with no restrictions.

For tank mixture applications, refer to the use directions and restrictions of the mixture product.

**When applied at 0.3 to 0.7 dry ozs. (0.008 to 0.016 pound active ingredient) per acre Aim Herbicide will provide:
Control of listed weeds up to 4 inches tall.**

| | |
|--------------------------------|-------------------|
| Bedstraw, catchweed | Nightshade, black |
| Fiddleneck, coast | |
| Flixweed | Pennycress, field |
| Lambsquarters (up to 3 inches) | Pigweed, redroot |
| Rocket, London | Velvetleaf |
| Mustard, tansy | Wallflower, bushy |
| Mallow, common | Cheeseweed |

Suppression of listed weeds up to 4 inches.

| | |
|------------------|------------------|
| Bindweed, field | Mustard, spp. |
| Filaree, redstem | Shepherdspurse |
| | Thistle, Canada |
| Kochia | Thistle, Russian |
| Lettuce, prickly | Wild buckwheat |

**When applied at 0.9 to 1.2 dry ozs. (0.023 to 0.031 pound active ingredient) per acre Aim Herbicide will provide:
Control of the following weeds up to 4 inches tall.**

| | |
|----------------------------------|--------------------|
| All weeds controlled above plus: | |
| Bittercross | |
| Buckwheat, wild | Pennycress, field |
| Filaree, Redstem | Pigweed, spp. |
| Amaranthus, spp | Nightshade, hairy |
| Kochia | Shepherdspurse |
| Lambsquarters | Sowthistle, annual |
| Mustard, tansy | Thistle, Russian |
| Mustard, tumble | Speedwell, ivyleaf |
| Meadowfoam | Spurry, com |

Tank Mixtures with other herbicides

Aim Herbicide may be tankmixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide. When tankmixing Aim Herbicide with other products, be sure the Aim Herbicide is mixed in the spray tank water first.

Restrictions:

Do not make applications less than 7 days apart.
Do not apply more than 3.8 dry ozs. (0.093 pound active ingredient) per acre per season.

HOPS

For Use In ID, OR AND WA Only

TIMING AND METHOD OF APPLICATION

Post-Directed Application For Sucker Management

Aim Herbicide is a contact herbicide for directed spray application to the basal portion of the hop plant for the management of sucker growth. Apply Aim Herbicide at 1.2 dry ozs. (0.03 pound active ingredient) per acre per application in a minimum of 20 gallons of spray solution by boom-type ground application equipment only to the basal portion of the hop plant (approximately the lower 1.5 feet) and to the sucker mat which extends from the base of the plant to approximately 1.5 to 2 feet into the row.

An alternate row treatment program may be followed to avoid the removal of excessive photosynthetic capacity from the crown area. When treating alternating rows on different days, the equivalent maximum rate must not exceed 0.7 dry ozs. (0.015 pound active ingredient) of Aim Herbicide per application per treated row area totaling 0.5 acres.

Coverage is essential to obtain good basal growth management. Use a nonionic surfactant (NIS) having at least 80 percent active ingredient at 0.25 % v/v (2 pints of NIS per 100 gallons of spray volume) or a quality crop oil concentrate (COC) at recommended rates.

If Aim Herbicide is used in a tank mixture, refer to the other product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

Postemergent Weed Control of Broadleaf Weeds

Aim Herbicide may be applied using shielded sprayers or hooded sprayers to control emerged and actively growing broadleaf weeds within or between the rows of the crop.

When applied at up to 1.2 dry ozs. (0.03 pound active ingredient) per acre Aim Herbicide will provide:
Control of the listed weeds up to 4 inches tall.

| | |
|-----------------------|----------------------|
| | Morningglory, pitted |
| Cheeseweed | Lettuce, prickly |
| Amaranth, Palmer | Nettle, burning |
| Burclover | Nettle, stinging |
| Cocklebur, common | Nightshade, spp. |
| Filaree, spp. | Pigweed, redroot, |
| Fiddleneck, coast | Pigweed, smooth |
| Lambsquarters, common | Redmaids |
| Rocket, London | Shepherdspurse |
| Mallow, common | Sowthistle |
| Morningglory, ivyleaf | Velvetleaf |

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

Precautions

Extreme caution must be taken during application to avoid upward drift of the spray solution and contact with the highly susceptible new growth. Avoid applications until newly trained vines have developed sufficient barking to avoid damage to the stem and are high enough up the string to avoid contact with the apical bud. Only use nozzles that will produce coarse or very coarse droplets of a Volume Median Diameter, VMD, greater than 350 microns. Do not exceed 30-psi spray pressure unless otherwise required by the manufacturer of drift reduction nozzles. Do not apply Aim Herbicide using air blast or air assisted sprayers or application devices.

Restrictions

Do not apply within 7 days of harvest.
Do not apply through any type of irrigation system.

Do not apply more than 4.8 dry ozs. (0.12 pound active ingredient) per acre per season.

Allow 14 days between treatments of Aim Herbicide.

Dealers Should Sell in Original Packages Only. Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent permitted by law FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC, Aim - trademarks of FMC Corporation
Accent, Accent Gold, Ally, Basis, Basis Gold, Express, Finesse, Harmony, - trademarks of E.I. DuPont de Nemours and Company
Achieve, Gramoxone, Touchdown, - trademark of ZENECA, Inc.
Amber, Beacon, Discover, Exceed, Equip, Northstar, Option, Peak, Spirit, Tough, - trademarks of Novartis
Assert - trademark of Helena Chemical Company
Lightning - trademark of American Cyanamid Company
Banvel, Clarity, Distinct, Marksman - trademarks of BASF Corporation
Curtail, Hornet, Scorpion, Starane - trademarks of Dow Agrosiences, LLC
Bronate, Hoelon, Liberty, Puma, Dropp, Finish, Prep, Folex, Ginstar - trademarks of Aventis Group
Laddok, Poast - trademarks of BASF Aktiengesellschaft
Permit - trademark of Nissan Chemical Industries, Inc.
Roundup, Roundup Ultra - trademarks of Monsanto Company
Salvo, Shotgun, Sword, - trademarks of Platte Chemical Company
Sencor, Def, Everest - trademarks of Bayer Aktiengesellschaft
Sterling, Bison - trademarks of Agrilience, LLC
Harvade - trademark of Uniroyal Chemical Company, Inc.
CottonQuik - trademark of Griffin, LLC

** In California only