

279-3241

09-21-2007

1/1

**SUPPLEMENTAL LABEL FOR THE USE OF AIM EC
HERBICIDE AS A HARVEST AID IN PEANUTS**



ACCEPTED
SEP 21 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

279-3241

**For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA**

EPA Reg. No. 279-3241 EPA Est. 279-

Active Ingredient:	By Wt.
Carfentrazone-ethyl	22.3%
Other Ingredients:	77.7%
	<u>100.0%</u>

This product contains 2.0 pounds active ingredient per gallon.
Contains Petroleum Distillates
U.S. Patent No. 5,125,958

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable precautionary statements, use directions and restrictions on the full Aim EC Herbicide label.

See Container Label for Statement of Practical Treatment and Other Precautionary Information.

All applicable directions, restrictions, Worker Protection Standard requirements and other precautions on the EPA-registered label are to be followed.

HARVEST AID TREATMENT

Aim EC may be applied to peanuts to defoliate and/or desiccate troublesome broadleaf weeds such as morningglories, pigweeds and velvetleaf that may be present at harvest. Aim EC may be used alone or as a tank mixture with other harvest aids.

Applications shall be made when the crop is mature and/or according to Extension Service recommendations in the use area.

Aim EC Use Rates

Apply Aim EC 1 to 2 fl. ozs. per acre, but not to exceed maximum labeled rates. If treatments of Aim EC have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment. (See restriction section below)

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

Adjuvant Recommendation

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre may be used in addition to the nonionic surfactant methylated seed oil or crop oil.

Coverage is essential for satisfactory performance.

Precaution

If applied as a tank mixture, refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

Restrictions

- Do not apply within 7 days of harvest.
- Do not apply more than one application at 2 fl. ozs. (0.031 pound active ingredient) per acre total as a harvest aid.
- Do not apply more than 6.1fl. ozs. (0.096 pound active ingredient) per acre total for preplant and in-season weed control.
- Do not graze or feed immature peanut plants or peanut hay to livestock.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Aim and FMC -FMC Trademarks
©2007 FMC Corporation