

279-3244

11-13-2001

146156-05816

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number.

279-3244

Date of

NOV 13 2001

Term of Issuance: Conditional
Registration until November
15, 2003

Name of Pesticide Product:

Capture 1.15 Bifenthrin

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

FMC Corporation
1735 Market Street
Philadelphia, Pennsylvania 19103

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered since you have agreed in writing to the following terms for conditional registration in accordance with FIFRA sec. 3(c)(7)(A):

1. You will submit and/or cite all data required for registration/reregistration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. The subject registration is conditioned under the same terms and condition for data generation as stipulated in our November 15, 1993 letter for use of this product on cotton and other crops.
3. FMC Corporation agreed that the current synthetic pyrethroid mitigation measures are interim in nature and may be reconsidered or modified after review and evaluation of the Spray Drift Task Force.

Signature of Approving Official:

Date:

- 4. You understand that this registration will expire on November 15, 2003. You further understand that it is EPA's stated intent to, by November 15, 2003, complete its review of all relevant data and other information that are available to the Agency, and to make FIFRA section 3(c)(5) or other appropriate regulatory decisions for cotton-use synthetic pyrethroids and other crops conditionally registered based on the Agency's review of such data/information and considering statutory and regulatory criteria for such decisions.
- 5. You will submit product information (pounds or gallons produced) for this product for the fiscal year in which the labeled crop uses are conditionally registered, in accordance with FIFRA section 29. The fiscal year begins October 1 and ends September 30. The production information will be submitted to the Agency no later than November 15 following the end of the preceding fiscal year.

The information should be submitted to:

U.S. Environmental Protection Agency
 Office of Pesticide Programs (7504C)
 Document Processing Desk
 Ariel Rios Building
 1200 Pennsylvania Avenue, NW
 Washington, DC 20460

- 6. You will make the following label changes before releasing the product for shipment.
 - a. Add the phrase EPA Registration Number 279-3244.
 - b. Make the New York State restriction the last paragraph under Application Instructions.
 - c. Under the Remarks section for "at plant" use for both field and sweet corn add,

"Granules must be incorporated into the top one inch of soil using tines, chains or other suitable equipment. Do not apply as a T-band treatment unless you can incorporate the granules."
 - d. Under the Remarks section for field and sweet corn foliar use add the following,

Apply by aircraft broadcasting over corn plants or with ground equipment by directing the granule into the whorl.

You should also note that, regardless of whether you satisfy all applicable conditions, this **conditional** registration will expire automatically on November 15, 2003. Sale and distribution of the subject product bearing labeling for subject uses after November 15, 2003 will be illegal.

Finally once the required data have been submitted and evaluated, EPA will entertain an application to amend the registration on the subject product to allow the unconditional use on cotton and other crops without any special restrictions on the duration of the registration.

A copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505)

Enclosure

4/8

RESTRICTED USE PESTICIDE
Toxic to fish and aquatic organisms.
For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.



Insecticide/Miticide

EPA Reg. No. 279-XXXX

EPA Est. 279-

Active Ingredient:

By Wt.

Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropanecarboxylate* 1.15%

Inert Ingredients:..... 98.85%
100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.
U.S. Patent No. 4,238,505

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician:

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

For Emergency Assistance Call (800) 331-3148.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing.

Personal Protective Equipment:

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. See other panels for additional precautionary statements.

User Safety Recommendations
Users should:
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

Physical/Chemical Hazards

Do not use or store near heat or open flame



FMC Corporation
Agricultural Products Group
Philadelphia PA 19103

Capture1 15G_11_1_2001

.....
.....
.....
.....
.....

ACCEPTED
with COMMENTS
in EPA Letter Dated

NOV 13 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 279-3244

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. For applications to sites with restricted entry intervals (REIs) of 7 days or more, notify workers of applications by warning them orally and by posting warning signs at entrances of treated areas.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) listed in the chart below:

Crop	Activity	REI
Eggplant	All	9 days
Beans (for processing only)	All	24 hours
Beans (for fresh market)	All	9 days
Peas (for processing only)	All	24 hours
Peas (for fresh market)	All	9 days
Head, Stem Brassica	All	5 days
Sweet Corn (for processing only)	All	24 hours
Sweet Corn (for fresh market)	All	18 days
Field Corn (Grown for seed)	Detasselling	18 days
Field Corn (Grown for seed)	Roguing	18 days
Field Corn (Grown for seed)	Other Activities	24 hours
Other Crops	All	24 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants, socks and shoes and waterproof gloves.

STORAGE AND DISPOSAL

Pesticide Storage

Do not store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Container: Do not rinse container. Return empty container to point of purchase.

Rotational Crops

Leafy vegetables and root crops may be rotated 30 days following the final application of bifenthrin. Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated seven months following the final application of bifenthrin. Straw may not be used for food or feed.

Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.


Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

Runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not apply this product within 20 yards of water (lakes, reservoirs, rivers, permanent streams, marshes, estuaries, natural ponds, and commercial fish farm ponds). Cover ponds to rate spill including end row spillage.

Use of Capture 1.15G is prohibited in all coastal counties.

Resistance

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details. If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local commodity representative or FMC staff for the best alternative method of control for your area.



6/8

FIELD CORN (GRAIN, SILAGE, AND SEED), POPCORN (AT PLANT USE)		
PEST	DOSAGE	REMARKS
Corn Rootworm Larvae Mexican Northern Southern Western Cutworms Wireworms Seed Corn Maggot Seed Corn Beetle White Grubs	6.4 to 8.0 oz product (0.005 - 0.006 lb ai) per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position in front of the press wheel centered over the row. Use the table below to determine the Capture 1.15G needs per acre. Use highest use rate under heavy insect pressure.
Cutworms Wireworms Seed Corn Maggot Seed Corn Beetle White Grubs	3.2 to 8.0 oz product (0.002 - 0.006 lb ai) per 1,000 linear feet of row	Apply as an in-furrow application directly into the seed furrow. Use highest use rate under heavy insect pressure.
Rootworm Larvae	8.0 oz product (0.006 lb ai) per 1,000 linear feet of row	In-furrow for light to moderate pressure

ounces product per 1000 linear feet of row	Capture 1.15G lbs product/acre (lbs ai/acre) based on row spacing					
	30"	32"	34"	36"	38"	40"
8.0	8.7 (0.100)	8.2 (0.094)	7.7 (0.088)	7.2 (0.083)	6.9 (0.079)	6.5 (0.075)
6.4	7.0 (0.080)	6.5 (0.075)	6.1 (0.071)	5.8 (0.067)	5.5 (0.063)	5.2 (0.060)
4.8	5.2 (0.060)	4.9 (0.056)	4.6 (0.053)	4.3 (0.050)	4.1 (0.047)	3.9 (0.045)
3.2	3.5 (0.040)	3.3 (0.038)	3.1 (0.035)	2.9 (0.033)	2.7 (0.032)	2.6 (0.030)

APPROXIMATE SETTINGS BASED ON SPEED - CALIBRATE FOR EXACT SETTINGS										
Planter type	4 mph		5 mph		6 mph		7 mph		8 mph	
	6.4 oz	8.0 oz								
JD Max 7200-7300	17	20	20	26	22	30	26	35	30	40
JD Max 7000-7100 odd notch	11	14	13	16	15	18	17	19	19	21
JD Max 7000-7100 even notch	15	17	18	20	22	24	24	26	27	29
Kinzi even notch	13	16	16	19	19	21	20	22	22	24
Kinze odd notch	11	13	13	15	15	17	17	19	19	21
White 6000 series	12	14	15	17	17	20	19	22	21	24
Case-IH 800-900 series	1/7	1/9	2/1	2/3	2/4	2/6	2/6	2/9	2/8	3/2

Always calibrate equipment prior to application each season to insure proper insecticide delivery.
 Utilize initial settings from the chart below or for clay granule insecticide delivery in the appropriate planter guide booklet.
 Steps for calibration:
 1. Fill insecticide hopper box
 2. Attach a Capture calibration device to delivery tube opening
 3. Set planter to initial setting
 4. Measure off 1,000 row feet and drive your planter the pre-measured distance at your desired planting speed
 5. Check the level in the calibration device and it should contain the number of ounces (3.2-8.0) required for your application
 6. If a row is over or under the desired rate adjust the settings and repeat the calibration

FIELD CORN (GRAIN, SILAGE, AND SEED), POPCORN (FOLIAR USE)		
PEST	DOSAGE lb product/acre (lb ai/a)	REMARKS
European corn borer (1st generation) Fall armyworm Southwestern corn borer Armyworm	3.5- 8.7 (0.04 - 0.10)	Apply as a broadcast application when insects first appear. Applications may be made by with ground or aerial equipment. Use higher labeled use rates under heavy insect infestations.

RESTRICTIONS—Corn
 Do not apply to soil where there is greater than 30% cover of crop residue remaining.
 Do not apply more than 0.1 pound active per acre per season as an at plant application
 Do not apply more than 0.3 pound active per acre per season including at plant plus foliar applications.
 Do not apply within 30 days of harvest.
 Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
 Do not make aerial or ground applications to corn if heavy rainfall is imminent.

SWEET CORN (AT PLANT USE)

PEST	DOSAGE	REMARKS
Corn Rootworm Larvae Mexican Northern Southern Western Cutworms Wireworms Seed Corn Maggot Seed Corn Beetle White Grubs	6.4 to 8.0 oz product (0.005 - 0.006 lb ai) per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position in front of the press wheel centered over the row. Use the table below to determine the Capture 1.15G needs per acre. Use highest use rate under heavy insect pressure.
Cutworms Wireworms Seed Corn Maggot Seed Corn Beetle White Grubs	3.2 to 8.0 oz product (0.002 - 0.006 lb ai) per 1,000 linear feet of row	Apply as an in-furrow application directly into the seed furrow. Use highest use rate under heavy insect pressure.
Rootworm Larvae	8.0 oz product (0.006 lb ai) per 1,000 linear feet of row	In-furrow for light to moderate pressure

ounces product per 1000 linear feet of row	Capture 1.15G lbs product/acre (lbs ai/acre) based on row spacing					
	30"	32"	34"	36"	38"	40"
8.0	8.7 (0.100)	8.2 (0.094)	7.7 (0.088)	7.2 (0.083)	6.9 (0.079)	6.5 (0.075)
6.4	7.0 (0.080)	6.5 (0.075)	6.1 (0.071)	5.8 (0.067)	5.5 (0.063)	5.2 (0.060)
4.8	5.2 (0.060)	4.9 (0.056)	4.6 (0.053)	4.3 (0.050)	4.1 (0.047)	3.9 (0.045)
3.2	3.5 (0.040)	3.3 (0.038)	3.1 (0.035)	2.9 (0.033)	2.7 (0.032)	2.6 (0.030)

APPROXIMATE SETTINGS BASED ON SPEED - CALIBRATE FOR EXACT SETTINGS

Planter type	4 mph		5 mph		6 mph		7 mph		8 mph	
	6.4 oz	8.0 oz								
JD Max 7200-7300	17	20	20	26	22	30	26	35	30	40
JD Max 7000-7100 odd notch	11	14	13	16	15	18	17	19	19	21
JD Max 7000-7100 even notch	15	17	18	20	22	24	24	26	27	29
Kinzi even notch	13	16	16	19	19	21	20	22	22	24
Kinze odd notch	11	13	13	15	15	17	17	19	19	21
White 6000 series	12	14	15	17	17	20	19	22	21	24
Case-IH 800-900 series	1/7	1/9	2/1	2/3	2/4	2/6	2/6	2/9	2/8	3/2

Always calibrate equipment prior to application each season to insure proper insecticide delivery.
 Utilize initial settings from the chart below or for clay granule insecticide delivery in the appropriate planter guide booklet.
 Steps for calibration:
 1. Fill insecticide hopper box
 2. Attach a Capture calibration device to delivery tube opening
 3. Set planter to initial setting
 4. Measure off 1,000 row feet and drive your planter the pre-measured distance at your desired planting speed
 5. Check the level in the calibration device and it should contain the number of ounces (3.2-8.0) required for your application
 6. If a row is over or under the desired rate adjust the settings and repeat the calibration

SWEET CORN (FOLIAR USE)

PEST	DOSAGE lb product/acre (lb ai/a)	REMARKS
European corn borer (1st generation) Fall armyworm Southwestern corn borer Armyworm	3.5- 8.7 (0.04 - 0.10)	Apply as a broadcast application when insects first appear. Applications may be made by with ground or aerial equipment. Use higher labeled use rates under heavy insect infestations.

RESTRICTIONS—Sweet Corn
 Do not make more than 0.2 pounds active ingredient per acre per season.
 Do not apply within one day of harvest.
 Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application. Do not apply within 14 days of harvest for corn that is hand harvested.
 Do not make aerial or ground applications to corn if heavy rainfall is imminent.

CUCURBITS:

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucurza) (Luffa spp.) (includes hechima, Chinese okra) (Momordica spp.) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin (Cucurbita spp.); Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (includes butternut squash, calabaza, hubbard squash (C. mota, C. pepo) includes acorn squash, spaghetti squash); Watermelon (includes hybrids and/or varieties of Citrullus spp.)

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not apply more than 0.3 lb active ingredient per acre per season.
Do not make more than two applications after bloom.
Do not make applications less than 7 days apart.
Do not apply within 3 days of harvest.

SUCCULENT PEAS AND BEANS:

Pea (Pisum spp.): Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea, Pigeon pea; Bean (Phaseolus spp.): Broadbean (succulent), Lima bean (green), Runner bean, Snap bean, Wax bean; Bean (Vigna spp.): Asparagus bean, Blackeyed pea, Chinese longbean, Cowpea, Moth bean, Southern pea, Yardlong bean, Jackbean, Soybean (immature seed), Sword bean

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present. Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not apply more than 0.2 lb. active ingredient per acre per season.
Do not apply within 3 days of harvest.

BRASSICAS:

Head and Stem Brassica Vegetables including: Broccoli, Chinese Broccoli (gai lan, white flowering broccoli), Brussels Sprouts, Cauliflower, Cavalo broccolo, Kohlrabi, Cabbage, Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy)

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not apply more than 0.5 lb. active ingredient per acre per season.
Do not make more than 5 applications after bloom
Do not make applications less than 7 days apart.
Do not apply within 7 days of harvest

LETTUCE, HEAD:

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present. Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not make applications less than 7 days apart.
Do not apply more than 0.20 pound active ingredient per acre per season.
Do not apply within 7 days of harvest.

EGGPLANT:

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not make applications less than 7 days apart.
Do not apply more than 0.2 lb active ingredient per acre per season.
Do not apply within 7 days of harvest.

COTTON:

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not apply more than 0.5 pound active per acre per season
Do not apply within 14 days of harvest.
Do not graze livestock in treated areas or cut treated crops for feed.
Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush[®], Ammo[®], Asana[®] XL, Baythroid[®], Capture[®], Danitol[®], Karate[®], Mustang[®], and Scout X-TRA[®].
This product is not for sale or use in the State of California.

PEPPERS, BELL AND NON-BELL:

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	3.5- 8.7 (0.04 - 0.10)	Make application of Capture as soon as pests are present Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not make applications less than 7 days apart.
Do not apply more than 0.20 pound active ingredient per acre per season.
Do not apply within 7 days of harvest.

CANOLA, CRAMBE, RAPESEED:

PEST	DOSAGE lb product/acre (lb a/a)	REMARKS
Wireworms Grubs Seed Maggots Cutworms	2.7 - 3.5 (0.033 - 0.04)	Make application of Capture as soon as pests are present. Thorough coverage is essential to achieve control.

RESTRICTIONS
Do not apply more than 0.08 lb active ingredient per acre per season.
Do not make applications less than 14 days apart.
Do not apply within 35 days of harvest.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

•FMC, Ammo, Capture, Mustang, U-Turn—trademarks of FMC Corporation

© 2001 FMC Corporation. All rights reserved.