

279-3246

09-19-2007

1/7

ACCEPTED

SEP 19 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

279-3246



For Use Only by Individuals/Firms Certified as Licensed Pesticide Applicators

EPA Reg. No. 279-3246

EPA Est.

FROM FMC

Table with 2 columns: Active Ingredient, Percentage. Includes Sulfentrazone* (62.1%), Cloransulam-methyl* (7.9%), and Other Ingredients (30.0%).

*Authority First DF Herbicide contains 0.7 pounds of active ingredient per pound of product (0.62 pounds a.i. of sulfentrazone and 0.08 pounds a.i. of cloransulam-methyl)

U.S. Patent No. 4,818,275 and 5,163,995

KEEP OUR OF REACH OF CHILDREN

CAUTION-AVISO

Si usted etiene esta etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID (2)

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER (3)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

ATTENTION (4)
-Although this label may appear similar to the label on a product you may have used, there may be important label differences. Users must read, understand and strictly follow all label directions, precautions and restrictions.
-It is the user's responsibility to be sure the product is approved for sale or use on the intended crop and for use in the specific geographic area.
-It is the user's responsibility to be aware of and to follow all State or local precautions or restrictions not appearing on this product label.
-Prior to purchase or use of this product, read the Conditions of Sale and Limitation of Warranty and Liability on page 2 of this label. If the terms and conditions are unacceptable, return the product immediately in the original and unopened container.

TABLE OF CONTENTS

Table with 2 columns: Section Title/ (Section Number), Page. Lists sections like Active Ingredient (1), Agricultural Use Requirements (9), etc.

2/7

PRECAUTIONARY STATEMENTS (5) Hazards to Humans and Domestic Animals

Caution

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE) (6)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards (7)

This pesticide is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface Water Advisory

Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

Physical/Chemical Hazards (8)

Do not use or store near heat or open flame.

AGRICULTURAL USE REQUIREMENTS (9)

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. (10)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Conditions of Sale and Limitation of Warranty and Liability (11)

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and, to the extent permitted by applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent permitted by applicable law, buyer assumes the risk of any such use.

To the extent permitted by applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Storage and Disposal (12)

Do not contaminate water, food or feed by storage or disposal.

Pesticide storage

Store product in original container only, away from other pesticides, fertilizer, food or feed.

Do not store below 40F or 4C degrees.

In Case of Spill

Avoid contact. Isolate areas and keep out animals and unprotected persons.

To Confine Spills.

Dike surrounding area, sweep up spillage, Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a large holding container. Identify contents per required hazardous waste labeling regulations.

3/7

Storage and Disposal (continued)(12)

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes can not be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Metal Containers

Metal containers should be triple rinsed (or equivalent). Then offered for recycling or reconditioning, or punctured and disposed of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

Plastic Containers

Plastic containers should be triple rinsed (or equivalent). Then offered to an approved pesticide container recycling program. Or punctured and disposed of in an approved waste disposal facility. Provided on-site incineration is allowed by State and local authorities, containers may be burned. Stay out of smoke.

Returnable/Refillable Sealed Containers

Do not break container seals. Do not empty remaining formulated product. Do not rinse inside of container. Return container intact to point of purchase.

RESISTANCE MANAGEMENT (13)

Some weeds are known to develop resistance to herbicides that have been used repeatedly. While the development of resistance is well understood, it is not easily predicted. Therefore herbicides should be used in conjunction with resistance management strategies in the area. Consult the local or State agricultural advisors for details. If weed resistance should develop in the area, this product used alone may not continue to provide sufficient levels of weed control. If the reduced levels of control can not be attributed to improper application timing, unfavorable weather conditions or abnormally high weed pressure, a resistant strain may have developed.

To reduce the potential for weed resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the recommended rates and in accordance with the use directions. Do not use less than recommended label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when weeds are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

DIRECTIONS FOR USE (14)

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION (15)

Authority First DF Herbicide is for preemergence control of broadleaf and grass weeds in soybeans only.

The mode of action of Authority First DF Herbicide involves uptake by weed roots and shoots. Preemergence and preplant incorporated applications of Authority First DF Herbicide require rainfall or irrigation to activate the herbicide. The amount of rainfall or irrigation required for activation following application depends on existing soil moisture, organic matter content and soil texture. If adequate moisture (1/2" to 1") is not received within 7 to 10 days after the Authority First DF treatment, a shallow cultivation may be needed to obtain desired weed

control. When sufficient moisture is received after dry conditions, Authority First DF Herbicide will provide control of susceptible germinating weeds.

Authority First DF Herbicide exhibits excellent crop safety. Poor growing conditions, such as excessive moisture, cool temperatures, and soil compaction or the presence of various pathogens may impact seedling vigor. Under these conditions, the active ingredients in Authority First DF, like other soil-applied herbicides, can contribute to crop response. However, these early symptoms are short-lived.

Observe all instructions, crop restrictions, mixing directions, application precautions, replanting directions, rotational crop guidelines and other label information of each product when tank mixing with Authority First DF.

Proper Handling Instructions

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Do not apply this product through any type of irrigation system.

Do not use flood irrigation to apply or incorporate this product.

Product must be used in a manner which will prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

ROTATIONAL CROP GUIDELINES (16)

Shown below are the minimum intervals in months from the time of Authority First DF Herbicide application until Authority First DF treated soil may be replanted with the crops listed. When Authority First DF Herbicide is tank mixed with other herbicide(s), refer to all the labels for re-cropping instructions, following the intervals that are the most restrictive. For crops not listed, the interval is 30 months in addition to a successful field bioassay.

CROP	INTERVAL (Months)
Alfalfa	12
Barley	12
Canola	24
Corn, field *	10
Corn, pop *	10
Corn, seed *	10
Corn, sweet	18
Cotton	18
Dry shelled beans and peas	12
Oats	12
Peanuts	12

4/7

Potatoes	18
Rice	10
Rye	12
Sorghum	12
Soybeans	Anytime
Sugarbeets **	30
Sunflower **	30
Tobacco ***	30
Wheat	4

- Do not use tail water from the first flood or furrow irrigation of treated fields to treat non target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

SPRAY DRIFT REDUCTION ADVISORY (19)

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops.

Where States and local governments have more stringent regulations, they must be observed.

Droplet Size Information

Reduce drift potential by applying large droplets. The optimum drift management strategy is to apply the largest droplets that will provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift when applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity and Temperature Inversions).

VMD – VMD is the expression of the droplet size of the spray cloud. The VMD value means that 50% of the droplets are larger than the expressed value and 50% of the droplets are smaller than the expressed value. Optimum spray clouds should be 450 microns with fewer than 10% of the droplets being 200 microns or smaller.

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows usually produce larger droplets.

Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types narrower spray angles produce larger droplets. Consider using low drift nozzles.

Application Height – Making applications at the lowest height practical reduces exposure of spray droplets to evaporation and wind movement.

Swath Adjustment – Swath adjustment distance must increase with increasing drift potential (higher wind, smaller droplets, etc.)

Wind – Drift potentials are lowest between wind speeds of 3 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications in wind conditions outside of this range could increase the risk of off-target effects and should be avoided. Note that local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity - When making applications in conditions of low relative humidity set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions – Do not apply Authority First during temperature inversions because the drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the following morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or a smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicate an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

* Corn (including field corn, popcorn and seed corn)
Observe an 18 month rotational interval if 6.45 – 8.0 oz. of Authority First DF Herbicide was applied to soils of 1.5% organic matter or less, and pH is above 7.

Hybrid Seed Production

Corn inbred lines grown for hybrid seed production may be injured in the growing season following an application of Authority First DF Herbicide. Inbred lines should be thoroughly tested for crop tolerance before rotating to production scale acreages. FMC will not accept responsibility for any crop injury on field corn grown for seed following an application of Authority First DF Herbicide.

** These crops require a 30-month rotational interval and a successful field bioassay.

*** Transplanted tobacco may be planted 10 months after application of a maximum application rate of 3.0 ounces per acre of Authority First. Tobacco in seedbed nurseries may be replanted 18 months after applications of 3.0 ounces per acre of Authority First and following a successful field bioassay. A rotational interval of 30 months and a successful field bioassay is required for all applications of Authority First greater than 3.0 ounces per acre.

REPLANTING INSTRUCTIONS (17)

If the initial planting of soybeans fails to produce a uniform stand, soybeans may be replanted in fields treated with Authority First DF Herbicide alone. Do not retreat fields with a second application of Authority First DF Herbicide. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP GUIDELINES on the Authority First DF Herbicide label. Where a tank mix is used, refer to the product's labels for any additional replant instructions.

APPLICATION INFORMATION (18)

DO NOT APPLY TO CROPS OTHER THAN SOYBEANS. APPLY WITH GROUND SPRAYERS ONLY

Use a standard low pressure herbicide boom sprayer equipped with suitable nozzles and screens. Apply uniformly using properly calibrated nozzles (10 to 40 psi) and screens and strainers no finer than 50 mesh. Use 10 to 40 gallons of spray solution per acre. Do not exceed 40 psi spray pressure unless required by the spray nozzle manufacturer.

Continuous agitation is required until all spray mixture has been applied. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow Authority First DF spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.

To avoid injury to sensitive crops, spray equipment used for Authority First DF Herbicide applications must be drained and thoroughly cleaned with water plus ammonia before being used to apply other products. See Spray Clean-out Section 21 on page 5.

Avoid all direct, and/or indirect spray contact with non-target plants. Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure.

RUNOFF AND WIND EROSION PRECAUTIONS

Do not apply under conditions which favor runoff or wind erosion of soil containing Authority First DF to non-target areas.

To prevent off-site movement due to runoff or wind erosion:

- Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, allow the soil surface to be settled by rainfall or irrigation.
- Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered ground.
- Do not apply to soils when saturated with water.

5/7

Sensitive Areas – Applications should be made when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species and non-target crops).

MIXING AND LOADING INSTRUCTIONS (20)

Authority First DF Herbicide Applied Alone

Select the proper Authority First DF Herbicide application rate from the following TIMING AND METHOD OF APPLICATION section of this label. Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of Authority First DF Herbicide for acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Apply the Authority First DF Herbicide spray mixture immediately after mixing. Do not store spray mixture.

Authority First DF Applied in Tank Mix Combination

Select the proper Authority First DF Herbicide application rate from TIMING AND METHOD OF APPLICATION section of label. Read and follow all applicable use directions, precautions and restrictions on the respective tank mix product labels. To ensure product compatibility, a jar test should be conducted before large volume mixing. Provided the jar test indicates the mixture is compatible, prepare the tank mixture as follows.

Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of Authority First DF Herbicide for the acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Next, add the recommended amount(s) of the additional tank mix product(s) in the following order: first dry formulations (e.g., wettable powders, dry flowables), next liquid suspensions (e.g., flowables) and finally liquids (e.g., EC's). Allow time for complete mixing and dispersion after each addition, adding water as necessary. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Use Authority First DF Herbicide tank mixtures immediately after mixing. Do not store tank mixtures.

Fertilizer Spray Mixtures

Applications of Authority First DF Herbicide alone, or with recommended tank mixtures, in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing in full spray tank quantities.

- Put 1 pint of fertilizer solution in a quart jar.
- Add the appropriate amount of herbicide based on the table below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

Herbicide Type	Herbicide Field Use Rate	Amount Herbicide Added Per Pint
Wettable Powder or Dry Flowable	0.5 pound	0.75 teaspoon
	1.0 pound	1.50 teaspoons
	2.0 pounds	3.00 teaspoons
	3.0 pounds	4.50 teaspoons
Emulsified Concentrates	1.0 pint	0.5 teaspoon
	1.0 quart	1.0 teaspoon
	2.0 quarts	2.0 teaspoons
	3.0 quarts	3.0 teaspoons

*Based on a spray volume of 25 gal. per acre. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- Close jar and shake well.
- Watch mixture for several seconds, again after 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below.

Authority First DF Herbicide Applied Alone with Liquid Fertilizer

In order to add Authority First DF Herbicide to a liquid fertilizer carrier, Authority First must be premixed in a slurry of product and clear water. Fill the spray tank one-half full with fertilizer solution. With agitator operating, add the Authority First DF Herbicide slurry to the spray tank. Use a minimum of one gallon of water for each container of Authority

First DF Herbicide. Stir until completely dissolved. Then add slurry to the spray tank through a 20-35 mesh screen. Rinse container used for pre-mixing and add rinsate to the spray tank. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use Authority First DF Herbicide spray mixture immediately after mixing. Do not store mixture.

Authority First DF Herbicide Applied in Tank Mix Combinations

Fill the spray tank one-half full with fertilizer solution. With the agitator operating, add a slurry of Authority First DF Herbicide as described in the preceding paragraph. Next, dilute the individual tank mix partners with sufficient water to form a free flowing dispersion, then add to the spray tank of fertilizer. While maintaining agitation, add the other products using the following order: slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use Authority First DF Herbicide tank mixtures immediately after mixing. Do not store tank mixtures.

SPRAY EQUIPMENT CLEAN-OUT (21)

After spraying Authority First DF Herbicide and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure:

- Drain sprayer tank, hoses, and spray boom and thoroughly rinse the inside of the sprayer tank with clean water to remove sediment and residues. Thoroughly flush sprayer hoses, boom and nozzles with clean water.
- Fill the tank 1/2 full with clean water, and add appropriate detergent or ammonia (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom, and nozzles.
- Convenient and thorough cleaning of the sprayer can be achieved if the cleaning solution is left in the spray tank; hoses, spray booms and spray nozzles overnight or during storage.
- Before using the sprayer, drain the spray system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately with the detergent or ammonia solution.
- Properly dispose of all cleaning solution and rinsate in accordance with Federal, State and local regulations and guidelines.

Do not drain or flush equipment on or near desirable trees or plants. Do not contaminate any body of water including irrigation water that may be used on other crops.

Should small quantities of Authority First remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

SOYBEANS (Conventional and GMO) (22)

TIMING AND METHOD OF APPLICATION

Authority First may be used alone or in tank mixture combinations for the control of the weeds listed in conventional or GMO soybean varieties.

Application Rates

Soil Organic Matter*	Authority First DF 70DF (Dry Ounces per Acre)**	Product Use Rates (Pound Active Ingredient Per Acre)	
		Sulfentrazone	Cloransulam-methyl
3% or less	6.45	0.25	0.032
Greater than 3%	8.00	0.31	0.040

*Do not apply Authority First DF Herbicide to soils classified as sand with less than 1% organic matter.
 ** Maximum application rates: See Preplant Surface and Preemergence Application below for specific recommendations.

Preplant Incorporated Application

Apply Authority First DF Herbicide alone or in tank mix combination with other herbicides registered for preplant incorporated application to soybeans. Incorporate the herbicide(s) into the top 1 to 3 inches of the final seedbed using equipment that provides thorough soil mixing.

6/7

When Authority First DF Herbicide is applied in tank mix combination with other herbicide(s), follow the incorporation directions for the tank mix partner(s). Follow applicable use instructions, including application rates, precautions and restrictions of each product used in the tank mixture.

Preplant Surface Application

Apply Authority First DF Herbicide alone or in tank mix combination with other herbicides registered for preplant soil surface application to soybeans. If applied in tank mix combination, follow use instructions, including application rates (note: apply 1/2 of the maximum application rate for suppression of weeds in Roundup Ready soybeans, maintaining control with sequential application(s) of registered postemergence herbicides), precautions and restrictions of each product used in the tank mixture.

Preemergence Application

Apply at planting time or within 3 days after planting. Authority First DF Herbicide may be applied alone or in tank mix combination with other herbicides registered for preemergence application to soybeans. When applied in a tank mix combination, follow applicable use instructions, including application rates (note: apply 1/2 of the maximum application rate for suppression of weeds in Roundup Ready soybeans, maintaining control with sequential application(s) of registered postemergence herbicides). Observe the precautions and restrictions of each product used in the tank mixture. Apply before planting, at planting time or prior to seed germination. Properly closed seed furrows are necessary when applying at planting time of before seed germination. Do not apply later than 3 days after planting (or after seed germination), as crop injury may result.

Weeds Controlled (23)

When used as directed, Authority First DF Herbicide will provide control or suppression of the following broadleaf weeds and grasses.

Smartweed, PA	<i>Polygonum pensylvanicum</i>
Smellmelon	<i>Cucumis melo</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
Grasses	
Barnyardgrass*	<i>Echinochloa crus-galli</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, southern*	<i>Digitaria ciliaris</i>
Crowfootgrass*	<i>Dactyloctenium aegyptium</i>
Foxtail, giant*	<i>Setaria faberi</i>
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, yellow*	<i>Setaria lutescens</i>
Goosegrass	<i>Eleusine indica</i>
Johnsongrass, seedling *	<i>Sorghum halapense</i>
Orchardgrass	<i>Dactylis glomerata</i>
Panicum fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Sedges	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

* Provides suppression or partial control only
 ** Will not control ALS resistant biotypes of these weed species

Reduced Rates for Roundup Ready Soybeans

Authority First may be used at reduced rates in conjunction with planned follow-up weed control applications with glyphosate based products such as Glyphomax XRT, Durango, Roundup or other glyphosate herbicide products labeled for use on Roundup Ready soybean varieties. Follow all Authority First application directions. Authority First will not control or reduce competition from ALS resistant biotypes.

Apply before planting, at planting time or prior to seed germination. Properly closed seed furrows are necessary when applying at planting time of before seed germination. Recommended postemergence treatments may include any product of combination of products labeled for use.

Reduced Rate Application Rates

Soil Organic Matter*	Authority First DF 70DF (Dry Ounces per Acre)	Product Use Rates (Pound Active Ingredient Per Acre)	
		Sulfentrazone	Cloransulam-methyl
3% or less	3.00	0.116	0.015
Greater than 3%	4.00	0.155	0.020

*Do not apply Authority First DF Herbicide to soils classified as sand with less than 1% organic matter.

Common Name	Scientific Name
Broadleaves	
Amaranth, Palmer	<i>Amaranthus, Palmer</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Anoda, spurred	<i>Anoda cristata</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Carpetweed	<i>Mollugo verticillata</i>
Cocklebur, common	<i>Xanthium Pensylvanicum</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryaeifolia</i>
Croton, tropic	<i>Croton glandulosus</i>
Daisy, American	<i>Eclipta alba</i>
Dayflower, common	<i>Commelina communis</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Groundcherry, clammy	<i>Physalis heterophylla</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Horseweed (Marestail)*	<i>Hippuris vulgaris</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Mexicanweed	<i>Caperonia castanaefolia</i>
Morningglory, entireleaf	<i>Ipomea hederacea integrisc</i>
Morningglory, ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, palmleaf	<i>Ipomea Wrightii</i>
Morningglory, pitted *	<i>Ipomea, lacunosa</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Mustard, wild	<i>Brassica kaber</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Poorjoe	<i>Diodia teres</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, common **	<i>Ambrosia artemisiifolia</i>
Ragweed, giant **	<i>Ambrosia trifida</i>
Senna, coffee	<i>Cassia occidentalis</i>
Teaweed	<i>Sida, prickly</i>

Preplant Burndown Application

Authority First DF Herbicide, used at 6.45 – 8.0 oz./acre as in the For All Soybeans Application Rates chart above, provides burndown of weeds listed below, when applied as follows. Thorough coverage is essential. Apply a minimum of ten gallons per acre finished spray volume. Use a non-ionic surfactant (NIS) having at least 80% active ingredient strength at 0.125-0.25% v/v (1-2 pints per 100 gallons of spray solution) plus ammonium sulfate (AMS) at 2.5% v/v. Crop oil concentrate (COC) at 1.2% v/v plus ammonium sulfate may be used. Burndown results may be slowed or reduced when the growth of the weeds is affected by unusual environmental factors just prior to or after application such as especially cool or widely fluctuating day and night air temperatures, drought, heat stress, or waterlogged soils. To broaden the spectrum of weeds controlled, Authority First DF

Herbicide may be tank mixed with other herbicides such as Aim EW Herbicide, 2,4-D products, glyphosate products, or paraquat products.

Weeds Controlled

When used as directed for burndown, Authority First DF Herbicide will provide control or suppression of the following broadleaf weeds.

Common Name	Scientific Name
Broadleaves	
Cocklebur, common	<i>Xanthium Pensylvanicum</i>
Horseweed (Marestail) * *	<i>Hippuris vulgaris</i>
Jimsonweed	<i>Datura stramonium</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Morningglory, entireleaf	<i>Ipomea hederacea tinegrisc</i>
Morningglory, ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, palmleaf	<i>Ipomea Wrightii</i>
Morningglory, pitted *	<i>Ipomea lacunosa</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea purpurea</i>
Ragweed, common * *	<i>Ambrosia artemisiifolia</i>
Ragweed, giant * *	<i>Ambrosia trifida</i>
Sicklepod	<i>Cassia obtusifolia</i>
Smartweed, PA	<i>Polygonum pensylvanicum</i>
Sunflower, common	<i>Helianthus annuus</i>
Velvetleaf *	<i>Abutilon theophrasti</i>

*For Velvetleaf control, use 28% nitrogen (UAN) or AMS with NIS or COC.
 * * Authority First will not control ALS resistant biotypes of these weed species

Precautions

Properly closed seed furrows are necessary when applying at planting time of before seed germination.
 Maintain spray tank agitation until the spray mixture is applied.

Restrictions (24)

Do not apply this product by air or through any type of irrigation system.
 Do not make more than one soil application per crop year.
 Do not apply more than 8.0 ounces of Authority First DF 70DF per acre per season (as a cumulative total of Authority First DF Herbicide soil application and FirstRate post-emergence application (1 oz/acre Authority First DF Herbicide contains 0.005 lb a.i. cloransulam-methyl; and 0.3 oz/acre FirstRate contains 0.016 lb a.i. cloransulam-methyl. Do not apply more than 0.055 lb. active ingredient (a.i.) per acre of cloransulam-methyl in a single growing season).
 Do not feed treated soybean forage or soybean hay to livestock.
 Do not harvest soybeans for 65 days after application of Authority First DF Herbicide.
 Do not apply Authority First DF Herbicide to soils classified as sands containing less than 1% organic matter.
 Do not drain or flush equipment on or near desirable trees or plants.
 Do not contaminate any body of water including irrigation water that may be used on other crops.

7/7

Authority, Authority First DF Herbicide and **FMC** — Trademarks of FMC Corporation, Philadelphia, PA 19103 USA
 FirstRate, Glyphomax XRT and Durango - Trademarks of Dow AgroSciences LLC, Indianapolis, IN 46268 USA.
 Roundup and Roundup Ready – Trademarks of Monsanto Technology LLC, St Louis, MO 63167 USA

LABEL TRACKING INFORMATON (25)

Label Code: AuthorityFirstDF_3_07-02-07

Replaces Label Code: Authority First 09-06

EPA Approval Date:

Revisions Include: Reformat
 Addition of Precautions and Restriction Section
 Addition of Reduced Rate Use
 Addition of Scientific Names for weeds
 Strengthen Spray Drift Section
 Added Table of Contents

FMC Corporation
 Agricultural Products Group
 1735 Market Street
 Philadelphia PA 19103
 215-299-6000

©2007 FMC Corporation All Rights Reserved