

# FMC 01-0004 Insecticide

For use by individuals/firms licensed or registered by the state to apply pesticide products. A ready to use insecticidal bait for application to the soil of flower and shrub beds, foundation perimeters, sports fields, parks, lawns and other ornamental turf for control of Mole Crickets.

EPA Reg. No. 279-3252

EPA Est. 279-

<b>Active Ingredient:</b>	<b>By Wt.</b>
Bifenthrin* .....	0.2%
Inert Ingredients .....	99.8%
	100.0%

\*Cis isomers 97% minimum, trans isomers 3% maximum.  
U.S. Patent No. 4,238,505

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See other panels for additional precautionary information.

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.	
NOTE TO PHYSICIAN	
This product is a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so should be avoided.	
For information Regarding the Use of this Product Call 1-800-321-1FMC (1362).	

## PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

**Caution**  
Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

### Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale of other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.4 lbs. A.I. (200 lbs. of this product) per Acre per year.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

## STORAGE AND DISPOSAL

**Prohibitions**  
Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage**  
Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**  
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal**  
Paper and plastic bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



FMC Corporation  
Agricultural Products Group  
Philadelphia PA 19103

**Net Weight**

**ACCEPTED**  
 JUN 18 2003  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 279-3252

272

## LAWN Applications

Broadcast this product with suitable application equipment to ensure uniform coverage over the treatment area. Apply the bait late in the afternoon to flower and shrub beds, lawns and other ornamental turf areas that have been well irrigated prior to application. For maximum efficacy, do not irrigate treated area immediately after application.

## APPLICATION RATES

The application rates listed in the following table will provide excellent control of Mole Crickets (Adults & Nymphs) under typical conditions. However, for residential lawn use, do not apply more than 0.2 lbs. A.I. per Acre per application.

Pest	Active Ingredient lbs. per acre	Application Rate FMC 01-0004 Insecticide	
		per acre	per 1000 sq. ft.
Mole Cricket (Adult) <sup>1</sup>	0.2 - 0.4	100 - 200 lbs.	2.3 - 4.6 lbs.
Mole Cricket (Nymph) <sup>1</sup>	lbs. A/A		

### Comments

<sup>1</sup>Mole Cricket: Achieving acceptable control of mole cricket nymphs and adults is best accomplished with a season-long integrated control program. A mole cricket management program should include monitoring the pest population and may include granular and/or liquid applications of Talstar® Insecticides and this product. Applications of this product may be followed by treatment with Talstar® Insecticides to control any individual mole crickets that may not have consumed the bait.

Tawny mole cricket nymphs will often be found infesting the same areas where late winter/early spring adult activity has been noted. Southern mole cricket adults will fly and breed in a more widespread area. The best time to control southern and tawny mole crickets with this bait is between June and December, depending on the geographic location and local weather conditions. All life stages of short-winged mole crickets may be present throughout the year.

### Perimeter Applications

Spread this product at a rate of 4 pounds per 1,000 square feet or 6.4 ounces per 100 square feet in a band at least 2 feet wide around foundations, including crevices where insects may hide.

**Resistance:** Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

## RECOMMENDED SPREADER SETTINGS

Note: Following will be a list of application spreaders and appropriate settings. This list will vary depending on market of spreaders. (This note will not appear on final printed label).

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

## Conditions of Sale and Limitation of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

In no event shall FMC or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Talstar and FMC—FMC Trademarks

(03/19/03)

