



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 21, 2025

Nina Marr
Registration Specialist
US Registration and Regulatory Affairs
FMC Corporation
c/o FMC Stine Research Center
P. O. Box 30
Newark, Delaware 19714-0030

Subject: Label Amendment - Registration Review Mitigation for Clomazone
Product Name: Cerano 5 MEG
EPA Registration Number: 279-3269
Case Number: 472340
Application Date: July 16, 2019

Dear Nina Marr:


The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Clomazone Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Concepción Rodríguez by phone at 202-566-0820, or via email at rodriguez.concepcion@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Julie Javier", with a stylized flourish at the end.

Julie Javier, Team Leader
Risk Mitigation and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

Cerano® 5 MEG**FOR AGRICULTURAL OR COMMERCIAL USE ONLY****5% Micro Encapsulated, Uniform Granule****For pre-emergence and post-emergence weed control in rice****EPA Reg. No. 279-3269****ACTIVE INGREDIENT**

Clomazone: 2-(2-chlorophenyl) methyl-4,

4-dimethyl-3-isoxazolidinone 5.0%

INERT INGREDIENTS 95.0%**TOTAL** 100.0%**EPA Est.** _____

% By Wt.

ACCEPTED**Oct 21, 2025**Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered underEPA Reg. No. **279-3269****KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID**If in Eyes**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes. Then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on Skin or Clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If Swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOTLINE NUMBERHave the product container or label with you when calling a poison control center, doctor or going for treatment.
You may also contact FMC Corporation at 1-800-331-3148 for emergency medical treatment information.

See other panels for additional precautionary information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust or vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

For medical emergencies involving this product, call toll free 1-800-331-3148.

Sold By:



[FMC Corporation]

2929 Walnut Street

Philadelphia, PA 19104

Nonrefillable Container**Net:** _____**OR****Refillable Container****Net:** _____

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical resistant gloves made of Barrier Laminate or Viton®;
- Shoes plus socks.

Wash thoroughly with soap and water after handling. Discard clothing and other absorbent materials that has been contaminated with this product. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides [40 CFR 170.305].

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

ENVIRONMENTAL HAZARDS

Except as directed on this label, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment wash waters. Apply this product only as specified on this label.

NON-TARGET ORGANISM ADVISORY STATEMENT:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing product drift.

USER SAFETY RECOMMENDATIONS

User should:

- Wash thoroughly with soap and water after handling.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Cerano® 5 MEG herbicide, referred to below as Cerano 5 MEG herbicide, must be used only in accordance with instructions on this label or as otherwise permitted by FIFRA. Always read the entire label, including the Conditions of Sale and Libation of Warranty and Liability.

To the extent consistent with applicable law, FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specified by FMC.

Crop	Pests	Rate of Application
Rice	Barnyard Grass Watergrass Spp. Sprangletop	12 lbs. Cerano® 5 MEG/acre (0.6 lb ai/A)

Cerano 5 MEG herbicide may be used at reduced rates (**do not use** less than 8 pounds per acre, (0.4 lb ai/A) for the control of sprangletop or for use on lighter textured soils, decomposed granitic soils or soils subject to excess drainage. When used at rates less than 12 pounds per acre (0.6 lb ai/A) additional use of labeled post-emergence herbicide applications may be required for control of watergrass species.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It

contains requirements for training, decontamination notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: A) Coveralls; B) Chemical-resistant gloves made of Barrier Laminate or Viton®; C) Shoes plus socks.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transpiration and Spills) at 1-800-424-9300.

GENERAL INFORMATION

Cerano 5 MEG herbicide is a pre-emergence and post-emergence herbicide for the control of watergrass, barnyardgrass (*Echinochloa* spp.) and sprangletop (*Leptochloa fascicularis*). Cerano 5 MEG herbicide controls weeds through foliar and root uptake as a chlorophyll-production inhibitor. Cerano 5 MEG is formulated as a 5% Micro encapsulated, uniform granule. **DO NOT** apply under conditions that would allow Cerano 5 MEG herbicide to come into contact with adjacent crops other than rice by drift as injury may occur.

RESTRICTIONS

- Apply this product to rice only as specified on this label. Do not apply to any other crop or site.
- Direct application to ditches used to transport irrigation water is prohibited.
- Do not apply more than 12 pounds per acre (0.6 lb ai/A) of Cerano 5 MEG herbicide per year on rice.
- Do not apply more than 12 pounds per acre (0.6 lb ai/A) of Cerano 5 MEG herbicide in a single application.
- Do not apply more than one application of Cerano 5 MEG per year on rice.
- **Pre-harvest intervals:** Do not apply later than 120 days before harvesting rice.
- Do not drain fields for a minimum of 14 days after application, unless the water is contained within a tailwater recovery system, or other system appropriate for preventing discharge from rice fields.
- Do not apply where runoff or irrigation water may flow directly onto agricultural land other than rice fields.
- Do not use this product for use in manufacturing or reformulation.
- Do not make applications of Cerano 5 MEG herbicide within 50 feet of any sensitive crops.
- Chemigation: Do not apply this product through any type of irrigation system.
- Do not fish or commercially grow fish, shellfish or crustaceans on treated acres during the year of treatment.
- Do not overlap or double apply ends of fields.
- Maintain a static water flood with no exposed soil for a minimum of 5 days after a post-flood application of Cerano 5 MEG herbicide.
- Do not apply sequential applications of other herbicides if rice is bleached or under stress from the use of Cerano 5 MEG herbicide.
- Do not apply Cerano 5 MEG herbicide to stressed rice.
- To facilitate green-up of bleached rice fields, fields need to receive a rapid flush of fresh water. Do not allow exposed soil in bleached fields.

PRECAUTIONS

- Applications of Cerano 5 MEG herbicide to fields that have been terraced or precision leveled with deep cuts may result in rice crop injury including stand loss. This precaution does not apply to normal maintenance leveling.
- Consult a UC Agricultural Extension Specialist or County Farm Advisors before applying Cerano 5 MEG herbicide to newly developed rice varieties.
- If temperature is in excess of 98° F for prolonged periods of time, (3 to 4 days), bleaching of rice and stand loss may occur.
- Lowering of water to allow exposed soil during the first 14 days following the Cerano 5 MEG herbicide application can cause bleaching of rice, possibly resulting in rice stand loss.
- For proper water circulation, functional water control devices at each end of each levee within a field are suggested.

WEED RESISTANCE MANAGEMENT

Cerano 5 MEG herbicide, which contains the active ingredient clomazone, is a group 13 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches) with a burndown herbicide.
- To minimize resistance, apply full labeled rates of Cerano 5 MEG herbicide.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.

- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 13 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad-spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 13 herbicides.
- Avoid making more than two applications of any Group 13 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields during and after harvest to reduce weed seed production.

DIRECTIONS FOR POST-FLOODED, PREPLANT USE ON RICE (WATER-SEEDED RICE AT PRE-SEEDING):

For control of watergrass and barnyardgrass, (*Echinochloa* spp.) and sprangletop (*Leptochloa fascicularis*), apply 12 pounds of Cerano 5 MEG herbicide per acre (0.6 lb ai/A) to a well-prepared seedbed, free of large clods. See rate exception under DIRECTIONS FOR USE.

Make application immediately after flooding and follow with the seeding of rice within 3 days after the Cerano 5 MEG herbicide application. Maintain water level in the checks with no exposed soil until the watergrass, barnyardgrass or sprangletop die, or for a minimum of 14 days after application.

Allowing water to recede or a prolonged soil exposure will result in loss of weed control and bleaching of rice. Excessively deep water will result in plant injury. Rice injury and/or stand thinning may be evident, especially when germinating rice is subject to stress conditions. Supplemental herbicide applications may be needed for season-long weed control.

DIRECTIONS FOR POST-FLOODED, POST-PLANT USE ON RICE (WATER-SEEDED RICE):

For control of watergrass and barnyardgrass (*Echinochloa* spp.) 1-leaf or less and sprangletop (*Leptochloa fascicularis*) 1-leaf or less apply 12 pounds of Cerano 5 MEG herbicide per acre (0.6 lb ai/A) to a well-prepared seedbed free of large clods. See rate exception under DIRECTIONS FOR USE.

Maintain a shallow flood in the checks after application with no exposed soil, or until watergrass, barnyard grass and sprangletop die, or for a minimum of 14 days.

Allowing water to recede or prolonged soil exposure will result in loss of weed control and bleaching of rice. Excessively deep floodwater will result in plant injury. Rice injury and/or stand thinning may be evident, especially when germinating rice is subjected to stress conditions. Supplemental herbicide applications may be needed for season-long weed control.

ROTATIONAL CROP RESTRICTIONS

After application of Cerano 5 MEG herbicide, rotational crops may not be planted prior to the intervals specified below:

Note: When using Cerano 5 MEG herbicide with other registered herbicides always refer to rotational restrictions and precautions on the other product label.

NO RESTRICTION	9 Months
Cotton*	Cotton
Peas	Corn (field, pop, seed, sweet)
Peppers	Cucurbits
Pumpkins (processing)	Dry Beans
Rice	Peanuts
Soybeans	Potatoes
Squash	Snap Beans
Sweet Potatoes	Sorghum
Tobacco	Sugar Beets
Tuberous & Corm Vegetables	Tomatoes (Transplanted)
(Arracacha, Cassava, Tanier, Yam Beans, True Yams)	
12 Months: All crops not listed	
Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Cerano 5 MEG herbicide treatment.	
*REQUIREMENTS FOR PLANTING TIME APPLICATIONS: If cotton is the rotational crop and is planted within 9 months after the application of Cerano 5 MEG herbicide, an organophosphate insecticide either disulfoton or phorate must be applied in-furrow with the seed at planting at a minimum of 0.75 pound per acre of active ingredient. Failure to apply either disulfoton or phorate in accordance with in-furrow label use direction can result in crop phytotoxicity (bleaching) and/or stand reduction. Do not use diuron at this planting time application as plant injury may result. Refer to the insecticide product labels for appropriate in-furrow application directions and maximum use rates. Monitor application equipment to ensure accurate and uniform placement of the insecticide.	

REPLANTING INSTRUCTIONS

If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with Cerano 5 MEG herbicide. **Do not** retreat fields with a second application of Cerano 5 MEG herbicide. **Do not** replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP RESTRICTIONS on this label.

Aerial Application Methods and Equipment: Provide accurate and uniform distribution of granule over the treated area.

APPLICATION INSTRUCTIONS

AERIAL APPLICATION: Fixed wing aircraft and helicopters can be used to apply Cerano 5 MEG, however, when making applications by fixed wing aircraft maintain appropriate buffer zones to prevent product drift out of the target area.

Dealers Must Sell in Original Packages Only.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent permitted by applicable law, buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

STORAGE AND DISPOSAL

Pesticide Storage: Store the product in original container only. Do not contaminate water, other pesticides, fertilizer, food, or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums with Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums with Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with CERANO 5MEG herbicide containing clomazone only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with Cerano 5MEG herbicide containing clomazone only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact FMC at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transportation and Spills) at 1-800-424-9300, day or night.

WILBUR-ELLIS® logo, Ideas to Grow With® and CERANO® are registered trademarks of Wilbur-Ellis Company LLC.
HERBICIDE™ logo is a trademark of Wilbur-Ellis Company LLC.

FMC is a trademark of FMC Corporation

Label code: D-4250 102125