

279-3320

04/17/2007

118



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: 279-3320

Date of Issuance: APR 17 2007

Term of Issuance: Expires April 17, 2009

Name of Pesticide Product: Echelon 0.57% Granular Herbicide on Fertilizer

NOTICE OF PESTICIDE:

[x] Registration
[] Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Submit within six months of the date of this Registration Notice, the results of a five batch analysis (830.1700), conducted on Echelon MUP.
3. Submit storage stability and corrosion characteristics studies (830.6317 and 830.6320, respectively) within one year of the date of this registration notice.

Make the following label changes:

- 4. Revise the EPA Registration Number to read: "EPA Reg. No. 279-.3320."
5. Add the following language to the "Precautionary Statements": "Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature of Joanne I. Miller

Date:

APR 17 2007

Page Number 2
EPA Reg. No. 279-3320

6. Under the heading "New Product Facts", remove this statement, "It is a violation of Federal law to use this product in a manner inconsistent with its labeling" and locate it **above** the heading.
7. Under the "First Aid" section, in the "If Swallowed" statement, delete this sentence: "Do not give liquids to the person.", and replace with "Have person sip a glass of water if able to swallow."
8. Under the "Precautionary Statement" make the following changes:
 - a. Revise the third sentence to read: "Avoid breathing dust."
 - b. Revise the fourth sentence to read: "Do not get on skin, in eyes or on clothing."
 - c. Delete the statement: "Do not contaminate water, food or feed by storage."
 - c. Delete the statement that begins with: "Do not graze..." and place under "New Product Facts".
9. Revise the "Pesticide Storage" statement to read: "Store product in original container only, away from other pesticides, fertilizer, food or feed. Reseal open bag by folding top down and securing. Store in a cool, dry place and avoid excess heat."

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

ECHELON®

0.57% Granular Herbicide on Fertilizer for consumers
For Selective Preemergence and Postemergence Control of Sedges,
Annual Grasses and Broadleaf weeds in Residential Lawns.

EPA Reg. No. 279-~~3320~~ EPA Est. 279-

Active Ingredient: By Wt.

Sulfentrazone: N-[2,4-dichloro-5-[4-(difluoro methyl-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl)]phenyl]methanesulfonamide	0.19%
Prodiamine: [N3,N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine	0.38%
Other Ingredients	99.43%
	100.0%

Contains 0.57% pounds of active ingredient per pound of product
U. S. Patent No. 4,818,275, Patents pending

**ACCEPTED
with COMMENTS
in EPA Letter Dated:**

APR 17 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

279-3320

Guaranteed Analysis XX-X-XX

Total Nitrogen (N)	XX.XX %
XX.XX% Urea Nitrogen	
Soluble Potash (K₂O)	XX.XX %
Sulfur (S) Total	XX.XX %
X.XX % Free Sulfur (S)	
X.XX % Combined Sulfur (S)	
Iron (Fe) Total	X.XX %
X.XX % Soluble Iron (Fe)	
Derived From: Poly Coated Sulfur Coated Urea, Sulfate Of Potash, Iron Sucrate.	
Chlorine (Cl) Max.	X.XX %

- The most effective crabgrass, foxtail, spurge and oxalis preventer you can buy.
- Halts crabgrass and other weeds before they appear
- Halts weeds already present
- Kills broad leaf weeds already in your lawn and prevents the growth on new weeds including broadleaf and grassy weeds and nutsedge

FMC Corporation
Agricultural Products Group
Philadelphia PA 19103

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back for additional precautionary information

New Product Facts

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

- Echelon is a selective pre-emergence and post-emergence turf herbicide for control of sedges, annual grass and broadleaf weeds in residential lawns.
- Do not use on Dichondra or St. Augustine lawns
- In early spring apply when lawn is dry to kill weeds already in the lawn including Plantain, Oxalis, Dandelion, Violet, Wild onion, Garlic and Black medic and prevent germination of Crabgrass, Foxtail, spurge, Barnyard grass, fall panicum, goosegrass, nutsedge, oxalis, Chickweed, Purslane and Pigweed:
 - Before March 1 in southern California. For best results apply by mid February
 - Before mid March in northern California
 - Early Spring (up to the time dandelions are in puffball stage) in other states
 - For Goosegrass control repeat application in 8 weeks
- Rain, snow or freezing conditions will not effect the performance of the product.
- In the fall (August-October) apply after temperatures have cooled and the fall rains have begun to prevent the germination of *Poa annua*, Chickweed, in the fall and crabgrass and oxalis in the spring
- Don't use on newly seeded grassy lawns until after the second mowing. On newly sprigged areas wait 4 months before application
- Do not plant grass seed for four months after using this product.
- Do not apply more frequently than twice a year. Allow at least one month interval between applications
- Do not rake or aerify after application.
- Do not apply this product to flower beds

Spreading Tips- Apply header strips. Then apply product in longest direction of lawn..

Diagram of Spreading TIPS

Spreader Settings

For best results, use a Scotts spreader and mow the lawn before applying this product

Scotts Rotary Spreaders	
Scotts Speedy Green	4 Make each pass about 5 feet from previous pass to overlap spread width
Scotts Easy Green	26.5 Make each pass about 30 inches from previous pass to overlap spread width
Scotts Drop Spreaders	
Scotts AccuGreen	6 Make sure spreader wheel completely overlaps previous wheel mark
Scotts PrecisionGreen	7 Make sure spreader wheel completely overlaps previous wheel mark

Additional Spreader Settings

- EZ Dorp Republic (#530) 6.5
- EZ Broadcast Republic (#540)....4
- Scotts Broadcast Tractor Spreader....4

(Non-Scotts spreader settings are approximate and may need to be adjusted for proper coverage)

FIRST AID	
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
For Information Regarding the Use of this Product Call 1-800-321-1FMC(1362)	

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Do not contaminate water, food or feed by storage. Do not graze or feed livestock forage cut from areas treated with Echelon turf herbicide.

Environmental Hazards

This pesticide is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface water advisory: Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store product in original container only, in a cool, dry place and avoid excess heat. Reseal opened bag by folding top down and securing

Pesticide Disposal

Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal

Do not reuse bag. Discard bag in Trash

Table2.

Weed controlled from Pre- emergence application of Echelon

Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>
Bluegrass annual /	<i>(Poa annua)</i>
Carpetweed	<i>(Mollugo verticillata)</i>
Carolina Geranium	<i>Geranium carolinianum</i>

Chickweed, common 2	(<i>Stellaria media</i>)
Chickweed, mouseear (from seed)	(<i>Cerastium vulgatum</i>)
Crabgrass (large, Smooth) 3	(<i>Digitaria spp.</i>)
Crowfootgrass	(<i>Dactyloctenium aegyptium.</i>)
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)
Foxtails, Annual	(<i>Setaria spp..</i>)
Goosegrass 4	(<i>Eleusine indica</i>)
Hairy Bittercress	<i>Cardamine hirsuta</i>
Henbit 2	(<i>Lamium amplexicaule</i>)
Itchgrass	(<i>Rottboellia exaltata</i>)
Knotweed, prostrate	(<i>Polygonum aviculare</i>)
Kochia	(<i>Kochia scoparia</i>)
Kyllinga, green	(<i>Kyllinga brevifolia</i>)
Kyllinga, false	(<i>Kyllinga gracillima</i>)
Lambsquarters, common	(<i>Chenopodium album</i>)
Lovegrass	(<i>Eragrostis spp.</i>)
Nutsedge, yellow	(<i>Cyperus esculentus</i>)
Panicum (Texas, Fall)	(<i>Panicum spp.</i>)
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)
Purslane, Common	(<i>Portulaca oleracea</i>)
Pusley, Florida	(<i>Richardia scabra L.</i>)
Sedge, globe	(<i>Cyperus globulosus</i>)
Sedge, cylindrical	(<i>Cyperus retrorsus</i>)
Sedge, Surinam	(<i>Cyperus surinamensis</i>)
Sedge, Texas	(<i>Cyperus polystachyos</i>)
Shephardspurse 2	(<i>Capsella bursa-pastoris</i>)
Signalgrass, broadleaf	(<i>Brachiaria playphylla</i>)
Speedwell	(<i>Veronica spp.</i>)
Sprangletop	(<i>Leptochola spp.</i>)
Spurge, prostrate	(<i>Euphorbia humistrata</i>)
Witchgrass	(<i>Panicum capillare L.</i>)
Woodsorrel, yellow (from Seed)	(<i>Oxalis stricta</i>)

1. Poa annua is a winter annual applications of Echelon should be made in August or September to established, non overseeded turf before Poa Annu germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August - September are approximate timings.
2. Apply Echelon in late summer, fall or winter before weed germinates for control.
3. Crabgrass control in cool season grasses from fall application:
Areas where the ground is subject to winter freezes, Echelon may be applied in the fall at rates listed in Table 1 for cool season grasses when soil temperatures are below 50F, but prior to the ground freezing. This will provide control of crabgrass the following spring.
4. In areas of extended growing season and heavy pressure weed control will be enhanced by second application of Echelon 60 -90 days after initial application

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds

EHELON herbicide will control or suppress the weeds listed in Table 3 when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1.

Table 3. Weeds Controlled or Suppressed by Echelon herbicide when application made post emergence to weeds.

BROADLEAVES	SCIENTIFIC NAMES
Bittercress	(<i>Cardamine spp.</i>)
Black medic	(<i>Medicago lupulina</i>)
Buttercups	(<i>Ranunculus spp.</i>)
Carolina geranium	(<i>Geranium carolinianum</i>)
Carpetweed	(<i>Mollugo verticillata</i>)
Chickweed, common	(<i>Stellaria media</i>)

Chickweed, mouseear	(<i>Cerastium vulgatum</i>)
Cinquefoil	(<i>Potentilla spp.</i>)
Clover	(<i>Trifolium spp.</i>)
Cudweed	(<i>Gnaphalium spp.</i>)
Dandelion	(<i>Taraxacum officinale</i>)
Dock, Curly	(<i>Rumex crispus</i>)
Evening primrose	(<i>Oenothera biennis</i>)
Fiddleneck	(<i>Amsinckia spp.</i>)
Filaree	(<i>Erodium spp.</i>)
Goldenrod	(<i>Solidago spp.</i>)
Ground ivy	(<i>Glechema hederacea</i>)
Henbit	(<i>Lamium amplexicaule</i>)
Knotweed, prostrate	(<i>Polygonum aviculare</i>)
Kochia	(<i>Kochia scoparia</i>)
Lambsquarters, common	(<i>Chenopodium album</i>)
Lawn burweed	(<i>Soliva pterosperma</i>)
Lespedeza, common	(<i>Lespedeza striata</i>)
Mallow, common	(<i>Malva neglecta</i>)
Parsley pierf	(<i>Alchemilla arvensis</i>)
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)
Pigweed, Tumble	(<i>Amaranthus albus</i>)
Pineapple weed	(<i>Matricaria matricarioides</i>)
Plantain, buckhorn	(<i>Plantago lanceolata</i>)
Puncture weed	(<i>Tribulus terrestris</i>)
Purslane, common	(<i>Portulaca oleracea</i>)
Pusley, Florida	(<i>Richardia scabra</i>)
Redweed	(<i>Melochia corchorifolia</i>)
Rocket, London	(<i>Sisymbrium irio</i>)
Smartweed, Pennsylvania	(<i>Polygonum pensylvanicum</i>)
Sorrel, Red	(<i>Rumex acetosella</i>)
Speedwell	(<i>Veronica spp.</i>)
Spurge, (annuals)	(<i>Euphorbia spp.</i>)
Spurge, prostrate	(<i>Euphorbia humistrata</i>)
Spurge, spotted	(<i>Euphorbia maculata</i>)
Star of Bethlehem	(<i>Ornithogalum umbellatum</i>)
Velvetleaf	(<i>Abutilon theophrasti</i>)
Violet, wild	(<i>Viola pratensis</i>)
Wild garlic	(<i>Allium vineale</i>)
Wild onion	(<i>Allium canadense</i>)
Woodsorrel, creeping	(<i>Oxalis corniculata</i>)
Woodsorrel, yellow	(<i>Oxalis stricta</i>)

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent permitted by law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC – Trademark of FMC Corporation