


279-3322

04/17/2007

1/13

 <p style="text-align: center;"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b> Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Reg. Number: 279-3322	Date of Issuance: APR 17 2007
	Term of Issuance: Expires April 17, 2009	
	Name of Pesticide Product: Echelon TM 0.3% Granular Herbicide on Fertilizer	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)		
Name and Address of Registrant (include ZIP Code): FMC Corporation 1735 Market Street Philadelphia, PA 19103		
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:</p> <ol style="list-style-type: none"> <li>1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>2. Submit within six months of the date of this Registration Notice, the results of a five batch analysis (830.1700) conducted on Echelon MUP.</li> <li>3. Submit storage stability and corrosion characteristics studies (830.6317 and 830.6320, respectively) within one year of the date of this registration notice.</li> </ol> <p>Make the following label changes:</p> <ol style="list-style-type: none"> <li>4. Revise the EPA Registration Number to read: "EPA Reg. No. 279-3322."</li> </ol>		
Signature of Approving Official:  Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	Date:  APR 17 2007	

5. Add the following language to the "Precautionary Statements": "Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals."

6. Under the "First Aid" section, in the "If Swallowed" statement, delete this sentence: "Do not give any liquid to the person.", and replace with "Have person sip a glass of water if able to swallow."

7. Under the "Precautionary Statement" make the following changes:

- a. Revise the third sentence to read: "Avoid breathing dust."
- b. Revise the fourth sentence to read: "Do not get on skin, in eyes or on clothing."
- c. Add this statement: "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco."

8. Revise the "Pesticide Storage" statement to read: "Store product in original container only, away from other pesticides, fertilizer, food or feed. Reseal open bag by folding top down and securing. Store in a cool, dry place and avoid excess heat."

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure

# ECHELON™

## 0.3% Granular Herbicide on Fertilizer

For Preemergence and early Postemergence Control of Crabgrass. Selective Preemergence and Postemergence Control of Annual Grasses, Broadleaf weeds and Sedges in Turf Sites Including Residential and Institutional Lawns, Athletic Fields, Commercial Sod Farms, Golf Course Fairways and Roughs, Roadsides, Utility right-of-ways, Railways and Industrial areas

**Active Ingredient:**

Sulfentrazone: N-[2,4-dichloro-5-[4-(difluoromethyl-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl)phenyl]methanesulfonamide .....

Prodiamine: [N3,N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine

Other Ingredients: .....

Echelon 0.3% Granular contains 0.3 pounds of active ingredient per pound of product  
U. S. Patent No. 4,818,275

**Guaranteed Analysis XX-X-XX**

Total Nitrogen (N) ..... XX.XX%

XX.XX% Urea Nitrogen

Soluble Potash (K<sub>2</sub>O) ..... XX.XX%

Sulfur (S) Total ..... XX.XX%

X.XX% Free Sulfur (S)

X.XX% Combined Sulfur (S)

Iron (Fe) Total ..... XX.XX%

X.XX% Soluble Iron (Fe)

Derived From: Poly Coated Sulfur Coated Urea,  
Sulfate Of Potash, Iron Sulfate

Chlorine (Cl) Max ..... X.XX%

By Wt.

0.1% **ACCEPTED**  
0.2% **with COMMENTS**  
**In EPA Letter Dated:**  
99.7%  
100.0%

APR 17 2007  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

279-3322

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

See additional precautionary information contained inside label booklet.

FMC Corporation  
Agricultural Products Group  
Philadelphia PA 19103

EPA Reg. No. 279-XXXX  
EPA Est. 279-

Net Contents

<b>FIRST AID</b>	
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
For Information Regarding the Use of this Product Call 1-800-321-1FMC(1362)	

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans (and Domestic Animals)

#### CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

- Users should remove clothing immediately if product gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface water advisory: Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

---

### DIRECTIONS FOR USE

---

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

**Non-Agricultural Use Requirements**  
 The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

**STORAGE AND DISPOSAL**  
 Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store product in original container only, in a cool, dry place and avoid excess heat.

**Spill:** In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. **Confine spills. Call FMC: (800) 331-3148.**

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal:** If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

---

**GENERAL INFORMATION**

---

**Turf Use Instructions**  
 Echelon is a selective pre-emergence and post-emergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in turf sites including residential and institutional lawns, athletic fields, commercial sod farms, golf course fairways and roughs, roadsides, utility right-of-ways, railways and industrial areas.

Echelon is a granular formulation containing 0.3 lbs. of active ingredient per pound of formulated product. The mode of action of Echelon herbicide involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

---

**SPECIFIC USE INSTRUCTIONS IN TURFGRASS**

---

**Established Turf**  
 This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury

may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

Broadcast Echelon with suitable application equipment to ensure uniform coverage over the treatment area. A drop or rotary spreader designed to apply granular herbicide or insecticide is the preferred application of this product. Calibrate application equipment prior to application. Consult manufacturer operation manual for proper calibration and operation of application equipment. Apply 2.87 to 8.60 pounds per 1,000 square feet to control broadleaf weeds and grasses. See Table 1 for application rates based on turf grasses

**When applied as directed under the conditions described, the following established grasses are tolerant to Echelon herbicide at the recommended use rates in a range from 0.250 to 1.125 lb. ai/acre (83 to 375 pounds/acre or 1.91 to 8.6 pounds/1,000 sq. ft.). Do not exceed 1.125 lb. ai/acre (375/acre) per calendar year.**

Table 1. Tolerant grasses.

<i>Grass Type</i>	<i>Recommended Use Rates</i> <i>Do not exceed maximum use rates per turf species in a single application</i>		
	<b>Lb. ai. per acre</b>	<b>Pounds per 1000 ft<sup>2</sup></b>	<b>Pounds per acre</b>
<b>Cool Season Grasses</b>			
Bentgrass, creeping <sup>1</sup> Fescue, fine <sup>2</sup> ( <i>Festuca rubra</i> ) Ryegrass, perennial ( <i>Lolium perenne</i> )	0.25-0.375 <sup>3</sup> lb. ai.	1.91 – 2.87 lbs	83 – 125 lbs
Bluegrass, Kentucky ( <i>Poa pratensis</i> ) Fescue, tall <sup>2</sup> ( <i>Festuca arundinacea</i> )	0.57-0.75 <sup>3</sup> lb. ai.	3.81 – 5.74 lbs	166 – 250 lbs
<b>Warm Season Grasses</b>			
Bahiagrass ( <i>Paspalum notatum</i> ) Buffalograss ( <i>Buchloe dactyloides</i> ) Carpentergrass ( <i>Axonopus affinis</i> ) Centipedegrass ( <i>Eremochloa ophiuroides</i> ) Kikuyugrass ( <i>Pennisetum clandestinum</i> ) Seashore Paspalum ( <i>Paspalum vaginatum</i> ) Zoysiagrass ( <i>Zoysia japonica</i> )	0.57-0.75 <sup>3</sup> lb. ai.	3.81 – 5.74 lbs	166 – 250 lbs
Bermudagrass ( <i>Cynodon dactylon</i> ) & hybrids	0.75-1.125 <sup>3</sup> lb. ai.	5.74 – 8.6 lbs	250 – 375 lbs

1 If bentgrass is mowed lower than 0.5 inches do not apply Echelon.

2 Use of this product on certain varieties of Chewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury.

3 Echelon may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate per turf type or 1.125 lbs. ai/acre (375 pounds/acre), whichever is lower.

Not all varieties of grasses have been tested. Consult your local FMC Technical Representative for additional recommendations on application rates for your local area or turf types.

## Weed Controlled With Preemergence Application

Table 2.

Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>
Bluegrass annual <sup>1</sup>	<i>(Poa annua)</i>
Carpeweed	<i>(Mollugo verticillata)</i>
Carolina Geranium	<i>Geranium carolinianum</i>
Chickweed, common <sup>2</sup>	<i>(Stellaria media)</i>
Chickweed, mouseear (from seed)	<i>(Cerastium vulgatum)</i>
Crabgrass (large, Smooth) <sup>3</sup>	<i>(Digitaria spp.)</i>
Crowfootgrass	<i>(Dactyloctenium aegyptium.)</i>
Cupgrass, Woolly	<i>(Eriochloa villosa)</i>
Foxtails, Annual	<i>(Setaria spp.)</i>
Goosegrass <sup>4</sup>	<i>(Eleusine indica)</i>
Hairy Bittercress	<i>Cardamine hirsute</i>
Henbit <sup>2</sup>	<i>(Lamium amplexicaule)</i>
Itchgrass	<i>(Rottboellia exaltata)</i>
Knotweed, prostrate	<i>(Polygonum aviculare)</i>
Kochia	<i>(Kochia scoparia)</i>
Kyllinga, green	<i>(Kyllinga brevifolia)</i>
Kyllinga, false	<i>(Kyllinga gracillima)</i>
Lambsquarters, common	<i>(Chenopodium album)</i>
Lovegrass	<i>(Eragrostis spp.)</i>
Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Panicum (Texas, Fall)	<i>(Panicum spp.)</i>
Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>
Purslane, Common	<i>(Portulaca oleracea)</i>
Pusley, Florida	<i>(Richardia scabra L.)</i>
Sedge, globe	<i>(Cyperus globulosus)</i>
Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Sedge, Texas	<i>(Cyperus polytachyos)</i>
Shephardspurse <sup>2</sup>	<i>(Capsella bursa-pastoris)</i>
Signalgrass, broadleaf	<i>(Brachiaria platyphylla)</i>
Speedwell	<i>(Veronica spp.)</i>
Sprangletop	<i>(Leptochola spp.)</i>
Spurge, prostrate	<i>(Euphorbia humistrata)</i>
Witchgrass	<i>(Panicum capillare L.)</i>
Woodsorrel, yellow (from Seed)	<i>(Oxalis stricta)</i>

1. *Poa annua* is a winter annual. Applications of Echelon should be made in August or September to established, non overseeded turf before *Poa Annua* germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings.
2. Apply Echelon in late summer, fall or winter before weed germinates for control.
3. Crabgrass control in cool season grasses from fall application:  
Areas where the ground is subject to winter freezes, Echelon may be applied in the fall at rates listed in Table 1 (83-250 pounds per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.
4. In areas of extended growing season and heavy pressure weed control will be enhanced by second application of Echelon 60-90 days after initial application



### Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds

Echelon herbicide will control or suppress the weeds listed in Table 3 when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1.

### Weeds Controlled or Suppressed When Application Made Postemergence to Weeds

Table 3

Common Name	Scientific Name
Bittercress	( <i>Cardamine</i> spp.)
Black medic	( <i>Medicago lupulina</i> )
Buttercups	( <i>Ranunculus</i> spp.)
Carolina geranium	( <i>Geranium carolinianum</i> )
Carpetweed	( <i>Mollugo verticillata</i> )
Chickweed, common	( <i>Stellaria media</i> )
Chickweed, mouseear	( <i>Cerastium vulgatum</i> )
Cinquefoil	( <i>Potentilla</i> spp.)
Clover	( <i>Trifolium</i> spp.)
Cudweed	( <i>Gnaphalium</i> spp.)
Dandelion	( <i>Taraxacum officinale</i> )
Dock, Curly	( <i>Rumex crispus</i> )
Evening primrose	( <i>Oenothera biennis</i> )
Fiddleneck	( <i>Amsinckia</i> spp.)
Filaree	( <i>Erodium</i> spp.)
Goldenrod	( <i>Solidago</i> spp.)
Ground ivy	( <i>Glechoma bederacea</i> )
Henbit	( <i>Lamium amplexicaule</i> )
Knotweed, prostrate	( <i>Polygonum aviculare</i> )
Kochia	( <i>Kochia scoparia</i> )
Lambsquarters, common	( <i>Chenopodium album</i> )
Lawn burweed	( <i>Soliva pterisperma</i> )
Lespedeza, common	( <i>Lespedeza striata</i> )
Mallow, common	( <i>Malva neglecta</i> )
Parsley piert	( <i>Alchemilla arvensis</i> )
Pigweed, Redroot	( <i>Amaranthus retroflexus</i> )
Pigweed, Tumble	( <i>Amaranthus albus</i> )
Pineapple weed	( <i>Matricaria matricarioides</i> )
Plantain, buckhorn	( <i>Plantago lanceolata</i> )
Puncture weed	( <i>Tribulus terrestris</i> )
Purslane, common	( <i>Portulaca oleracea</i> )
Pusley, Florida	( <i>Richardia scabra</i> )
Redweed	( <i>Melochia corchorifolia</i> )
Rocket, London	( <i>Sisymbrium irio</i> )
Smartweed, Pennsylvania	( <i>Polygonum pennsylvanicum</i> )
Sorrel, Red	( <i>Rumex acetosella</i> )
Speedwell	( <i>Veronica</i> spp.)
Spurge, (annuals)	( <i>Euphorbia</i> spp.)
Spurge, prostrate	( <i>Euphorbia humistrata</i> )
Spurge, spotted	( <i>Euphorbia maculata</i> )
Star of Bethlehem	( <i>Ornithogalum umbellatum</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )
Violet, wild	( <i>Viola pratensis</i> )
Wild garlic	( <i>Allium vineale</i> )
Wild onion	( <i>Allium canadense</i> )
Woodsorrel, creeping	( <i>Oxalis corniculata</i> )
Woodsorrel, yellow	( <i>Oxalis stricta</i> )

### Application to Reseeded, Overseeded or Sprigged areas

Reseeding, overseeding or sprigging of treated areas within three to twelve (3-12) months after application of this product could inhibit the establishment of desirable grasses. [See table for exact intervals from application].

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

### Overseeding Intervals after Echelon Applications

Table 4

Total Amount of Echelon 4SC Applied Pounds Product/Acre	Interval (Months Before Overseeding)		
	North	Transition	South
83 - 125	3	3	3
166 - 250	4	4	4
375	6	5	5

### Sod Production

It is recommended that sod be established for at least six (6) months before an application of Echelon herbicide.

Do not apply this product within three (3) months of harvest.

### Other Use Precautions

Do not apply to golf course putting greens or tees.

Do not use on turgrasses other than those listed on this label.

Do not graze or feed livestock forage cut from areas treated with Echelon herbicide.

Do not apply directly to landscape ornamentals or ornamental beds.

Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are *desirable* species.

Do not apply to creeping bentgrass if mowing height is less than 0.5 inch.

Do not harvest treated sod before 90 days after application.

Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.

Do not apply with adjuvants or surfactants.

---

**ADDITIONAL USE INSTRUCTIONS**

---

**Railroad Rights-of-Way**

Controls many weeds and maintains bare ground on railroad rights-of-way, including railroad yards, railroad crossings and railroad bridge abutments.

**Highway, Roadside, Pipeline and Utility Rights-of-Way**

Controls many weeds and maintains bare ground in highway, roadside, pipeline and utility rights-of-way. Such areas would include, but are not limited to, guardrails, road shoulders, electric utility substations, pipeline pumping stations, around electric transmission towers, around distribution line poles plus other areas where complete vegetation control is desired.

**Industrial Areas, Fence Rows and Other Non-crop Sites**

Controls weeds and maintains bare ground in industrial areas including production facilities, tank farms, storage areas, parking areas, lumber yards, airports, military installations, along fence rows, and in similar non-crop sites where complete vegetation control is needed.

---

**METHOD AND RATE OF APPLICATION FOR ADDITIONAL USES**

---

For residual control of germinating weeds in non-crop land, apply as a broadcast treatment using 250 to 375 pounds (0.75 to 1.125 pounds active ingredient) per acre by ground in a suitable granular applicator.

DO NOT apply Echelon Herbicide to soils classified as sand with less than 1% Organic Matter.

**Timing**

For best results, apply Echelon alone or in combination with other herbicides for residual control of weeds in late summer, fall, or early spring to insure adequate moisture for soil activation.

**Weeds Controlled**

Echelon when applied postemergence at 250 to 375 pounds per acre, will control the following weeds in non-cropland areas (Table 5). Use the higher labeled rates to extend length of control. Use the higher rates on sites with fine soil textures and on sites with more than 2% organic matter.

## Weeds Controlled from Preemergence Application to Bare Ground Prior to Weed Seed germination

Table 5

Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>
Bluegrass annual	<i>(Poa annua)</i>
Carpetweed	<i>(Mollugo verticillata)</i>
Carolina Geranium	<i>Geranium carolinianum</i>
Chickweed, common	<i>(Stellaria media)</i>
Chickweed, mouseear (from seed)	<i>(Cerastium vulgatum)</i>
Crabgrass ( large, Smooth)	<i>(Digitaria spp.)</i>
Crowfootgrass	<i>(Dactyloctenium aegyptium.)</i>
Cupgrass, Woolly	<i>(Eriochloa villosa)</i>
Foxtails, Annual	<i>(Setaria spp..)</i>
Goosegrass	<i>(Eleusine indica)</i>
Hairy Bittercress	<i>Cardamine hirsuta</i>
Henbit 2	<i>(Lamium amplexicaule )</i>
Itchgrass	<i>(Rottboellia exaltata)</i>
Knotweed, prostrate	<i>(Polygonum aviculare)</i>
Kochia	<i>(Kochia scoparia)</i>
Kyllinga, green	<i>(Kyllinga brevifolia)</i>
Kyllinga, false green	
Lambsquarters, common	<i>(Chenopodium album)</i>
Lovegrass	<i>(Eragrostis spp.)</i>
Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Nutsedge, purple	<i>(Cyperus rotundus)</i>
Panicum (Texas, Fall)	<i>(Panicum spp.)</i>
Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>
Purslane, Common	<i>(Portulaca oleracea)</i>
Pusley, Florida	<i>(Richardia scabra L.)</i>
Sedge, globe	<i>(Cyperus globulosus)</i>
Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Sedge, Texas	<i>(Cyperus polytachyos)</i>
Shephardspurse	<i>(Capsella bursa-pastoris)</i>
Signalgrass, broadleaf	<i>(Brachiaria platyphylla)</i>
Speedwell	<i>(Veronica spp.)</i>
Sprangletop	<i>(Leptochloa spp.)</i>
Spurge, prostrate	<i>(Euphorbia humistrata)</i>
Witchgrass	<i>(Panicum capillare L.)</i>
Woodsorrel, yellow (from Seed)	<i>(Oxalis stricta)</i>

**CONDITIONS OF SALES AND LIMITATIONS OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent permitted by law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC – Trademark of FMC Corporation