



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 8, 2021

Catherine Rice
Product Registration Manager
FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Subject: PRIA Label and CSF Amendment – Revisions to support maximum nitrogen fertilizer content of 46%, add state-specific fertilizer language and other label updates; revise basic CSF and add new alternate 1 CSF
Product Name: Echelon 0.16% Granular Herbicide on Fertilizer
EPA Registration Number: 279-3325
Application Date: October 22, 2020
Decision Number: 567160

Dear Ms. Rice:

The amended label and CSFs referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, are acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Please note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/02/2021
- Alternate 1 CSF dated 08/02/2021

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those

claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, you may contact Jamie Harrington at harrington.jamie@epa.gov.

Sincerely,



Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

NOTE to EPA Product Manager: Text in brackets [] is optional.

| | | | |
|----------------------|--------------|-----------|------------------|
| PRODIAMINE | GROUP | 3 | HERBICIDE |
| SULFENTRAZONE | GROUP | 14 | HERBICIDE |

ECHELON®

0.16% Granular Herbicide on Fertilizer

For Preemergence and early postemergence Control of Crabgrass, selected annual grasses, sedges and annual broadleaf weeds in Turf Sites - Residential and Institutional Lawns, Athletic Fields, Commercial Sod Farms, Golf Course Fairways and Roughs, Roadsides, Utility right-of-ways, Railways and Industrial areas

EPA Reg. No. 279-3325


EPA Est. _____

| Active Ingredients: | By Wt. |
|--------------------------|---------|
| Sulfentrazone | 0.053% |
| Prodiamine | 0.107% |
| Other Ingredients: | 99.840% |
| Total: | 100.0% |

This product contains 0.0016 pounds of active ingredients (0.00053 lb ai sulfentrazone and 0.00107 lb ai prodiamine) per pound of product

Guaranteed Analysis XX-X-XX

Total Nitrogen (N) XX.XX% {Note to reviewer: Total Nitrogen value cannot exceed 46%}
 XX.XX% Urea Nitrogen
 Soluble Potash (K₂O) XX.XX%
 Sulfur (S) Total XX.XX%
 X.XX% Free Sulfur (S)
 X.XX% Combined Sulfur (S)
 Iron (Fe) Total XX.XX%
 X.XX% Soluble Iron (Fe)
 Derived From: Poly Coated Sulfur Coated Urea,
 Sulfate Of Potash, Iron Sucrate
 Chlorine (Cl) Max X.XX%



KEEP OUT OF REACH OF CHILDREN

CAUTION

See [other panels][booklet][back] for additional precautionary information.

FMC Corporation
2929 Walnut Street
Philadelphia PA 19104

Net Contents

| FIRST AID | |
|---|--|
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| MEDICAL HOTLINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information. | |

For Technical Support or information regarding the use of this product, call 1-800-321-1FMC(1362)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should remove clothing/PPE immediately if product gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants and aquatic organisms in adjacent areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface water advisory: Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product may only be used in accordance with the Application Rates listed on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

WEED RESISTANCE-MANAGEMENT

For resistance management, please note that Echelon 0.16% Granular Herbicide on Fertilizer contains both a Group 3/[Prodiamine] and a Group 14/[Sulfentrazone] herbicide. Any weed population may contain plants naturally resistant to Group 3 and/or Group 14 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 and/or 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control

the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout area before herbicide application for weed identification and growth stage.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your local extension specialist or university researchers for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.

PRODUCT INFORMATION

Echelon 0.16% Granular Herbicide on Fertilizer is a granular herbicide on fertilizer carrier formulation containing 0.16 pounds of active ingredients per pound of formulated product. The mode of action of this product involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

This product is a selective pre-emergence and post-emergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in turf sites including residential and institutional lawns, athletic fields, commercial sod farms, golf course fairways and roughs, roadsides, utility right-of-ways, railways and industrial areas.

SPECIFIC USE INSTRUCTIONS FOR USE ON TURFGRASS

Established Turf

This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

Broadcast this product with suitable application equipment to ensure uniform coverage over the treatment area. A drop or rotary spreader designed to apply granular herbicide or insecticide is the preferred application of this product. Calibrate application equipment prior to application. Consult manufacturer operation manual for proper calibration and operation of application equipment.

Echelon 0.16% Granular Herbicide on Fertilizer application may cause temporary discoloration to exposed leaf surfaces on certain cultivars of zoysiagrass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply this product on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.

Echelon 0.16% Granular Herbicide on Fertilizer has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties and or cultivars have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply this product to a small area prior to treatment of larger areas.

Application Rates

| Turfgrass Use Rate Conversion | | | | |
|-------------------------------|----------------------------|-------------------------|-----------------------|----------------------|
| lbs product per 1000 sq ft | lbs Sulfentrazone per acre | lbs Prodiamine per acre | lbs total ai per acre | lbs product per acre |
| 3.60 | 0.08 | 0.17 | 0.25 | 157 |
| 5.39 | 0.125 | 0.25 | 0.375 | 235 |
| 8.1 | 0.19 | 0.38 | 0.57 | 355 |
| 10.8 | 0.25 | 0.50 | 0.75 | 470 |
| 16.1 | 0.375 | 0.750 | 1.125 | 700 |

When used as directed under the conditions described, apply this product at 157 - 700 lbs product per acre (3.60 to 16.1 lbs product per 1000 sq ft) per application.

The maximum single and annual application rate is 700 lbs product per acre (equivalent to 0.375 lbs sulfentrazone and 0.750 lbs prodiamine per acre) per calendar year.

Refer to Table 1 for the application rates for each grass type.

Table 1. Application Rates for Tolerant grasses

| Grass Type | Use Rates <i>Do not exceed maximum use rates per turf species in a single application</i> | |
|--|---|--|
| | lbs product / 1000 sq ft | lbs product / acre |
| Cool Season Grasses | | |
| Bentgrass, creeping ¹ Fescue, fine ² (<i>Festuca rubra</i>) Ryegrass, perennial (<i>Lolium perenne</i>) | 3.60 – 5.39 ³ | 157 – 235 ³ |
| Bluegrass, Kentucky (<i>Poa pratensis</i>) Fescue, tall ² (<i>Festuca arundinacea</i>) | 8.1 – 10.8 ³ | 355 - 470 ³ |
| Sequential Application for Mixed Cool Season Turf/Cool Season Blends consisting of: | | |
| Fescue, fine ² (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>), Bluegrass, Kentucky (<i>Poa pratensis</i>) Fescue, tall ² (<i>Festuca arundinacea</i>) | Initial application 5.39 | Initial application 235 |
| | Follow-up application 3.60 - 5.39 30- 60 days later | Follow-up application 157 - 235 30 – 60 days later |
| Warm Season Grasses | | |
| Bahiagrass (<i>Paspalum notatum</i>) Buffalograss (<i>Buchloe dactyloides</i>) Carpetgrass (<i>Axonopus affinis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kikuyugrass (<i>Pennisetum clandestinum</i>) Seashore Paspalum (<i>Paspalum vaginatum</i>) Zoysiagrass (<i>Zoysia japonica</i>) | 8.1 – 10.8 ³ | 355 - 470 ³ |
| Bermudagrass (<i>Cynodon dactylon</i>) & hybrids | 10.8 – 16.1 ³ | 470 – 700 ³ |
| <p>1 If bentgrass is mowed lower than 0.5 inches do not apply this product.</p> <p>2 Use of this product on certain varieties of Chewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury.</p> <p>3 This product may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate per turf type or 700 lbs product per acre (equivalent to 0.375 lbs sulfentrazone and 0.750 lbs prodiamine per acre), whichever is lower.</p> | | |

Table 1A. Applications for St. Augustinegrass

- Application of this product to St. Augustinegrass may cause temporary discoloration to exposed leaf surfaces of St. Augustinegrass. Treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing.
- Do not apply this product to St. Augustinegrass if temporary discoloration cannot be accepted.

| Grass Type | Use Rates <i>Do not exceed maximum use rates per turf species in a single application</i> | |
|--------------------------|---|--|
| | lbs product / 1000 sq ft | lbs product / acre |
| Warm Season Grass | | |
| <i>St. Augustine</i> | Initial application 8.1 | Initial application 355 |
| | Follow-up application 8.1 45- 60 days later | Follow-up application 355 45 – 60 days later |

Preemergence Control of Annual, Biennial & Perennial Broadleaf Weeds and sedges

This product, applied alone or in recommended tank mixtures, will provide control of the following weeds for pre-emergence applications.

Table 2. Weed Controlled with Preemergence Application

| Common Name | Scientific Name |
|--|------------------------------------|
| Barnyardgrass | <i>(Echinochloa crus-galli)</i> |
| Bluegrass annual ¹ | <i>(Poa annua)</i> |
| Carpetweed | <i>(Mollugo verticillata)</i> |
| Carolina Geranium | <i>Geranium carolinianum</i> |
| Chickweed, common ² | <i>(Stellaria media)</i> |
| Chickweed, mouseear (from seed) | <i>(Cerastium vulgatum)</i> |
| Crabgrass (large, Smooth) ³ | <i>(Digitaria spp.)</i> |
| Crowfootgrass | <i>(Dactyloctenium aegyptium.)</i> |
| Cupgrass, Woolly | <i>(Eriochloa villosa)</i> |
| Foxtails, Annual | <i>(Setaria spp.)</i> |
| Goosegrass ⁴ | <i>(Eleusine indica)</i> |
| Hairy Bittercress | <i>Cardamine hirsute</i> |
| Henbit ² | <i>(Lamium amplexicaule)</i> |
| Itchgrass | <i>(Rottboellia exaltata)</i> |
| Knotweed, prostrate | <i>(Polygonum aviculare)</i> |
| Kochia | <i>(Kochia scoparia)</i> |
| Kyllinga, green | <i>(Kyllinga brevifolia)</i> |
| Kyllinga, false | <i>(Kyllinga gracillima)</i> |
| Lambsquarters, common | <i>(Chenopodium album)</i> |
| Lovegrass | <i>(Eragrostis spp.)</i> |
| Nutsedge, yellow | <i>(Cyperus esculentus)</i> |
| Panicum (Texas, Fall) | <i>(Panicum spp.)</i> |
| Pigweed, Redroot | <i>(Amaranthus retroflexus)</i> |
| Purslane, Common | <i>(Portulaca oleracea)</i> |
| Pusley, Florida | <i>(Richardia scabra L.)</i> |
| Sedge, globe | <i>(Cyperus globulosus)</i> |
| Sedge, cylindrical | <i>(Cyperus retrorsus)</i> |
| Sedge, Surinam | <i>(Cyperus surinamensis)</i> |
| Sedge, Texas | <i>(Cyperus polystachyos)</i> |

| | |
|--|------------------------------------|
| Shephardspurse ² | (<i>Capsella bursa-pastoris</i>) |
| Signalgrass, broadleaf | (<i>Brachiaria playphylla</i>) |
| Speedwell | (<i>Veronica spp.</i>) |
| Sprangletop | (<i>Leptochola spp.</i>) |
| Spurge, prostrate | (<i>Euphorbia humistrata</i>) |
| Witchgrass | (<i>Panicum capillare L.</i>) |
| Woodsorrel, yellow (from Seed) | (<i>Oxalis stricta</i>) |
| <ol style="list-style-type: none"> 1. <i>Poa annua</i> is a winter annual. Applications of this product should be made in August or September to established, non overseeded turf before <i>Poa Annua</i> germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings. 2. Apply this product in late summer, fall or winter before weed germinates for control. 3. <u>Crabgrass control in cool season grasses from fall application:</u> Areas where the ground is subject to winter freezes, this product may be applied in the fall at rates listed in Table 1 (157 - 470 lbs product per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring. 4. In areas of extended growing season and heavy pressure weed control will be enhanced by second application of this product 60-90 days after initial application. | |

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds and sedges

This product, applied alone or in recommended tank mixture, will provide control of the following weeds for post emergence applications.

Table 3. Weeds Controlled or Suppressed When Application Made Postemergence to Weeds

| Common Name | Scientific Name | Controlled | Suppressed |
|------------------------------|----------------------------------|------------|------------|
| Bittercress | (<i>Cardamine spp.</i>) | | X |
| Black medic | (<i>Medicago lupulina</i>) | | X |
| Buttercups | (<i>Ranunculus spp.</i>) | | X |
| Carolina geranium | (<i>Geranium carolinianum</i>) | 1 | |
| Carpetweed | (<i>Mollugo verticillata</i>) | | X |
| Chickweed, common | (<i>Stellaria media</i>) | 1 | |
| Chickweed, mouseear | (<i>Cerastium vulgatum</i>) | | X |
| Cinquefoil | (<i>Potentilla spp.</i>) | | X |
| Clover | (<i>Trifolium spp.</i>) | | X |
| Crabgrass (Large and Smooth) | (<i>Digitaria spp.</i>) | 2 | |
| Cudweed | (<i>Gnaphalium spp.</i>) | | X |
| Dandelion | (<i>Taraxacum officinale</i>) | | X |
| Dock, Curly | (<i>Rumex crispus</i>) | | X |
| Evening primrose | (<i>Oenothera biennis</i>) | | X |
| Fiddleneck | (<i>Amsinckia spp.</i>) | | X |
| Filaree | (<i>Erodium spp.</i>) | | X |
| Goldenrod | (<i>Solidago spp.</i>) | 1 | |
| Goosegrass | (<i>Eleusine indica</i>) | 2 | |
| Ground ivy | (<i>Glechema hederacea</i>) | 1 | |
| Henbit | (<i>Lamium amplexicaule</i>) | | X |
| Knotweed, prostrate | (<i>Polygonum aviculare</i>) | | X |
| Kochia | (<i>Kochia scoparia</i>) | 1 | |
| Lambsquarters, common | (<i>Chenopodium album</i>) | | X |

| | | | |
|---|--------------------------------------|---|---|
| Lawn burweed | (<i>Soliva pterosperma</i>) | | X |
| Lespedeza, common | (<i>Lespedeza striata</i>) | | X |
| Mallow, common | (<i>Malva neglecta</i>) | | X |
| Nutsedge, Yellow | (<i>Cyperus esculentus</i>) | C | |
| Parsley piert | (<i>Alchemilla arvensis</i>) | 1 | |
| Pigweed, Redroot | (<i>Amaranthus retroflexus</i>) | 1 | |
| Pigweed, Tumble | (<i>Amaranthus albus</i>) | 1 | |
| Pineapple weed | (<i>Matricaria matricarioides</i>) | | X |
| Plantain, buckhorn | (<i>Plantago lanceolata</i>) | 1 | |
| Puncture weed | (<i>Tribulus terrestris</i>) | | X |
| Purslane, common | (<i>Portulaca oleracea</i>) | | X |
| Pusley, Florida | (<i>Richardia scabra</i>) | 1 | |
| Redweed | (<i>Melochia corchorifolia</i>) | | X |
| Rocket, London | (<i>Sisymbrium irio</i>) | | X |
| Smartweed, Pennsylvania | (<i>Polygonum pensylvanicum</i>) | 1 | |
| Sorrel, Red | (<i>Rumex acetosella</i>) | | X |
| Speedwell | (<i>Veronica spp.</i>) | 1 | |
| Spurge, (annuals) | (<i>Euphorbia spp.</i>) | 1 | |
| Spurge, prostrate | (<i>Euphorbia humistrata</i>) | 1 | |
| Spurge, spotted | (<i>Euphorbia maculata</i>) | 1 | |
| Star of Bethlehem | (<i>Ornithogalum umbellatum</i>) | 1 | |
| Velvetleaf | (<i>Abutilon theophrasti</i>) | | X |
| Violet, wild | (<i>Viola pratincola</i>) | | X |
| Wild garlic | (<i>Allium vineale</i>) | | X |
| Wild onion | (<i>Allium canadense</i>) | | X |
| Woodsorrel, creeping | (<i>Oxalis corniculata</i>) | | X |
| Woodsorrel, yellow | (<i>Oxalis stricta</i>) | 1 | |
| <p>1. Weeds are suppressed at lower label rates (<470 lbs product per acre). For optimum control apply rates of at least 470 lbs product per acre in a single application. Do not exceed the application rate specified for the turf species in Table 1.</p> <p>2. This product controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.</p> | | | |

Application to Reseeded, Overseeded or Sprigged areas

Reseeding, overseeding or sprigging of treated areas within three to twelve (3-12) months after application of this product could inhibit the establishment of desirable grasses. [See table for exact intervals from application].

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

Table 4. Overseeding Intervals after Product Applications

| Total Amount of Product Applied lbs product / acre | Interval (Months Before Overseeding) | | |
|---|--------------------------------------|------------|-------|
| | North | Transition | South |
| 157 - 235 | 3 | 3 | 3 |
| 355 - 470 | 4 | 4 | 4 |
| 700 | 6 | 5 | 5 |

Sod Production

Sod should be established for at least six (6) months before an application of this product.

Do not apply this product within three (3) months of harvest.

TURFGRASS USE RESTRICTIONS

Do not apply to golf course putting greens or tees.

Do not use on turfgrasses other than those listed on this label.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply directly to landscape ornamental foliage or ornamental beds containing dormant bulbs or non-woody perennials.

Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are *desirable* species.

Do not apply to creeping bentgrass if mowing height is less than 0.5 inch.

Do not harvest treated sod before 90 days after application.

Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.

Do not apply this product through any type of irrigation system.

Do not apply by air.

ADDITIONAL USE INSTRUCTIONS

Railroad Rights-of-Way

Controls many weeds and maintains bare ground on railroad rights-of-way, including railroad yards, railroad crossings and railroad bridge abutments.

Highway, Roadside, Pipeline and Utility Rights-of-Way

Controls many weeds and maintains bare ground in highway, roadside, pipeline and utility rights-of-way. Such areas would include guardrails, road shoulders, electric utility substations, pipeline pumping stations, around electric transmission towers, around distribution line poles plus other areas where complete vegetation control is desired.

Industrial Areas, Fence Rows and Other Non-crop Sites

Controls weeds and maintains bare ground in industrial areas including production facilities, tank farms, storage areas, parking areas, lumber yards, airports, military installations, along fence rows, and in similar non-crop sites where complete vegetation control is needed.

METHOD AND RATE OF APPLICATION FOR NON-TURFGRASS USES

| Use rate conversion | | | | |
|----------------------------|----------------------------|-------------------------|-----------------------|----------------------|
| lbs product per 1000 sq ft | lbs Sulfentrazone per acre | lbs Prodiamine per acre | lbs total ai per acre | lbs product per acre |
| 10.8 | 0.25 | 0.50 | 0.75 | 470 |
| 16.1 | 0.375 | 0.750 | 1.125 | 700 |

For residual control of germinating weeds in non-crop land, apply as a broadcast treatment using 470 - 700 lbs product per acre (10.8 – 16.1 lbs product per 1000 sq ft) by ground in a suitable granular applicator.

The maximum single and annual application rate is 700 lbs product per acre (equivalent to 0.375 lbs sulfentrazone and 0.750 lbs prodiamine per acre) per calendar year.

DO NOT apply this product to soils classified as sand with less than 1% Organic Matter.

Timing

For best results, apply this product alone or in combination with other herbicides for residual control of weeds in late summer, fall, or early spring to ensure adequate moisture for soil activation.

Weeds Controlled

Echelon 0.16% Granular Herbicide on Fertilizer when applied postemergence at 470 - 700 lbs product per acre, will control the following weeds in non-cropland areas (Table 5). Use the higher labeled rates to extend length of control. Use the higher rates on sites with fine soil textures and on sites with more than 2% organic matter.

Table 5. Weeds Controlled from Preemergence Application to Bare Ground Prior to Weed Seed Germination

| Common Name | Scientific Name |
|---------------------------------|-----------------------------------|
| Barnyardgrass | <i>(Echinochloa crus-galli)</i> |
| Bluegrass annual | <i>(Poa annua)</i> |
| Carpetweed | <i>(Mollugo verticillata)</i> |
| Carolina Geranium | <i>Geranium carolinianum</i> |
| Chickweed, common | <i>(Stellaria media)</i> |
| Chickweed, mouseear (from seed) | <i>(Cerastium vulgatum)</i> |
| Crabgrass (large, Smooth) | <i>(Digitaria spp.)</i> |
| Crowfootgrass | <i>(Dactyloctenium aegyptium)</i> |
| Cupgrass, Woolly | <i>(Eriochloa villosa)</i> |
| Foxtails, Annual | <i>(Setaria spp.)</i> |
| Goosegrass | <i>(Eleusine indica)</i> |

| | |
|--------------------------------|----------------------------------|
| Hairy Bittercress | <i>Cardamine hirsuta</i> |
| Henbit | <i>(Lamium amplexicaule)</i> |
| Itchgrass | <i>(Rottboellia exaltata)</i> |
| Knotweed, prostrate | <i>(Polygonum aviculare)</i> |
| Kochia | <i>(Kochia scoparia)</i> |
| Kyllinga, green | <i>(Kyllinga brevifolia)</i> |
| Kyllinga, false green | <i>(Kyllinga gracilima)</i> |
| Lambsquarters, common | <i>(Chenopodium album)</i> |
| Lovegrass | <i>(Eragrostis spp.)</i> |
| Nutsedge, yellow | <i>(Cyperus esculentus)</i> |
| Nutsedge, purple | <i>(Cyperus rotundus)</i> |
| Panicum (Texas,Fall) | <i>(Panicum spp.)</i> |
| Pigweed, Redroot | <i>(Amaranthus retroflexus)</i> |
| Purslane, Common | <i>(Portulaca oleracea)</i> |
| Pusley, Florida | <i>(Richardia scabra L.)</i> |
| Sedge, globe | <i>(Cyperus globulosus)</i> |
| Sedge, cylindrical | <i>(Cyperus retrorsus)</i> |
| Sedge, Surinam | <i>(Cyperus surinamensis)</i> |
| Sedge, Texas | <i>(Cyperus polystachyos)</i> |
| Shephardspurse | <i>(Capsella bursa-pastoris)</i> |
| Signalgrass, broadleaf | <i>(Brachiaria playphylla)</i> |
| Speedwell | <i>(Veronica spp.)</i> |
| Sprangletop | <i>(Leptochola spp.)</i> |
| Spurge, prostrate | <i>(Euphorbia humistrata)</i> |
| Witchgrass | <i>(Panicum capillare L.)</i> |
| Woodsorrel, yellow (from Seed) | <i>(Oxalis stricta)</i> |

USE RESTRICTIONS (for Non-Turfgrass uses)

Do not apply this product through any type of irrigation system.
Do not apply by air.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Reseal open bag by folding top down and securing. Store in a cool, dry place and avoid excess heat.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. **Confine spills. Call CHEMTREC (Transportation and spills): (800) 424-9300.**

To confine spill: If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling:

[Paper and Plastic Bags:] Non-refillable. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALES AND LIMITATIONS OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Turf or plant injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC – Trademark of FMC Corporation

2021 FMC Corporation. All rights reserved

[Optional State Specific Language]

{Note to Reviewer: These statements are typically added right after the Fertilizer composition section. Typically states have not raised questions concerning where these statements are on the labels, as long as the information is present. Only states where the product will be registered will be included on the commercial label.}

FLORIDA APPLICATIONS: For use on golf courses or other athletic turf in FL, we recommend you follow SL191, “Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Base on Mehlich I Extractant”, available on the internet at <http://edis.ifas.ufl.edu/SS404>.

FLORIDA APPLICATIONS: For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf

Optional State Specific Language for Virginia

In Virginia this lawn/turf fertilizer contains phosphorous and is intended only for nonagricultural use on (1) turf during its first growing season, (2) on turf areas being repaired or renovated, and (3) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not intended for the routine maintenance of turf.

Optional State Specific Language for Maryland

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article. Annotated Code of Maryland performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.