

279-3337 (01-26-2010 (

1/8.1



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Callista O. Chukwunenye
FMC Corporation
1735 Market St.
Philadelphia, PA 19103

JAN 26 2010

Dear Dr. Chukwunenye:

SUBJECT: Master Label Amendment
Spartan Charge Herbicide
EPA Registration No. 279-3337
Your Application Dated July 31, 2009

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

1. Remove the word "General" in all headings (pages 5 and 12), and replace with "Product".

A stamped copy is enclosed for your records. Please submit one (1) copy of your final printed labeling before you release the product for shipment. This amended labeling supersedes all previously accepted ones.

Sincerely yours,

A handwritten signature in black ink, appearing to read "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

2/31

SPARTAN CHARGE

For Use Only by Individuals/Firms Certified
As Licensed Pesticide Applicators

EPA Reg. No. 279-3337

EPA Est. XXX

Active Ingredient: (1)	By Wt.
Carfentrazone-ethyl*	3.53%
Sulfentrazone**	31.77%
Other Ingredients:	64.7%
Total:	100.0%

*SPARTAN CHARGE Herbicide contains 0.35 pounds per US gallon of the active ingredient Carfentrazone-ethyl.

** SPARTAN CHARGE Herbicide contains 3.15 pounds per US gallon of the active ingredient Sulfentrazone.

U.S. Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID (2)

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER (3)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

SPARTAN CHARGE IS FORMULATED AND PACKAGED IN USA.

FROM **FMC**

ATTENTION (4)

-Although this label may appear similar to the label on a product you may have used, there may be important label differences. Users must read, understand and strictly follow all label directions, precautions and restrictions.

-It is the user's responsibility to be sure the product is approved for sale or use on the intended crop and for use in the specific geographic area.

-It is the user's responsibility to be aware of and to follow all State or local precautions or restrictions not appearing on this product label.

-Prior to purchase or use of this product, read the Conditions of Sale and Limitation of Warranty and Liability on page 2 of this label. If the terms and conditions are unacceptable, return the product immediately in the original and unopened container.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN 26 2010
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

279-3337

TABLE OF CONTENTS

Section Title/ (Section Number)	Page
Active Ingredient (1)	1
Agricultural Use Requirements (9)	3
Application Information (20)	8
Attention (4)	1
Cabbage (38)	26
Conditions of Sale, Warranty and Liability (11)	3
Corn (29)	15
Crop Rotation Intervals (25)	11
Directions for Use (14)	5
Dry Shelled Beans and Peas (34)	21
Environmental and Soil Factors (17)	6
Environmental Hazards (7)	2
Fallow Systems (28)	13
First Aid (2)	1
General Information (15)	5
General Postemergent Weeds Controlled (27)	12
Horseradish (39)	27
Hot Line Number (3)	1
Label Tracking (41)	30
Lima Beans, Succulent (35)	23
Maximum Annual Use Rate (23)	10
Mixing and Loading Instructions (19)	7
Peanut (37)	25
Physical/Chemical Hazards (8)	3
PPE Requirements (6)	2
Precautionary Statements (5)	2
Potato (30)	16
Rate Conversion Chart (24)	11
Replanting Instructions (26)	12
Restricted Entry Interval (REI) (10)	2
Resistance Management (13)	5
Soil Classification Chart (18)	6
Sod production (40)	28
Soybeans (31)	17
Spray Equipment Clean-out (22)	10
Spray Drift Reduction Advisory (21)	9
Storage and Disposal (12)	4
Sugarcane (32)	19
Sunflowers (33)	20
Tobacco (36)	24

PRECAUTIONARY STATEMENTS (5)

Hazards to Humans and Domestic Animals

Caution

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE) (6)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves made of waterproof material such as polyethylene or polyvinyl chloride, and shoes plus socks. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards (7)

This pesticide is toxic to algae, marine/estuarine invertebrates, and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface Water Advisory

This product can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

Physical/Chemical Hazards (8)

Do not use or store near heat or open flame.

AGRICULTURAL USE REQUIREMENTS (9)

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. (10)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, chemical resistant gloves, and shoes plus socks.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY (11)

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent consistent with applicable law, buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

5/21

Storage and Disposal (12)

Do not contaminate water, food or feed by storage or disposal.

Pesticide storage

Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool dry place and avoid excess heat. Do not store below 32F degrees.

In Case of Spill

Avoid contact. Isolate areas and keep out animals and unprotected persons.

To Confine Spills.

Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a large holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: (For containers greater than 5 gallons) Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. (For containers 5 gallons or less) Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Triple rinse (or equivalent). Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Containers - Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

RESISTANCE MANAGEMENT (13)

Some weeds are known to develop resistance to herbicides that have been used repeatedly. While the development of resistance is well understood, it is not easily predicted. Therefore herbicides should be used in conjunction with resistance management strategies in the area. Consult the local or State agricultural advisors for details. If weed resistance should develop in the area, this product used alone may not continue to provide sufficient levels of weed control. If the reduced levels of control cannot be attributed to improper application timing, unfavorable weather conditions or abnormally high weed pressure, a resistant strain may have developed.

To reduce the potential for weed resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the recommended rates and in accordance with the use directions. Do not use less than recommended label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when weeds are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

DIRECTIONS FOR USE (14)

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION (15)

SPARTAN CHARGE is a selective herbicide that provides postemergent contact and soil residual weed control. SPARTAN CHARGE may be applied as a burndown prior to planting, early preplant, or as a preemergent application before or after weed emergence for control of susceptible broadleaf weeds. SPARTAN CHARGE is a 3.5 pound per gallon suspoemulsion containing the active ingredients carfentrazone-ethyl and sulfentrazone. Applications of SPARTAN CHARGE must be made before crop seed germination to prevent injury to the emerging crop seedlings. When applications after planting are delayed, injury may occur if seeds are germinating or if they are located near the soil surface.

Observe all instructions, crop restrictions, mixing directions, application precautions, replanting directions, rotational crop guidelines and other label information of each product when tank mixing with SPARTAN CHARGE. In addition to general application information, refer to the specific directions of use for a particular crop/use pattern as set forth below.

Proper Handling Instructions

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Do not apply this product through any type of irrigation system.

Do not use flood irrigation to apply or incorporate this product.

This product must be used in a manner which will prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

GENERAL APPLICATION INSTRUCTIONS (16)

SPARTAN CHARGE is to be mixed with water, liquid fertilizer, or mixtures of water and liquid fertilizer and applied in fallow systems or as a preplant burndown or preemergence herbicide to labeled crops. SPARTAN CHARGE provides postemergent contact and soil residual control of susceptible weed species.

Emerged, susceptible broadleaf weeds are easiest to control when they are small (less than 3 inches tall) and actively growing. Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved postemergent weed control will be poor. Always use the higher application recommended rate of this product, for the appropriate soil texture and organic matter, when weed growth is dense or heavy, or when weeds are growing in an undisturbed or non-cultivated area. Reduced weed control may occur if weeds are experiencing drought stress, disease or insect damage, or when weeds are thickly covered with dust. For control of weeds not listed on this label SPARTAN CHARGE may be tank-mixed with other herbicides such as glyphosate. Read and follow all manufacturers' label directions ~~recommendations~~ for the companion herbicide(s)

except for specific use directions ~~recommendations~~ on this label. The use of a quality spray adjuvant is required for optimum control of emerged weeds. Refer to the individual crop ~~recommendation~~ sections of this label for specific adjuvant type and use rates.

The residual activity of SPARTAN CHARGE applications requires adequate moisture for herbicidal activation. The amount of residual activity is dependent on several factors. These factors include, but are not limited to, existing soil moisture at application, soil type, organic matter, and till. Where irrigation is not available and rainfall has not provided activation, particularly for surface applications of SPARTAN CHARGE, a shallow incorporation (less than 2") is recommended for destruction of any existing weeds and to incorporate SPARTAN CHARGE. Herbicide incorporation will initiate the process of activation with existing soil moisture. In circumstances where rainfall has not occurred and/or irrigation is not possible, alternative or additional weed management practices may be required.

Under normal growing conditions, SPARTAN CHARGE exhibits excellent crop safety. Soil applications of SPARTAN CHARGE must be made before crop seed germination to prevent injury to the emerging crop seedlings. SPARTAN CHARGE applied after crop emergence will cause severe injury to the crop. Poor growing conditions, such as excessive moisture, cool temperatures, and soil compaction or the presence of various pathogens may impact seedling vigor. Under these conditions, the active ingredients in SPARTAN CHARGE can contribute to crop response. Refer to the specific directions of use for a particular crop/use pattern as set forth below for additional information.

ENVIRONMENTAL AND SOIL FACTORS INFLUENCING SPARTAN CHARGE APPLICATIONS (17)

Do not apply to soils classified as sand with less than 1% organic matter.

The user is required to read and follow the specific SPARTAN CHARGE use directions and restrictions for each crop as defined in subsequent sections of this label. The user is cautioned that some crops and weeds respond differently to SPARTAN CHARGE. This response is governed by the SPARTAN CHARGE application rate, various soil factors and inherent crop sensitivity. See individual crop use sections for specific directions on the use of SPARTAN CHARGE for optimum weed control and crop safety results in each crop.

INFLUENCE OF CLAY, SOIL TYPE, AND PH ON SPARTAN CHARGE USE RATES AND CROP RESPONSE

Following an application of SPARTAN CHARGE to soil, germinating seeds and seedlings take up SPARTAN CHARGE from the soil solution. The amount of SPARTAN CHARGE in the soil solution, and available for weed uptake, is determined primarily by soil type, organic matter, and soil pH. SPARTAN CHARGE adsorbs to the clay and organic matter fractions of soils; effectively limiting the amount of active ingredient immediately available to control weeds. Soils typically increase in clay content through the series from coarse to fine as noted in the following Soil Classification Chart (Table 1).

Table 1. SOIL CLASSIFICATION CHART (18)

COARSE	MEDIUM	FINE
Sand	Sandy clay loam	Silty clay loam
Loamy sand	Sandy clay	Silty clay
Sandy loam	Loam	Clay loam
	Silt loam	Clay
	Silt	

Soil organic matter content can vary widely and independently of soil type and requires an accurate analysis of representative soil samples to determine its content.

Soil pH also exerts a dramatic affect on SPARTAN CHARGE availability in the soil solution. As soil pH increases, SPARTAN CHARGE availability increases. Accurate soil pH information will require an accurate analysis of representative soil samples.

The total amount of SPARTAN CHARGE available, in any given soil, is determined by the interaction of soil type (clay content), % organic matter, and pH. The application timing (relative to the emergence of the crop and weeds) and amount of rainfall and/or irrigation received will ultimately determine, in conjunction with the soil parameters and pH, the amount of SPARTAN CHARGE in soil solution.

Irrigation with highly alkaline water (high pH) following a SPARTAN CHARGE soil application can also significantly increase the amount of SPARTAN CHARGE available in the soil solution. Irrigation with water having a pH greater than 7.5 could result in adverse crop response. This response will ultimately depend on initial SPARTAN CHARGE application rate, timing, amount and pH of irrigation water and sensitivity of the crop and its growth stage when irrigated. The risk of

adverse crop response will lessen with the advance in growth stage among most crops.

The following Crop Specific Use Directions have been designed with specific SPARTAN CHARGE recommendations for each crop based on the soil type, soil organic matter, and soil pH interactions described above. The user is cautioned that crop tolerance and weed control performance are based on strict adherence to these recommendations.

MIXING AND LOADING INSTRUCTIONS (19)

Water or liquid fertilizer solutions may be used as the carrier for SPARTAN CHARGE when applied alone or in tank mixtures with other registered herbicides. A jar test is recommended to determine the compatibility of SPARTAN CHARGE and the fertilizer solution. When mixing with fertilizer solutions it is important to premix SPARTAN CHARGE in clear water. See directions for applying SPARTAN CHARGE alone with liquid fertilizer under section 20

A crop oil concentrate, methylated seed oil, nonionic surfactant (NIS) wetting agent labeled, or other equivalent adjuvant labeled for use with herbicides is required for optimum control of emerged weeds. Read and follow all applicable use directions, precautions and restrictions on the surfactant label.

SPARTAN CHARGE Applied Alone

Select the proper SPARTAN CHARGE application rate from the following tables in the crop section of this label. Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of SPARTAN CHARGE for acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Apply the SPARTAN CHARGE spray mixture immediately after mixing.

Do not store spray mixture.

Do not prepare spray mixtures in nurse tanks.

SPARTAN CHARGE Applied in Tank Mix Combination

Select the proper SPARTAN CHARGE application rate from TIMING AND METHOD OF APPLICATION section of label. Read and follow all applicable use directions, precautions and restrictions on the respective tank mix product labels. To ensure product compatibility, a jar test should be conducted before large volume mixing (see MIXTURE COMPATIBILITY TESTING chart below). Provided the jar test indicates the mixture is compatible, prepare the tank mixture as follows.

Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of SPARTAN CHARGE for the acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, SPARTAN CHARGE and other liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Allow time for complete mixing and dispersion after each addition, adding water as necessary. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Use SPARTAN CHARGE tank mixtures immediately after mixing.

Do not store tank mixtures.

Do not prepare spray mixtures in nurse tanks.

SPARTAN CHARGE Applied Alone with Liquid Fertilizer

When adding SPARTAN CHARGE to a liquid fertilizer carrier, SPARTAN CHARGE should be premixed in clear water before adding to fertilizer solution. Adding SPARTAN CHARGE to fertilizer mixtures without first mixing with water can result in incompatibility.

Fill the spray tank one-half full with fertilizer solution. With agitator operating, add the SPARTAN CHARGE slurry to the spray tank. Use a minimum of one gallon of water for each container of SPARTAN CHARGE. Then add slurry to the spray tank through a 20-35 mesh screen. Rinse container used for pre-mixing and add rinsate to the spray tank. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use SPARTAN CHARGE spray mixture immediately after mixing.

Do not store mixture.

Do not prepare spray mixtures in nurse tanks.

Jar Testing Fertilizer Spray Mixtures

Applications of SPARTAN CHARGE alone, or with recommended tank mixtures, in conjunction with clear liquid fertilizer solutions (28-32% nitrogen only) may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing in full spray tank quantities.

- 1) Add 1 pint of fertilizer solution in a quart jar.

- 2) Add the appropriate amount of herbicide based on the MIXTURE COMPATIBILITY table below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).
- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again after 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, and then follow directions noted below.

MIXTURE COMPATIBILITY TESTING

Herbicide Type	Herbicide Field Use Rate	Amount Herbicide Added Per Pint
Wettable Powder or Dry Flowable	0.5 pound	0.75 teaspoon
	1.0 pound	1.50 teaspoons
	2.0 pounds	3.00 teaspoons
	3.0 pounds	4.50 teaspoons
Emulsified Concentrates	1.0 pint	0.5 teaspoon
	1.0 quart	1.0 teaspoon
Liquid Flowables	2.0 quarts	2.0 teaspoons
	3.0 quarts	3.0 teaspoons

*Based on a spray volume of 25 gallons per acre. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

Adjuvant Requirements Recommendation

The use of methylated seed oil (MSO) or a crop oil concentrate (COC) adjuvant, labeled for use with herbicides, is required for optimum control of emerged weeds. A nonionic surfactant adjuvant and water conditioning agent is recommended when SPARTAN CHARGE is tank-mixed with glyphosate. Read and follow all applicable use directions, precautions and restrictions on the surfactant label.

APPLICATION INFORMATION (20)

Ground Application

Use a boom and nozzle sprayer equipped with the appropriate nozzles and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that that produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population. Thorough coverage is essential for control of susceptible broadleaf weeds. Be aware that overlaps and slower ground speeds while starting, stopping, or turning while spraying may result in excessive application and subsequent crop response.

Continuous agitation is required until all spray mixture has been applied. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow SPARTAN CHARGE spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.

To avoid injury to sensitive crops, spray equipment used for SPARTAN CHARGE applications must be drained and thoroughly cleaned with water CHARGE ammonia before being used to apply other products. See Spray Clean-out Section 22 on page 6.

Avoid all direct, and/or indirect spray contact with non-target plants. Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure.

Do not apply when wind speed favors drift beyond the area of treatment.

Aerial Application

SPARTAN CHARGE may be applied by air using properly calibrated nozzle types and arrangements that will provide optimum coverage while producing minimal amounts of fine droplets. Apply sufficient spray volume to achieve adequate coverage. Apply a minimum of five (5) gallons of finished spray per acre. Do not apply when wind speed favors drift beyond the area intended for treatment.

Runoff and Wind Erosion Precautions

Do not apply under conditions which favor runoff or wind erosion of soil containing SPARTAN CHARGE to non-target areas.

To prevent off-site movement due to runoff or wind erosion:

- Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, allow the soil surface to be settled by rainfall or irrigation.

- Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered ground.
- Do not apply to soils when saturated with water.
- Do not use tail water from the first flood or furrow irrigation of treated fields to treat non target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

SPRAY DRIFT REDUCTION ADVISORY (21)
 AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops.

Where States and local governments have more stringent regulations, they must be observed.

Droplet Size Information

Reduce drift potential by applying large droplets. The optimum drift management strategy is to apply the largest droplets that will provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift when applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity and Temperature Inversions).

VMD – VMD is the expression of the droplet size of the spray cloud. The VMD value means that 50% of the droplets are larger than the expressed value and 50% of the droplets are smaller than the expressed value. Optimum spray clouds should be 450 microns with fewer than 10% of the droplets being 200 microns or smaller.

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows usually produce larger droplets.

Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types narrower spray angles produce larger droplets. Consider using low drift nozzles.

Application Height – Making applications at the lowest height practical reduces exposure of spray droplets to evaporation and wind movement.

Swath Adjustment – Swath adjustment distance must increase with increasing drift potential (higher wind, smaller droplets, etc.)

Wind – Drift potentials are lowest between wind speeds of 3 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications in wind conditions outside of this range could increase the risk of off-target effects and should be avoided. Note that local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity - When making applications in conditions of low relative humidity set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions – Do not apply SPARTAN CHARGE during temperature inversions because the drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the following morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or a smoke generator. Smoke that layers and moves laterally in a concentrated clod (under low wind conditions) indicate an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas – Applications should be made when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species and non-target crops).

Off-Target Movement of SPARTAN CHARGE – Drift of dilute spray mixtures containing SPARTAN CHARGE must be prevented. Observation of the preceding environmental conditions, correct application equipment design, calibration and application practices will significantly diminish the risk of off-target spray drift. SPARTAN CHARGE can cause significant symptomology by drift on to sensitive crops and other plants. This symptomology may manifest initially as discreet, localized spots where contact by SPARTAN CHARGE drift mixtures. Depending on concentration of the spray solution and droplet size (effectively determining the concentration of SPARTAN CHARGE) and also depending on the inherent sensitivity of the plants involved, these spots or lesions may not coalesce. These effects will usually not have lasting effects on plant growth, but will likely reduce the value of affected fruit of foliage where grade or quality is associated with appearance. In severe drift instances with particularly sensitive crops, defoliation of affected foliage could result. Failure to follow these guidelines and environmental prohibitions that then result in off-target movement or drift of SPARTAN CHARGE on to unintended crops or plants, irrespective of severity, constitutes misapplication of this product. FMC accepts no responsibility or liability for potential crop effects that may result from such misapplication of SPARTAN CHARGE.

SPRAY EQUIPMENT CLEAN-OUT (22)

After spraying SPARTAN CHARGE and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure.

1. Drain sprayer tank, hoses, and spray boom and thoroughly rinse the inside of the sprayer tank with clean water to remove sediment and residues. Thoroughly flush sprayer hoses, boom and nozzles with clean water.
2. Fill the tank 1/2 full with clean water, and add appropriate detergent or ammonia (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom, and nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the cleaning solution is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, drain the spray system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately with the detergent or ammonia solution.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State and local regulations and guidelines.

Do not drain or flush equipment on or near desirable trees or plants.

Do not contaminate any body of water including irrigation water that may be used on other crops.

Should small quantities of SPARTAN CHARGE remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

MAXIMUM ALLOWABLE SPARTAN CHARGE USE PER ACRE PER 12 MONTH PERIOD* (23)

Refer to the crop section of this label for specific product use directions.

Crop	Ounces SPARTAN CHARGE Per Acre	Pounds Active SPARTAN CHARGE** Per Acre
Corn	10.2	0.28
Dry peas & beans	10.2	0.28
Fallow	10.2	0.28
Limas (succulent)	7.6	0.21
Soybeans	8.5	0.23
Sunflowers	10.2	0.28
Peanut	12.2	0.33
Potato	10.2	0.28
Sugarcane	15.2	0.42
Tobacco	15.2	0.42
Cabbage	15.2	0.42
Horseradish	10.2	0.28
Sod production	15.2	0.42

*The total allowed usage per twelve-month period includes all applications made to the field per twelve-month interval. This includes fallow treatments, burndown treatments, planting time and all in-season treatments. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

** Based on total active carfentrazone-ethyl and sulfentrazone

RATE CONVERSION CHART (24)

SPARTAN CHARGE		CARFENTRAZONE-ETHYL		SULFENTRAZONE	
Product oz/A	lb ai*	Product oz/A**	lb ai	Product oz/A8***	lb ai
3.75	0.10	0.65	0.01	2.9	0.09
5.75	0.15	1.0	0.015	4.5	0.14
8.5	0.23	1.5	0.02	6.7	0.21
10.2	0.28	1.8	0.03	8.0	0.25
15.25	0.41	2.7	0.04	12.0	0.37

* Total pounds active of sulfentrazone + carfentrazone-ethyl

** Based on Aim 2EC formulation

*** Based on SPARTAN 4F formulation

CROP ROTATIONAL INTERVALS (25)

Shown below are the minimum intervals in months from the time of SPARTAN CHARGE application until SPARTAN CHARGE treated soil may be replanted with the crops listed. When SPARTAN CHARGE is tank mixed with other herbicide(s), refer to all those labels for re-cropping instructions, following the intervals that are the most restrictive. For crops not listed, the interval is 12 months in addition to a successful field bioassay.

The field bioassay is a test strip of the intended crop planted across the previously treated field and grown to maturity. The test strip should include low spots, knolls, and variable pH and soil types. If crop responses are not observed, the crop may be planted the following year.

CROP ROTATION INTERVALS*

CROP	INTERVAL (Months)
Alfalfa	12
Barley	4
Cabbage (transplant only)	Anytime
Canola, Crambe	24
Corn, field	Anytime
Corn, pop	Anytime
Corn, seed	Anytime
Corn, sweet	4
Cotton	12
Dry Shell Peas & Beans	Anytime
Horseradish	Anytime
Lima Beans	Anytime
Mint	Anytime
Peanuts	Anytime
Potatoes	Anytime
Rice	10
Rye	4
Sorghum	10**
Soybeans	Anytime
Sugar Beets	36
Sugarcane	Anytime
Sunflowers	Anytime
Sweet Potatoes	12
Tobacco	Anytime
Triticale	4
Turf	Anytime
Wheat	4

* For all other crops not listed, the rotation interval is a minimum of 12 months.

** 18 month rotation for rates above 10.2 fluid ounces per acre

Hybrid Corn Seed Production

Corn inbred lines grown for hybrid seed production may be injured in the growing season following an application of SPARTAN CHARGE. Inbred lines should be thoroughly tested for crop tolerance before rotating to production scale acreages. FMC will not accept

responsibility for any crop injury on field corn grown for seed following an application of SPARTAN CHARGE.

REPLANTING INSTRUCTIONS (26)

If the initial planting of labeled crops fails to produce a uniform stand, only labeled crops for SPARTAN CHARGE or the tank mix partner; whichever is most restrictive, may be replanted. Do not retreat fields with a second application of SPARTAN CHARGE or other herbicide containing sulfentrazone. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the CROP ROTATION INTERVALS on this label. When replanting use minimum soil tillage to preserve the herbicide barrier and achieve maximum weed control.

GENERAL POSTEMERGENT WEEDS CONTROLLED Pre-Plant Burndown (27)

(Refer to individual crop sections for preemergent weeds controlled).

When used as directed, SPARTAN CHARGE will provide postemergent control of the following weeds (less than 3 inches tall) as specified:

Weeds Controlled	SPARTAN CHARGE use rate fluid ounce (lb ai) per acre
Lambsquarters (up to 3 inches tall)	3.75 (0.10)
Morningglory, ivyleaf (up to 3 leaves)	
Morningglory, pitted (up to 3 leaves)	
Nightshade, Eastern black	
Pigweed, redroot	
Velvetleaf	
Waterhemp (up to 2 inches tall)	
Weeds Controlled	SPARTAN CHARGE use rate fluid ounce (lb ai) per acre
All the weeds controlled at 3.75 fluid ounces per acre (0.10 lb/acre) plus the weeds listed below:	4.75 (0.13)
Cheeseweed	
Filaree, redstem	
Flixweed	
Lambsquarters, common	
Mallow, common	
Morningglory, entireleaf	
Morningglory, ivyleaf	
Morningglory, pitted	
Morningglory, scarlet	
Nightshade, hairy	
Pennycress, field	
Pigweed, smooth	
Sesbania, hemp	
Smartweed (PA), seedling	
Tansymustard	
Waterhemp	
Weeds Controlled	SPARTAN CHARGE use rate fluid ounce (lb ai) per acre
All the weeds controlled at 4.75 fluid ounces per acre (0.13 lb/acre) plus the weeds listed below:	6.0 (0.16)
Amaranth, spiny	
Anoda, spurred	
Bedstraw, catchweed	
Buffalobur	
Carpetweed	
Cocklebur	
Copperleaf, hophornbeam	
Cotton, GMO varieties	

Cotton, volunteer		
Dayflower		
Eclipta		
Fiddleneck, coast		
Groundcherry, smooth (seedling)		
Groundcherry, Wright's		
Jimsonweed		
Kochia		
Rocket, London		
Morningglory, ivyleaf		
Morningglory, tall		
Nightshade, American black		
Nightshade, black		
Sheperdspurse		
Spiderwort, tropical		
Thistle, Russian		
Wallflower, bushy		
Weeds Controlled		SPARTAN CHARGE use rate fluid ounce (lb ai) per acre
All the weeds controlled at 6.0 fluid ounces per acre (0.16 lb/acre) plus the weeds listed below:		8.5 (0.23) – 15.2 (0.375)
Amaranth, Palmer		
Ammania, purple		
Buckwheat, wild		
Burclover		
Filaree, broadleaf		
Filaree, white		
Lettuce, prickly		
Mallow, Venice (up to 2 inches tall)		
Meadowfoam		
Mustard spp.		
Redmaids		
Spurry, com		
Spurry, clover		

FALLOW SYSTEMS (28) (see Table 2 for required recommended application rates).

SPARTAN CHARGE may be used in fallow cropping systems only where crops are seeded and harvested on alternate years for soil moisture conservation using rates recommended in Table 2. Follow crop rotational restrictions when replanting following SPARTAN CHARGE applications.

Table 2

SPARTAN CHARGE Use Rate Table			
Fallow Applications			
Broadcast Rate	Fluid Ounces (lb ai) SPARTAN CHARGE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	3.75(0.10)- 5.0(0.14)	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 6.5(0.18)
1.5-3.0	3.75(0.10)- 5.75(0.16)	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.5(0.23)
>3	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.5(0.23)	6.5(0.18)- 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

Adjuvant Requirements recommendation

For optimum control of emerged weeds a nonionic surfactant, crop oil concentrate, methylated seed oil, or equivalent adjuvant is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints/100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.5 to 2.0% v/v (1.5 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2.0 to 4.0% v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre may be used in addition to the selected NIS, COC, or MSO. When an adjuvant is to be used with this product, FMC recommends use of a Chemical Producers and Distributors Association certified adjuvant.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

For all products used in tank mixes refer to the specific product labels for all restrictions on tank mixing and observe all label precautions, instructions, and rotational cropping restrictions.

Precautions

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General SPARTAN CHARGE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE under specific local conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor. If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

Kochia (ALS and Triazine Resistant)	Pigweed, redroot
Lambsquarters, common	Pigweed, smooth
Morningglory, ivyleaf	Thistle, Russian
Morningglory, tall	Waterhemp, common
Nightshade, Eastern Black	Waterhemp, tall

Restrictions

Do not apply more than 10.2 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent SPARTAN CHARGE runoff from rain or snowmelt that may occur following application.

CORN (Field Corn, Seed Corn, Popcorn)

(29)

Preplant Burndown, Early Preplant, and Preemergence Applications (see Table 3 for recommended application rates).

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to emergence of corn to control or suppress weeds using rates recommended in table 3. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to

16/31

achieve maximum weed control. Apply SPARTAN CHARGE using the rates in Table 3 below.

For applications 14-21 or more days prior to planting, use the mid to high rate in the appropriate rate range for the soil and organic matter type listed in Table 3. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 3

SPARTAN CHARGE Use Rate Table (Corn)			
Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fluid Ounces (lb ai) SPARTAN CHARGE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	3.75(0.10)-	3.75(0.10)-	5.0(0.14)-
	5.75(0.16)	5.75(0.16)	6.7(0.18)
1.5 – 3.0	3.75(0.10)-	5.0(0.14)-	5.75(0.16)-
	5.75(0.16)	7.6(0.21)	8.6(0.23)
>3.0	5.0(0.14)-	5.75(0.16)-	7.6(0.21)-
	7.6(0.21)	8.6(0.24)	10.2(0.28)

*Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.

Adjuvant Requirements recommendation

For optimum control of emerged weeds a nonionic surfactant, crop oil concentrate, methylated seed oil, or equivalent adjuvant is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints/100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.5 to 2.0% v/v (1.5 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2.0 to 4.0% v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre may be used in addition to the selected NIS, COC, or MSO.

For all products used in tank mixes refer to the specific product labels for all restrictions on tank mixing and observe all label precautions, instructions, and rotational cropping restrictions.

Precautions

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General SPARTAN CHARGE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE under specific local conditions. Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

BROADLEAVES	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrifusc</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>

Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Thistle, Russian	<i>Lactuca seriola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
SEDGES	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent SPARTAN CHARGE runoff from rain or snowmelt that may occur following application.

POTATOES (30)

Table 4

Spartan Charge Use Rate Table (Potatoes) Preemergence Application			
Broadcast Rate	Fl oz (lb ai) Spartan Charge per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	3.8(0.1)-	3.8(0.10)-	4.8(0.13)-
	5.7(0.16)	5.7(0.16)	6.7(0.18)
1.5-3.0	3.8(0.10)-	4.8(0.13)-	5.7(0.16)-
	5.7(0.16)	7.6(0.21)	7.6(0.21)
>3	5.7(0.16)-	6.7(0.18)-	7.6(0.21)-
	7.6(0.21)	8.6(0.23)	10.2(0.28)

*Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

Ground and Aerial Applications (30.1)

Apply Spartan Charge by aerial application as a preemergence treatment following planting and after dragoff, but prior to potato emergence. Optimum performance can be achieved if Spartan Charge is applied to the soil surface and either rainfall or overhead irrigation is used to activate the product. If no moisture is received within 7 days following application in areas without irrigation, a shallow incorporation (less than 2 inches) may be needed prior to weed and potato emergence to activate the product. Select the appropriate use rate based on soil texture and organic matter as shown in Table 4 above. For control of emerged weeds at the time of the Spartan application, an appropriate burndown herbicide and adjuvants labeled for potatoes may be tankmixed with Spartan to control these weeds. Do not apply Spartan Charge if the potatoes have emerged from the soil as undesirable crop response may occur. Spartan Charge may be tankmixed with other soil-applied herbicides labeled for use in potatoes to improve weed management and increase weed control spectrum.

Apply Spartan Charge in a minimum of 10 gallons of spray by ground application and 5 gallons of spray by air.

Chemigation Applications (30.2)

Spartan Charge may be applied to potatoes through sprinkler irrigation systems including center pivot, lateral move, end tow, solid set or hand move irrigation systems. Apply Spartan Charge prior to potato emergence using sufficient water (0.25 to 0.5 inch per acre) to provide thorough soil surface coverage, but to avoid runoff of irrigation water. Spartan Charge may be applied with other products labeled for chemigation use in potatoes.

It is important to note that irrigation with highly alkaline water (high pH) following a Spartan Charge soil application may significantly increase the amount of sulfentrazone available in soil solution. Irrigation with water having a pH greater than 7.5 could result in adverse crop response. This response will ultimately depend on initial Spartan Charge application rate, application timing, amount and

18731

pH of irrigation water; the sensitivity of the crop and the crop growth stage when irrigated. The risk of adverse crop response will lessen with advances in the crop growth stage.

Weeds Controlled

When applied, according to directions, Spartan Charge will provide control of:

Amaranth, Palmer	Nightshade, Eastern black
Filaree, redstem	Pigweed, redroot
Kochia (ALS and Triazine Resistant)	Pigweed, smooth
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall

Also control all those weeds which are susceptible to carfentrazone application.

For information on other weeds not listed above, refer to Weeds Controlled section (Table 6) in this label.

Precautions

Potato varieties may vary in their response to herbicide applications. When using Spartan Charge on an untested variety, always determine the crop tolerance before planting. Some potato varieties, including Sangre, Shepody and Snowden, have shown sensitivity to Spartan 4F. Caution should be used when planting these varieties on marginal coarse soils.

These Crop Specific Use directions are based upon the interactive effects of Spartan Charge (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General Spartan Charge Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weeds Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with Spartan Charge. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan Charge under specific local conditions.

Restrictions

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply Spartan Charge after potato emergence from the soil as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre per twelve-month period. The twelve-month period is considered to begin upon the initial Spartan application.

SOYBEANS (31) (Food, feed and Industrial)

Preplant Burndown, Early Preplant, and Preemergence Applications (see Table 5 for recommended application rates).

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or within 3 days after planting soybeans to control or suppress weeds using rates recommended in Table 5. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products, be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 5

SPARTAN CHARGE Use Rate Table (Soybeans)			
Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz SPARTAN CHARGE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	5.75-7.75	7.75-8.5	8.5
1.5-3	7.75-8.5	8.5	8.5
>3	8.5	8.5	8.5
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

Precautions

19/31

When applying SPARTAN CHARGE with other registered herbicides, refer to specific label information on precautions, instructions, limitations, application methods and timings, and weeds controlled.

SPARTAN CHARGE is especially effective against a wide range of economic broadleaf weeds. The same processes that SPARTAN CHARGE affects in these weeds can, under certain conditions, be affected in soybeans. These conditions include high pH (7.5 and above), cool weather, prolonged and excessive moisture, seedling diseases, and any other condition, including poor agronomic practices, that are unfavorable to vigorous crop growth. Such effects in soybeans are often observed as stunting and discoloration. The duration of these effects are somewhat dependent on the duration of the adverse growing conditions. These effects lessen and generally diminish with a return to normal growing conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

BROADLEAVES	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrusc</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Smartweed, PA (seedling)	<i>Polygonum pennsylvanicum</i>
Thistle, Russian	<i>Lactuca serriola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
SEDGES	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 8.5 fluid ounces per acre of SPARTAN CHARGE per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent SPARTAN CHARGE runoff from rain or snowmelt that may occur following application. Do not apply after crop seed germination.

SUGARCANE (32)

Table 6

Spartan Charge Use Rate Table (Sugarcane)			
Planting Time and Lay-by Applications			
Broadcast Rate	Fl Oz (lb ai) per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	5.7 (0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	10.2(0.28)
1.5-3	7.6(0.21)- 10.5(0.29)	10.2(0.28)- 12.8(0.35)	12.8(0.35)
>3	10.2(0.28)- 12.8(0.35)	12.8(0.35)- 15.2(0.42)	15.2(0.42)

20/31

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

Apply Spartan Charge as a broadcast or banded preemergence soil applied treatment for the control of broadleaf weeds, grasses and sedges in sugarcane. Refer to the Spartan Charge Product Use Rate Section and Table 6 for specific use information.

Planting Time Applications (32.1)

Apply Spartan Charge preemergence to newly planted or ratoon sugarcane. Use the higher rate on clay soils and/or soils with organic matter content higher than 2 percent. Apply either by air in a minimum of 5 gallons of spray per acre or by ground equipment in a minimum of 15 gallons of spray per acre. Spartan Charge may be applied with other herbicides registered for use in sugarcane.

Aerial Applications (32.2)

Spartan 4F may be applied by air in a minimum of 5 gallons of finished spray per acre. Spartan Charge may be applied with other herbicides or insecticides registered for aerial application in sugarcane.

Lay-by Applications (32.3)

Apply Spartan Charge as a directed spray to sugarcane at lay-by timing. Use the higher rate on clay soils and/or soils with organic matter content higher than 2 percent. Apply as a directed spray with ground equipment in a minimum of 15 gallons of spray per acre. Spartan Charge may be applied with other herbicides registered for use in sugarcane.

Weeds Controlled

When applied according to directions, Spartan Charge will provide control of:

Morningglory, entireleaf	Morningglory, tall
Morningglory, ivyleaf	Pigweed, red root
Morningglory, red	Nutsedge, yellow

For information on other weeds not listed above, refer to Weed Controlled section (27) in this label.

Precautions

These Crop Specific Use directions are based upon the interactive effects of Spartan Charge (sulfentrazone + carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General Spartan Charge Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with Spartan Charge. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan Charge under specific local conditions.

Restrictions

- Do not apply within 120 days of harvest.
- Do not use on soils classified as sand, which have less than 1% organic matter.
- Do not allow spray to contact crop leaves.
- Do not apply more than 15.2 fluid ounces per acre of Spartan Charge per acre per twelve-month period. The twelve-month period is considered to begin upon the initial Spartan application.

SUNFLOWERS (33)

Fall Application, Preplant Burndown, Early Preplant, and Preemergence Applications (see Table 7 for recommended application rates)

Fall Application

SPARTAN CHARGE may be applied in the fall following crop harvest or in existing fallow fields to control or suppress weeds the following season. The SPARTAN CHARGE Rotational Crop Guidelines in Section 25 must be followed if crops are planted the next season. SPARTAN CHARGE should be applied to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product. Do not mechanically incorporate in the fall or spring after application because this activity may destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils to prevent SPARTAN CHARGE runoff from rain or snow that may occur following application. SPARTAN CHARGE may be tank mixed with herbicides to control emerged weeds. Sequential applications may be needed depending on weed size. In situations where weed size may interfere with SPARTAN CHARGE reaching the soil surface, a separate burndown application prior to the application of SPARTAN CHARGE will be required. Use full, recommended rates of burndown herbicides in combination with SPARTAN CHARGE, or sequential applications as needed. Higher

aerial spray volumes are required when there is a dense weed population or canopy.

SPARTAN CHARGE can be tank mixed with other herbicides. Observe all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

Preplant Burndown, Early Preplant, and Preemergence Applications

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or up to 3 days after planting sunflowers to control or suppress weeds using rates recommended in table 7. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 7

SPARTAN CHARGE Use Rate Table (Sunflowers)			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb ai) SPARTAN CHARGE per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5	3.8(0.10)- 5.0(0.14)	3.8(0.10)- 5.75(0.16)	5.0(0.14)- 6.7(0.18)
1.5-3.0	3.8(0.10)- 5.75(0.16)	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.6(0.23)
>3	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 8.6(0.23)	7.75(0.23)- 10.2(0.28)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.

Precautions

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and pH of 7.8 or higher, or on highly eroded soils, hill tops, or in areas of calcareous outcroppings. SPARTAN CHARGE use rates should be reduced or SPARTAN CHARGE should not be used in those areas. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General SPARTAN CHARGE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

2731

Amaranth, Palmer	Pigweed, red root
Filaree, redstem	Pigweed, smooth
Kochia (ALS and Triazine Resistant)	Sida, prickly
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall
Nightshade, Eastern black	

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, or if the seedling is close to the soil surface as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre of SPARTAN CHARGE per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent SPARTAN CHARGE runoff from rain or snowmelt that may occur following application.

DRY SHELLED BEANS AND PEAS (34)

Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea moth bean, lentil, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lab lab bean; pea (*Pisum*) (includes dry field pea) and pigeon pea (see Table 8 for recommended application rates).

Fall Applications

SPARTAN CHARGE may be applied in the fall as a preplant treatment to control or suppress weeds prior to planting the following spring. SPARTAN CHARGE should be applied to the stubble or soil surface and allow moisture from rainfall or snow to move the product into the soil. Do not mechanically incorporate in the fall or spring as this can destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils or to existing snow cover to prevent SPARTAN CHARGE runoff from rain or snow melt that may occur following application. SPARTAN CHARGE may be tank mixed with other residual soil herbicides that are labeled for fall use on dry bean and dry peas. If weeds are emerged at the time of SPARTAN CHARGE application, use a burndown herbicide such as glyphosate or paraquat at the full-labeled rate in combination with SPARTAN CHARGE or split application as needed. Select the appropriate rate from the table below within the correct soil type and organic matter range. When applying SPARTAN CHARGE in the fall, use a mid to high rate within the rate range for the appropriate soil type and organic matter.

Preplant Burndown, Early Preplant, and Preemergence Applications

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or up to 3 days after planting dry shelled peas and beans to control or suppress weeds. Properly closed seed furrows are required when applying at planting time. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 8

SPARTAN CHARGE Use Rate Table (Dry Shelled Beans and Peas)			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb ai) SPARTAN CHARGE per acre*		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5%	3.0(0.08) – 3.75(0.10)	3.75(0.1) – 5.75(0.16)	3.75(0.10) – 5.75(0.16)
1.5-3.0 %	3.75(0.10) – 5.75(0.16)	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)
>3.0 %	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 8.6(0.23)	6.7(0.18) – 10.2(0.28)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.

Precautions

~~Read Crop Liability Disclaimer (section 12) before using SPARTAN CHARGE in dry peas and beans.~~ Best results are achieved with SPARTAN CHARGE when applications are made early preplant and greater than 14 days before planting.

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1.5%) and pH of 7.8 or higher, or on highly eroded soils (such as hilltops), or in areas of calcareous outcroppings. SPARTAN CHARGE use rates should be reduced or SPARTAN CHARGE should not be used in those areas. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General SPARTAN CHARGE Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

If adequate moisture (1/2" to 1" of rainfall or irrigation) is not received within 7 to 10 days and also if dry conditions persist throughout the growing season, erratic preemergent weed control may result. Additional moisture is needed throughout the growing season to maintain herbicide activity and prevent weed escapes.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

Amaranth, Palmer	Pigweed, red root
Filaree, redstem	Pigweed, smooth
Kochia (ALS and Triazine Resistant)	Sida, prickly
Lambsquarters, common	Thistle, Russian
Morningglory, ivyleaf	Waterhemp, common
Morningglory, tall	Waterhemp, tall
Nightshade, Eastern black	

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or to existing snow cover to prevent SPARTAN CHARGE runoff from rain or snow melt that may occur following application.

LIMA BEANS, SUCCULENT (35)

Preplant Burndown, Early Preplant, and Preemergence Applications (see Table 9 for recommended application rates).

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting lima beans to control or suppress weeds. Properly closed seed furrows are required when applying at planting time or before seed germination. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

For applications 14-21 or more days prior to planting, use the mid to high rate in the appropriate rate range for the soil and organic matter type in Table 9. Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown

24731

herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label (section 20).

Table 9

SPARTAN CHARGE Use Rate Table (Lima Beans, Succulent)			
Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb ai) SPARTAN CHARGE per acre*		
% Organic Matter	Soil Texture		
	Coarse	Medium	Fine
<1.5%	3.0(0.08)- 5.0 (0.14)	3.8(0.10)- 7.75(0.21)	4.8(0.13)- 7.75(0.21)
1.5 – 3.0 %	3.8(0.10)- 5.75(0.16)	5.0(0.14)- 7.75(0.21)	5.7(0.16)- 7.75(0.21)
>3.0 %	5.0(0.14)- 7.75(0.21)	5.75(0.16)- 7.75(0.21)	6.7(0.18)- 7.75(0.21)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.

Precautions

When applying SPARTAN CHARGE to coarse textured soils, it is recommended that growers allow a minimum of 7-14 days from application to planting. Best results are achieved with SPARTAN CHARGE when applications are made early preplant and greater than 14 days before planting.

Some adverse crop response may occur on coarse textured soils with low organic matter (less than 1%) and pH of 7.8 or higher, or on highly eroded soils, or in areas of calcareous outcroppings. SPARTAN CHARGE use rates should be reduced in those areas. Inadequate seed furrow closure or shallow planting (less than 1.0 inch) may result in undesirable crop response. As expected, poor growing conditions such as excessive moisture, low temperatures, soil compaction and diseases may also cause undesirable crop response.

These Crop Specific Use directions are based upon the interactive effects of SPARTAN CHARGE and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General SPARTAN CHARGE product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with SPARTAN CHARGE. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on SPARTAN CHARGE under specific local conditions.

Thorough coverage is essential for control of small susceptible broadleaf weeds. If thorough coverage is not achieved, postemergent weed control will be poor. Optimum broad-spectrum control of annual and perennial weeds requires a tank-mix of with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat.

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

Copperleaf, hophornbeam	Pigweed, redroot
Morningglory, entireleaf	Pigweed, smooth
Morningglory, ivyleaf	

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 7.75 fluid ounces per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or to existing snow cover to prevent SPARTAN CHARGE runoff from rain or snow melt that may occur following application.

TOBACCO (Burley, Flue-Cured and Dark) (36)

Table 10:

Spartan Charge Use Rate Table (Tobacco) Preemergence and Preplant Incorporated Applications			
Broadcast Rate	Fl oz (lb ai) Spartan Charge per acre		
Soil Texture			
% Organic Matter	Coarse	Medium	Fine
<1.5	5.7 (0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	10.2(0.28)
1.5-3.0	7.6(0.21)- 10.2(0.28)	10.2(0.28)- 12.8(0.35)	12.8(0.35)
>3	10.2(0.28)- 12.8(0.35)	12.8(0.35)- 15.2(0.42)	15.2(0.42)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

Spartan Charge may be surface applied or preplant incorporated (to a depth no greater than 2 inches) from 14 days to 12 hours days prior to transplanting tobacco. Incorporating Spartan Charge deeper than 2 inches can result in inconsistent weed control.

Broadcast apply the appropriate Spartan Charge rate from Table 10 above, in a minimum of 10 gallons per acre of water, to the soil prior to transplanting.

Non-Bedded (Fields where raised beds are NOT formed prior to transplanting)

Perform all accepted cultural practices for land preparation, fertilizer/fungicide incorporation, etc. prior to the application of Spartan Charge. Once the field has been prepared for planting, Spartan Charge may be surface applied or lightly preplant incorporated from 14 days to 12 hours prior to transplanting.

If Spartan Charge is surface applied and it is necessary to remove equipment tracks from the field after application but prior to transplanting, any light finishing equipment may be used providing the soil is not disturbed to a depth greater than 2 inches.

If timely cultivations are not performed following a pre-transplant surface application, reduced/unacceptable weed control may occur in the drill.

Bedded (Fields where raised beds ARE formed PRIOR to transplanting)

Apply Spartan Charge to formed beds as a surface application from 14 days to 12 hours prior to transplanting. If it is customary to drag/knock down beds prior to transplanting, this procedure must be performed prior to the Spartan Charge application.

When incorporating prior to bedding, Spartan Charge must be thoroughly and uniformly incorporated to a depth no greater than 2 inches to avoid concentrating Spartan Charge in the bed.

If initial transplanting fails to produce a uniform stand, tobacco may be replanted. DO NOT re-treat field with a second application of Spartan Charge, or any other herbicide containing sulfentrazone. DO NOT re-bed. Re-transplant into previously formed, treated beds.

For broad spectrum and optimum grass weed control a grass herbicide application will be required.

Weeds Controlled

When Applied according to directions, Spartan Charge will provide control of:
Filaree, redstem
Amaranthus, livid
Galinsoga, hairy
Lambsquarters, common
Morningglory, ivyleaf
Morningglory, tall
Pigweed, redroot
Pigweed, smooth
Sida, prickly
Signalgrass, broadleaf
Smartweed, Pennsylvania

PEANUTS (37)

Preplant Burndown, Early Preplant, and Preemergence Applications (see Table 11 for recommended application rates).

Apply SPARTAN CHARGE alone or with other herbicides or liquid fertilizers as a burndown or preemergence treatment prior to planting or within 3 days after planting peanuts to control or suppress weeds using rates recommended in Table 11. Properly closed seed furrows are required when applying at planting time or before seed

26/31

germination. When planting into soil treated preplant with SPARTAN CHARGE, minimize soil disturbance to maintain the herbicide barrier on the soil surface to achieve maximum weed control.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide such as glyphosate, glufosinate, or paraquat. When tank mixing SPARTAN CHARGE with other products, be sure the SPARTAN CHARGE is added to the spray tank water first. For specific mixing instructions refer to the Mixing and Loading instructions section of this label.

Table 11

SPARTAN CHARGE Use Rate Table (Peanuts)			
Fall, Preplant Burndown, Early Preplant, and Preemergence			
Broadcast Rate	Fl oz (lb ai) SPARTAN CHARGE per acre*		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5%	3.0(0.08) – 3.75(0.10)	3.75(0.10) – 5.75(0.16)	3.75(0.10) – 5.75(0.16)
1.5-3.0 %	3.75(0.10) – 5.75(0.16)	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)
>3.0 %	5.0(0.14) – 7.75(0.21)	5.75(0.16) – 7.75(0.21)	6.5(0.18) – 10.2(0.28)
Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories Use higher rates for soils of pH less than 7.0 and lowest rate for pH greater than 7.0 within the rate range.			

Precautions

When applying SPARTAN CHARGE with other registered herbicides, refer to specific label information on precautions, instructions, limitations, application methods and timings, and weeds controlled.

SPARTAN CHARGE is especially effective against a wide range of economic broadleaf weeds. The same processes that SPARTAN CHARGE affects in these weeds can, under certain conditions, be affected in peanuts. These conditions include high pH (7.5 and above), cool weather, prolonged and excessive moisture, seedling diseases, and any other condition, including poor agronomic practices, that are unfavorable to vigorous crop growth. Such effects in peanuts are often observed as stunting and discoloration. The duration of these effects are somewhat dependent on the duration of the adverse growing conditions. These effects lessen and generally diminish with a return to normal growing conditions. **Thorough coverage is essential for control of small susceptible broadleaf weeds.** If thorough coverage is not achieved, postemergent weed control will be poor

When used as directed, SPARTAN CHARGE will provide preemergent control of the following weeds (refer to section 27 for postemergent weeds controlled):

BROADLEAVES	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrus</i>
Morningglory, Ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Smartweed, PA (seedling)	<i>Polygonum pensylvanicum</i>
Thistle, Russian	<i>Lactuca serriola</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
SEDGES	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

Restrictions

Do not apply SPARTAN CHARGE Herbicide after crop emergence, at cracking, or if the seedling is close to the soil surface, as undesirable crop response may occur.

Do not apply more than 10.2 fluid ounces per acre of SPARTAN CHARGE per twelve-month period. The twelve-month period is considered to begin upon the initial SPARTAN CHARGE application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not apply to frozen soils or existing snow cover to prevent SPARTAN CHARGE runoff from rain or snowmelt that may occur following application. Do not apply after crop seed germination.

CABBAGE (Transplanted Only) (38)

Table 12

Spartan Charge Use Rate Table (Cabbage)			
Fall or Spring Early Preplant, Preemergence, and Preplant Incorporated Applications			
Broadcast Rate	Fl oz (lb ai) Spartan Charge per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5%	2.9(0.08)- 3.8(0.10)	3.8(0.10)- 5.7(0.16)	3.8(0.10)- 7.6(0.21)
1.5-3.0 %	3.8(0.10)- 7.6(0.21)	7.6(0.21)- 11.4(0.31)	7.6(0.21)- 11.4(0.31)
>3.0 %	7.6(0.21)- 11.4(0.31)	7.6(0.21)- 15.2(0.42)	7.6(0.21)- 15.2(0.42)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories.
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

Early Preplant (Fall Application or Spring Application) (38.1)

Spartan Charge may be applied in the states of MN, ND, SD, MT, CO, NE, WY, ID, WA, OR, WI, or MI only in the fall or spring preceding the growing season to control weeds prior to or up to the planting or transplanting of cabbage. Spartan Charge may be applied in the spring from 60 days prior to planting up to planting time. Spartan Charge should be applied to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product into the soil. Do not mechanically incorporate in the fall or spring after application as this may destroy the herbicide barrier and weed escapes can occur. Do not apply to frozen soils to prevent Spartan Charge runoff from rain or snow that may occur following application. Spartan Charge may be tankmixed with other burndown herbicides to control emerged weeds in the fall or spring or with residual soil herbicides that are labeled for fall use on cabbage. Use the full, recommended rates of burndown herbicides in combination with Spartan Charge, or split applications as needed. Observe all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

Preplant Incorporated (PPI) (38.2)

Spartan Charge may be applied as a preplant incorporated treatment in the spring prior to transplanting of cabbage. Do not incorporate to depths greater than 2 inches. Spartan Charge can be tankmixed with other burndown or soil-applied herbicides labeled for use in cabbage. Use the full, recommended rates of burndown herbicides or split applications as needed. Observe all precautions, instructions and rotational cropping guidelines of each product's label when tank mixing including all references to potential carryover and crop injury warnings or restrictions.

Transplant Cabbage (38.3)

Spartan Charge may be applied pre-emergence as a broadcast or banded treatment to transplanted cabbage only. Applications should be made broadcast or banded treatment prior to transplanting. Spartan Charge may be applied as a banded treatment into the row middles within 72 hours after transplanting.

Weeds Controlled

When Applied according to directions, Spartan Charge will provide control of:

Galinsoga, hairy	Waterhemp, common
Lambsquarters, common	Waterhemp, tall
Pigweed, redroot	

For information on other weeds not listed above, refer to Weed Controlled section (27) in this label.

Precautions

These Crop Specific Use directions are based upon the interactive effects of Spartan Charge (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General Spartan Charge Product Use

28/31

Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with Spartan Charge. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan Charge under specific local conditions.

Restrictions

Do not apply more than 15.2 fluid ounces per acre of Spartan Charge per application or per twelve-month period. The twelve-month period is considered to begin upon the initial Spartan application.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not incorporate to depths greater than 2 inches.

HORSERADISH (39)

Table 13

Spartan Charge Use Rate Table (Horseradish)			
Fall or Spring Early Preplant, Preemergence, and Preplant Incorporated Applications			
Broadcast Rate	Fl oz (lb ai) Spartan Charge per acre		
	Soil Texture		
% Organic Matter	Coarse	Medium	Fine
<1.5%	2.9(0.08)- 5.7(0.16)	3.8(0.10)- 5.7(0.16)	3.8(0.10)- 5.7(0.16)
1.5-3.0 %	5.7(0.16)- 7.6(0.21)	7.6(0.21)- 10.2(0.28)	7.6(0.21)- 10.2(0.28)
>3.0 %	7.6(0.21)- 9.8(0.27)	7.6(0.21)- 10.2(0.28)	7.6(0.21)- 10.2(0.28)

Refer to the previous information on soil types under the COARSE, MEDIUM, and FINE categories
Use higher rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate range.

Spartan Charge may be applied as an preplant premerge or preplant incorporated treatment by ground in a minimum of 15 gallons of finished spray.

Early Preplant (Fall Application or Spring Application) (MN, ND, SD, MT, CO, NE, WY, ID, WA, OR, WI, MI) (39.1)

Spartan Charge may be applied in the fall or spring preceding the growing season to control or suppress weeds prior to or up to the planting of horseradish. Spartan Charge may be applied in the spring from 60 days prior to planting up to planting. Spartan Charge should be applied to the harvested crop stubble or soil surface without incorporation. Moisture in the form of rain or snow will move and activate the product into the soil. Do not mechanically incorporate in the fall or spring after application as this may destroy the herbicide barrier and weed escapes may occur. Do not apply to frozen soils to prevent Spartan runoff from rain or snow that may occur following application. Spartan 4F may be tankmixed with other burndown herbicides to control emerged weeds in the fall or spring or with residual soil herbicides that are labeled for use on horseradish. Use full, recommended rates of burndown herbicides in combination with Spartan Charge, or split applications as needed. Observe all precautions, instructions, and rotational cropping guidelines of each product label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

Preplant Incorporated (PPI) (39.2)

Spartan Charge may be applied as a preplant incorporated treatment in the spring prior to planting of horseradish. Do not incorporate to depths greater than 2 inches. Spartan Charge can be tankmixed with other burndown or soil-applied herbicides labeled for use on horseradish. Use the full, recommended rates of burndown herbicides or split applications as needed. Observe all precautions, instructions and rotational cropping guidelines of each product's label when tank mixing including all references to potential carryover and crop injury warnings or restrictions.

Pre-Emergence (PRE) (39.3)

Spartan Charge may be applied pre-emergence as a broadcast or banded treatment on horseradish. Applications should be made broadcast prior to planting, broadcast soon after planting but at least 5 days before crop emergence. Spartan Charge may be applied as a banded treatment into the row middles after crop emergence. Use the higher Spartan Charge rates on clay soils and/or soils with greater than 1% organic matter. Spartan Charge may be applied with other pesticides registered for use on horseradish.

Weeds Controlled

When applied according to directions, Spartan Charge will provide control of:

Lambsquarters, common	Pigweed, redroot
Morningglory, ivyleaf	Waterhemp, common
Nutsedge, yellow	Waterhemp, tall

2/31

For information on other weeds not listed above, refer to Weed Controlled section (Table 6) in this label.

Precautions

These Crop-Specific Use directions are based upon the interactive effects of Spartan Charge (sulfentrazone and carfentrazone) and the primary soil and environmental factors, which affect its activity on various weed species and tolerance among crops. The user is required to observe the instructions and guidance previously presented under General Application Instructions, General Spartan Charge Product Use Rates, Rotational Crop Guidelines, Replanting Instructions, Weed Controlled and any other section of this label pertinent to the anticipated crop use. It is important to note that not all varieties or cultivars of a given crop species have been evaluated under treatment with Spartan Charge. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan Charge under specific local conditions.

Restrictions

Do not apply more than 10.2 fluid ounces per acre of Spartan Charge per application or per twelve-month period. The twelve-month period is considered to begin upon the initial Spartan application.

Do not apply directly on the crop after the crop emerges or if the seedling sprouts are close to the soil surface.

Do not use on soils classified as sand, which have less than 1% organic matter.

Do not incorporate to depths greater than 2 inches.

SOD PRODUCTION (40)

Spartan Charge may be applied to established seeded, sodded or sprigged turfgrasses following the second mowing for the control of key grass, sedge and broadleaf weeds. Turf grasses should have developed a good root system, a uniform stand with healthy root systems to fill in the exposed edges prior to application. Sod injury could result from application of this product on sod that is not well established or has been weakened by stresses such as unfavorable weather conditions, diseases, chemical, recent harvesting or mechanical influences.

Turf Grass Tolerance

When applied as directed, the following established turf grasses are tolerant to Spartan Charge herbicide at the listed recommended use rates.

Table 14 Tolerant grasses

Grass Type	Maximum Use Rate For Single Application
Cool Season Grasses **	Fl oz (lb ai) Spartan Charge Per Acre
Bentgrass, creeping	5.1 (0.14)
Fescue, fine * (Festuca rubra) Fescue, tall * (Festuca arundinacea) Ryegrass, perennial (Lolium perenne) Bluegrass, Kentucky (Poa pratensis) Bluegrass, Rough (Poa trivialis)	5.1(0.14)-10.2(0.28)
Warm Season Grasses **	
Bahiagrass (Paspalum notatum) Buffalograss (Buchloe dactyloides) Carpetgrass (Axonopus affinis) Centipedegrass (Eremochloa ophiuoides) Kikuyugrass (Pennisetum clandestinum) Seashore Paspalum (Paspalum vaginatum) Zoysiagrass (Zoysia japonica) Bermudagrass (Cynadon dactylon) Bermudagrass Hybrids (Cyn Bluegrass, St. Augustinegrass (Stenotaphrum secundatum)	10.2(0.28)-15.2(0.42)

* Applications of Spartan Charge to certain varieties of Chewings, Fine Fescue or Tall Fescue may result in undesirable plant response.

** It is important to note that not all varieties or cultivars have been evaluated under treatment with Spartan Charge. Consult university or extension weed management specialists for additional information on specific local varieties or cultivars and any other pertinent information on Spartan Charge under specific local conditions.

Applications to Reseeded, Overseeded or Sprigged Areas

Reseeding, overseeding or sprigging may be done following Spartan Charge applications to turfgrasses. If reseeded, overseeding or sprigging is done within 1 month following a Spartan Charge treatment, the establishment of desirable grasses may be inhibited. Overseeding of bermudagrass with perennial ryegrass may be done two (2) to four (4) weeks following a Spartan Charge application provided slight grass plant response can be tolerated.

Optimum reseeded and overseeding results may be obtained with the use of mechanical or power seeding equipment, and where proper soil cultivation, irrigation and fertilization practices are followed.

Adjuvant use

Good spray coverage is required for optimum control of weeds. Temporary discoloration of some sod species may result from use of surfactant. Use of surfactants is not recommended.

Postemergence Control of Sedges

Spartan Charge may be applied at the rate of four (4) to twelve (12) fluid ounces per acre to established turf grasses for the control or suppression of sedges. Select the correct Spartan Charge use rate from Table 14

When applied as directed, Spartan Charge will provide control or suppression of the following sedges.

Table 15

Common Name	Scientific Name
Kyllinga, green	<i>Kyllinga brevifolia</i>
Kyllinga, false green	<i>Kyllinga gracillima</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, cylindrical	<i>Cyperus retrorsus</i>
Sedge, globe	<i>Cyperus globulosus</i>
Sedge, Surinam	<i>Cyperus surinamensis</i>
Sedge, Texas	<i>Cyperus polystachyos</i>

Purple nutsedge: For optimum control of purple nutsedge, split applications are listed recommended below. Apply 4-8 ounces per acre as an initial application followed by a second application when evidence of actively growing purple nutsedge is visible. Do not exceed the maximum rate per acre based on the turf variety as listed in Table 14; tolerant grasses.

Split Application Rates for Optimum Purple Nutsedge Control

Grass Type	First Application (fl. oz (lb ai) per acre)	Second Application (fl. Oz (lb ai). per acre)
Cool Season Grasses	2.5(0.07)-5.1(0.14)	2.5(0.07)-7.6(0.21)
Warm Season Grasses	5.1(0.14)-7.6(0.21)	5.1(0.14)-7.6(0.21)

Allow 35 days after first application for second application.

Postemergence Control of Grassy Weeds

Spartan Charge will control or suppress specific annual grasses (Table 16) when applied at a rate of 4 to 12 fl oz/acre. Apply the highest rate consistent with the rate needed for turfgrass tolerance in Table 14. Rates lower than 12 fl oz/acre will generally control grasses for at least 60 days. Spartan 4F works best if applied when the annual grasses are small (pre tiller stage) and actively growing.

Table 16

Common Name	Scientific Name
Goosegrass	<i>Elysiue indica</i>

Postemergence Control of Broadleaf Weeds

Spartan Charge herbicide will control or suppress the weeds listed in the broadleaf chart below when applied alone shortly after weeds have emerged. Spartan Charge may be applied at the rate of four (4) to twelve (12) fluid ounces per acre to established turf grasses for the control or suppression of broadleaf weeds. Select the correct Spartan Charge use rate from Table 14. For optimum results, Spartan applications should be made shortly after weeds have emerged.

Spartan Charge may be tankmixed with other herbicides, insecticides and fungicides registered for use on turfgrasses. Read and follow the label recommendations of the tank mix partner to determine turfgrass specie tolerance, use rates and application requirements. Follow all label restrictions, use directions and precautionary statements before use.

When applied as directed, Spartan Charge will provide control or suppression of the following broadleaf weeds.

Broadleaves	Scientific Names
Bittercress	<i>Cardamine spp.</i>
Black Medic	<i>Medicago lupulina</i>
Buttercup	<i>Ranunculus spp.</i>
Carolina geranium	<i>Geranium carolinianum</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Cinquefoil	<i>Potentilla spp.</i>
Clover	<i>Trifolium spp.</i>
Cudweed	<i>Gnaphalium spp.</i>
Dandelion	<i>Taraxacum officinale</i>
Dock, curly	<i>Rumex crispus</i>
Evening primrose	<i>Oenothera biennis</i>
Fiddleneck	<i>Amsinckia spp.</i>
Filaree	<i>Erodium spp.</i>
Garlic, wild	<i>Allium vineale</i>
Goldenrod	<i>Solidago spp.</i>
Ground ivy	<i>Glechema hederasea</i>
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lawn burweed	<i>Soliva pterosperma</i>
Lespedeza, common	<i>Lespedeza striata</i>
Mallow, common	<i>Malva neglecta</i>
Onion, wild	<i>Allium canadense</i>

Parsley piert	<i>Alchemilla arvensis</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Pineapple weed	<i>Matricaria matricariodes</i>
Plantain, buckhorn	<i>Plantago lanceolata</i>
Puncture weed	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Redweed	<i>Melochia corchorifolia</i>
Rocket, London	<i>Sisymbrium ino</i>
Smartweed, PA	<i>Polygonum pensylvanicum</i>
Sorrel, red	<i>Rumex acetosella</i>
Speedwell	<i>Veronica spp.</i>
Spurge, annual	<i>Euphorbia spp.</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Star of Bethlehem	<i>Ornithogalum umbellatum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Violet, wild	<i>Viola pratensis</i>
Woodsorrel, creeping	<i>Oxalis comiculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>

Precautions
 The use of additional surfactants may cause temporary undesirable effects to turfgrasses.

Restrictions
 Sod production areas must be established three (3) months prior to the initial treatment of Spartan Charge.
 Do not apply Spartan Charge to golf course greens or tees.
 Do apply Spartan Charge to turf grasses not listed on this label.
 Do not apply with surfactants without on-site evaluations for spray mixture compatibility and physical effects to turf grasses.
 Do not graze or feed forage harvested from Spartan Charge treated areas.
 Do not apply to landscape ornamental plants or ornamental beds.
 Do not harvest sod within three (3) months of Spartan Charge application.

LABEL TRACKING INFORMATION (41)

Label Code: Spartan Charge 07-31-09
 Replaces Label Code: 10-15-08
 EPA Approval Date:
 Philadelphia, PA 19103 USA
 FMC Corporation
 Agricultural Products Group
 1735 Market Street
 Philadelphia PA 19103
 215-299-6000

SPARTAN CHARGE and FMC — Trademarks of FMC Corporation, Philadelphia, PA 19103 USA

©2009 FMC Corporation All Rights Reserved