



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (H7505C)
1200 Pennsylvania Ave NW
Washington, D.C. 20460

EPA Reg.
Number:

279-3339

Date of Issuance:

APR 18 2008

Term of Issuance: Conditional

Name of Pesticide Product:

Lawn Insect Control 2

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Andy Eimanis
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

- 1. Revise the EPA Registration Number to read, EPA Reg. No. "279-3339".

Signature of Approving Official:

Dani Daniel
Insecticide-Rodenticide Branch
Registration Division (7505P)

Date:

APR 18 2008

2. Rewrite your introductory label statement to read as: **For Residential Lawns, Landscaped areas only. Not for use in commercial green houses or on grass grown for seed or on golf courses and sod farms.**

Failure to add this statement otherwise would cause this label to be classified as a restricted use label due to bifenthrin.

3. Reformat the **Precautionary Statement** for easier reading as follows:

Precautionary Statements

Hazards to Humans and Domestic Animals

Cautions: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid

- If swallowed: Call a poison control center...
- If on skin or clothing: ... Call a poison control center or doctor for treatment advice.
- If in eyes: Hold eyes open ...Call a poison control center or doctor for treatment advice.

4. Use the following Note to Physician:

No specific antidote is available. Treat the patient symptomatically. This product contains a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so must be avoided.

5. Page 3 make the following changes:

- Delete the titles **“Use Directions Option 1** at the top of the page, and delete the second **“Directions for Use”** near the center of the page, this is too confusing and unnecessary.
- Under the heading **Lawn Insect Control 2** change the word **“various”** to the word **“listed”**. Also under the same title and every where throughout the label remove **“ornamental and flower gardens”**. This is not a use site for this label.
- Remove the word **“Garden”** from the title **“Lawns and Garden Applications”**, this is not a use site for this product.

- Make the following changes to the last paragraph on page 3:
 - a. Change “recommended applications rate” to “listed application rate”;
 - b. Change “recommended rates” to “listed rates”
 - c. Insert into the last sentence beginning at 125 lb per acre “per year” on residential lawns.
 - d. At the very end of the paragraph add the following statement: **Avoid mowing lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.**

6. On page add “Restrictions” section:

- **Avoid runoff or puddling or irrigation water following application. Avoid application of Lawn Insect Control 2 to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant. Do not apply more than 250 lb (0.5lb of Imidacloprid and 0.4 lb of bifenthrin per acre per year.**
- **Do not apply by air.**
- **Not for use in nurseries or commercial greenhouse.**

Additional changes for page 4 include:

- Remove the word “do” from the second paragraph. The sentence will now read: **“In New Your State, make a single...”**
- Insert the word “and” between the **Equipment and Settings** in the title concerning the spreader.
- Remove the word “recommended” from the sentence under the heading **Specific Pest Instructions.**

7. Page 5 in the first paragraph following the **Conditions of Sales and Limitation of Warranty and Liability** change the word “should” to the word “must.”

8. Delete page 6 in its entirety.

9. Every where the term “ants” appears add the statement: **“except carpenter ants and pharaoh ants”.**

10. Every where spiders appear add the statement: **“excluding black widow and brown recluse”.**

11. Remove **centipedes** from the label. Centipedes are public health pest and require data.
12. Remove the word "immediately" from the first "Optional Re-entry statement".
13. Make the following changes for the **Speed Claims**:
 - Fasting Acting against insects, spiders, and ticks. **Rewrite to read as "Fast acting against listed insects."**
 - Starts killings (to kill) lawn bugs/insects immediately. Remove the word "immediately"
 - Start killing immediately! Remove the word "immediately"
14. Make the following changes for **Residual claims** on page 10: Long lasting! {Up to} month control {Guaranteed or your money back}
15. Make the following changes on page 11 "**Usage site claims**":
 - Remove "**ornamental and flower gardens**"
 - Remove "**use around outside of homes, porches and patios**"
 - Remove "**for use on ornamentals, and flower beds/gardens**"
16. Within eighteen months of the date of this registration, submit to the Agency the required one year storage stability study (830.6317) for the proposed product under warehouse conditions. The corrosion characteristics study (830.6320) may be carried out concurrently. It is recommended that observations be made at 0, 3, 6, 9, and 12 months.
17. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact me (703) 305-5409.

Enclose:

Lawn Insect Control 2

For Residential Lawns, Landscaped areas, Ornamental & Flower Garden use only.

[icon]
ANT

[icon]
GRUB

[icon]
LAWN

[icon] [icon]
ORNAMENTAL LANDSCAPED AREA

EPA Reg. No. 279- EPA Est. 279-

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....	0.20%
Bifenthrin*	0.16%

OTHER INGREDIENTS:	<u>99.64%</u>
	100.00%

*Cis isomers 97% minimum, trans isomers 3% maximum.

Product is covered by one or more of the following patents: U.S. Patent No. 4,742,060, 5,783,203 and 4,238,505. Other patents pending.

Not for sale, use, or distribution in or into Nassau, Suffolk, Queens, or Kings counties of New York

KEEP OUT OF REACH OF CHILDREN

CAUTION

See (back) panel for additional precautionary statements (and directions for use).

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

APR 18 2008

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

279-3339



FMC Corporation
Agricultural Products Group
Philadelphia PA 19103

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals) Caution

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

First Aid

If swallowed: A poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **If in eyes:** eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Note to Physician:

This product is a pyrethroid. If large amounts have been ingested, the milk, cream or other digestible fats and oils may increase absorption and so should be avoided.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

Faint, illegible text or markings, possibly bleed-through from the reverse side of the page.

Use Directions OPTION 1

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into drink containers.

Product Disposal: Partially filled bag - Securely wrap original bag in several layers of newspaper and put in trash.

Container Disposal: Nonrefillable bag. Do not reuse or refill empty bag. Offer for recycling, if available or dispose of empty bag in trash.

Lawn Insect Control 2 is designed for use as a ready-to-use granular insecticide to control various adult and larvae above and below surfaces of lawns, ornamental & flower gardens. Please read the following "Directions for Use" carefully and fully before applying this product.

The active ingredients in **Lawn Insect Control 2** have sufficient residual activity so that applications for control of subsurface feeders can be made preceding the egg laying activity. The need for an application for control of subsurface feeders can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control of subsurface feeders will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

The active ingredients in **Lawn Insect Control 2** have sufficient knockdown and residual activity to provide curative and residual control of surface feeding pests. Applications for control of surface feeding pests can be made when infestations are anticipated based on historical monitoring of the site, previous records or experience, current season adult trapping or presence of insects at economic thresholds as determined by scouting and/or recommendations of local State extension personnel or other qualified specialists.

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Total amount applied cannot exceed 125 lb (0.25 lb of imidacloprid, 0.2 lb of bifenthrin) per acre per year.

DIRECTIONS FOR USE

LAWN and GARDEN APPLICATIONS

Broadcast **Lawn Insect Control 2** to provide uniform coverage over the treatment area. Locate the pest to be controlled in the following table and follow the recommended application rate to provide excellent control under typical conditions. When treating for multiple pests, apply the higher of the recommended rates. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity. Repeat application should be limited to no more than once per 7 days. Usage rates greater than recommended or multiple applications may be made at the applicators discretion, not to exceed 125 lb per acre on residential lawns .

Pest	Application Rate		
	Lb product per 1000sqft	Lb product per Acre	Lb total ai per Acre
Armyworms Cutworms Sod Webworms	1.1 to 2.9	50 (65)-125	0.18 (0.23) – 0.45
Annual Bluegrass Weevil (Hyperodes) Billbugs Black Turfgrass Ataenius (Adult) Leafhoppers Mealybugs	1.7 to 2.9	75-125	0.27 – 0.45
Subsurface feeder larvae (such as Asiatic garden beetle, Japanese beetle, Black turfgrass ataenius, Northern masked chafer, Black vine weevil Oriental beetle, European chafer, <i>Phyllophaga spp.</i> ,	2.3 to 2.9	100-125	0.36 – 0.45

European crane fly, Southern masked chafer)			
Ants Centipedes Chinch Bugs Fleas (larvae & adults) Imported Fire Ants (adults) Millipedes Mole Crickets (nymphs, & adults) Scorpions Ticks Pillbugs Sowbugs	2.9	125	0.45

In New York State, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application **Lawn Insect Control 2** if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

SPREADER EQUIPMENT SETTINGS

Broadcast application should be performed with a calibrated commercial broadcast spreader to evenly distribute the granules. Approximate settings for several models are provided in the table below.

{Note to PM: Following will be a list of application spreaders and appropriate settings. This list will vary depending on market spreaders. This note will not appear on final label}

SPECIFIC PEST INSTRUCTIONS

For optimum control of particular pests, follow recommended procedures outlined below:

Ants: Mound Treatment, including Fire Ants: Treat ant mounds by applying 1/2 cup of **Lawn Insect Control 2** per mound. Water with sufficient force to break its apex and allow the insecticide to flow into the ant tunnels, but not sufficient to significantly damage the mound that may lead to relocation. For best results, apply in cool weather (65 to 80°F) or in early morning or late evening hours. Treat new mounds as they appear. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide. For optimum control or for heavily infested areas, combine both mound and broadcast treatments

Ants: Broadcast Treatment, including Fire Ants: To control ants actively foraging, uniformly apply **Lawn Insect Control 2** at a rate of 125lb per 1000sqft. over treatment area. For optimum control or for heavily infested areas, combine both mound and broadcast treatments.

Armyworms, Cutworms, and Sod Webworms: Irrigate the treated area with up to 0.1 inches of water immediately after application to activate (release from the granule) the insecticide.

Annual Bluegrass Weevil (Hyperodes) adults: Time application for when adult weevils leave their overwintering sites and move into grass areas.

Billbug adults: Apply when adult billbugs are first observed during April and May.

Black Turfgrass Ataenius adults: Apply during May and July to control the first and second generation of black turfgrass ataenius adults.

Chinch Bugs: Irrigate the treated area with up to 0.25 inches of water immediately after application to activate (release from the granule) the insecticide. Higher applications rates may be required to control populations that contain both nymphs and adults during the summer. In temperate regions, Spring applications will provide control of over-wintered chinch bugs.

Flea larvae: Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release from the granule) the insecticide.

Mole Crickets: Apply as late in the day as possible to moist soil (irrigate before application if necessary), and water in with up to 0.5 inches of water immediately after treatment. Treat grass areas that receive pressure from adult mole crickets in the spring immediately

prior to peak egg hatch when young nymphs are located near the soil surface. Use higher and more frequent applications to control larger, more damaging, nymphs later in the year.

Scorpions: Treat at dusk to prevent outdoor scorpions from entering the building. Treat areas where scorpions reside hidden during the day such as in cracks and voids. Remove trash, debris, or firewood that scorpions may use as resting sites prior to treatment.

Ticks (Including ticks that transmit Lyme Disease and Rocky Mountain Spotted fever): For extended control, do not rely on spot applications only. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Re-treat if ticks are reintroduced from surrounding areas on host animals. Do not allow public use of treated areas during application.

Deer ticks (*Ixodes sp.*) Apply in mid to late-spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

Subsurface feeders: Apply prior to egg laying, followed by sufficient watering or rain to drive active ingredient through the thatch. Avoid applying when soil is waterlogged as adequate distribution will not be achieved.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. EXCEPT AS WARRANTED BY THIS LABEL, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC - Registered Trademark of FMC Corporation
© 2007 FMC Corporation All Rights Reserved

Use Directions Option 2

SEASON-LONG GRUB CONTROL

White grubs can cause extensive damage to turf by feeding on grass roots. Affected turf turns yellow, then brown, and is easily pulled up in pieces with your hands.

ONE APPLICATION STOPS GRUBS ALL SEASON LONG!

Lawn Insect Control 2 works best when applied before grubs hatch. The active ingredient will remain in the root zone of the turf until the eggs of grubs hatch. Newly hatched grubs are highly susceptible to Lawn Insect Control 2 and are killed before they become large enough to cause turf damage.

Lawn Insect Control 2 prevents turf damage by killing grubs when they are newly hatched, before they can cause damage.

For residential use on lawns and landscaped areas.

DO NOT use in vegetable gardens.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

In New York State, this product may NOT be applied to any grass or turf area (lawns within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch)

In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

PRODUCT FACTS

INSECTS CONTROLLED

Use LAWN INSECT CONTROL to kill and control:

- Fleas

Use LAWN INSECT CONTROL to kill and provide * season long (3-month) control of:

- Fire Ants, Ants (including Carpenter, Red Harvester, Pavement, Odorous, Pyramid, Argentine, Pharaoh, Black Turf, White-footed Ant, Centipedes, Chiggers, Crazy Ant, Little Black Ant, Small Honey Ant, Black Turf Ant, Lawn Ant, Cornfield Ant, Ghost Ant, Thief Ant, Acrobat Ant, Citronella Ant, Big-headed Ant, Allegheny Mound Ant), American Dog Ticks, Armyworms (including Lawn Armyworm, Fall Armyworm), Billbugs (including Bluegrass Billbug, Hunting Billbug), Black Turfgrass Ataenius adults, Bristletail, Chinch Bugs, Crickets (including House Cricket, Common Short-tailed Cricket, Indies Short-tailed Cricket, Arizona Cricket, Tropical Cricket, Variable Field Cricket, Fall Field Cricket, Southeastern Field Cricket, Texas Field Cricket, Spring Field Cricket, Northern Wood Cricket, Western Striped Cricket, Eastern Striped Cricket, Sand Field Cricket, Vocal Field Cricket, Southern Wood Cricket), Cutworms (including Black Cutworm, Bronzed Cutworm, Granulate Cutworm, Variegated Cutworm), Dichondra Flea Beetle, Earwigs, European Crane Fly Larvae, Fiery Skipper, Fire Ants [Note to PM: When products is labeled for mound treatments label must state the following: (excluding mound treatments)], Firebrat, Frit Fly, Grasshoppers, Greenbug, Ground Beetle, Ground Pearl, Ladybeetles (including Convergent Ladybeetle, Seven-spotted Ladybeetle, Two-spotted Ladybeetle, Thirteen-spotted Ladybeetle, Twice-stabbed Ladybeetle, Asian Ladybeetle, Mealybug Ladybird), Leafhoppers, Lucerne Moth, Mealybugs, Millipedes, Mole Crickets (including Tawny Mole Cricket, Southern Mole Cricket, Short-Winged Mole Cricket), Pillbugs, Arizona Stripetail Scorpion, Southern Unstriped Scorpion, Lesser Stripetail Scorpion, Yellow Ground Scorpion, California Common Scorpion, Northern Scorpion, Eastern Sand Scorpion, Seed Maggots, Sowbugs, Spiders (including Marbled Cellar Spider, Longbodied Cellar Spider, Shortbodied Cellar Spider, Thinlegged Wolf Spider, Burrowing Wolf Spider, Pirate Wolf Spider), Springtails, Striped Grassworm, Ticks (including Brown Dog Tick and ticks that may transmit Lyme Disease), Two-lined Spittle Bugs, Webworms (including Sod Webworm, Grass Webworm, Tropical Sod Webworm), Weevils (including Hyperodes and Bluegrass) and Wireworms.
- (Sub-surface Feeders:) White Grubs (Larvae of Japanese Beetles, Oriental Beetles, Chafers, Asiatic Garden Beetles, May/June Beetles, Green June Beetles, Sugarcane Beetles and Black Turfgrass Ataenius) and Larvae of Black Vine and Annual Bluegrass Weevils.

WHERE TO USE

Lawn and Landscape areas.

WHEN TO APPLY

Apply early/proactively [in the Spring/Summer] to prevent infestation.

Works best when applied before grubs hatch. For best results, apply once, anytime from Spring through mid-Summer.

(For extended surface feeder (feeding) insect control (into the Fall), follow-up your Spring/Summer treatment with Ortho Bug B Gon Max Insect Killer for Lawns.)

(For extended surface feeder (feeding) insect control (into the Fall), apply Ortho Bug B Gon Max Insect Killer for Lawns.)

DO NOT apply Lawn Insect Control 2 when turf areas are waterlogged. The product may not work effectively under these conditions.

HOW TO APPLY

For best results, treated area should be thoroughly watered immediately after application.

On Lawns (*Illustration*)

- Using a spreader designed for application of granular materials, uniformly apply product. (see spreader settings chart)
- A 10 (20) pound bag will treat up to 3484 (6968) sq. ft.
- For season long control, apply 3lbs of product for every 1000 sq. Re-apply after 3-months if necessary.

On landscape areas (*Illustration*)

- Apply 0.4 to 0.7lb. evenly over each 500 sq. ft. area using a hand held spreader. Repeat if there are signs of renewed insect activity.
- Product can be applied just before planting or after plants emerge.

This product applies 0.25 lb. of imidacloprid, 0.2 lb. bifenthrin per acre per application.

Spreader (Application) Tips

Spreader settings are approximate. You may need to adjust for proper coverage. Apply header strips. Then apply product in the longest direction of lawn. For best results, use a Scotts® spreader.

(Illustrations to correspond with application tips)

Spreader	Spreader Settings Covers X sq. ft.	Directions
Drop Spreader Scotts® AccuGreen®	X	Overlap wheel marks to ensure uniform coverage.
Scotts EasyGreen® Rotary Spreader	X	Make each pass about 30 inches from the previous one.
Broadcast/Rotary Spreaders Scotts® Basic™, Standard™, Deluxe™, EdgeGuard®, Lawn Pro™, or SpeedyGreen®	X	Make each pass about five feet from the previous one.

When using other spreaders, calibrate the spreader settings based on the manufacturer's recommendations.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place to avoid excess heat.. For bags after partial use, reseal opened bag by folding top clown and securing.

Pesticide Container Disposal;

If empty: Non-refillable bag. Do not reuse or refill bag.. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency. Never place unused product down any indoor or outdoor drain.

[Note to PM: Optional Re-entry statements]

People and pets can (re-enter) enjoy the yard (lawn) (area) (immediately) after application is made. If watering in, allow area (lawn) (yard) to dry before re-entering. *(People and Pets icon)*

OR

People and pets can enjoy the yard (lawn) (area) once product has been watered in and allowed to dry. *(People and Pets icon)*

OR

People and pets may re-enter yard (lawn) (area) after area (it) is watered and allowed to dry. *(People and Pets icon)*

OR

People (children) and pets can enjoy the lawn (yard) (area) after lawn (yard) (area) (it) is watered and dry. *(People and Pets icon)*

OR

People (children) and pets can enjoy the area (yard) (lawn) once watered and dry. *(People and Pets icon)*

OR
People and pets may enter treated area (yard) (lawn) after granules have been watered and area (yard) (lawn) has dried. *(People and Pets icon)*

(Precautionary & Environmental Statements)

ORTHO AND YOUR ENVIRONMENT

Your home and yard are places for family and pets to enjoy. That's why Ortho products are designed with care to provide effective solutions to insect problems inside and outside your home. For best results and a healthy environment, please follow instructions for appropriate usage, storage and disposal.

U.S. Patent No. 4,238,505

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

THE ORTHO GUARANTEE

If you are not satisfied with the results after using this product, return the original proof-of-purchase and original receipt to your ORTHO Dealer or send to: The ORTHO Group, Consumer Affairs Dept., P.O. Box 1749, Columbus, OH 43216.

(phone icon) Questions, Comments or Medical Information
call 1-800-225-2883 www.ortho.com

Appendix

[Note to PM: Optional Front Panel Claims]

Barrier

- ✓ (Lawn) Insect Protection

Coverage:

- ✓ A 10lb bag treats 3,484 sq ft of lawn
- ✓ A 14.35lb bag treats 5,000 sq ft of lawn

Speed claims:

- ✓ Kills in 24 hours
- ✓ Kills overnight
- ✓ Fast Acting (formula)
- ✓ Fast Acting against insects, spiders and ticks
- ✓ *Fast-Acting and Long-Lasting!!*
- ✓ *Starts killing (to kill) lawn bugs/insects immediately*
- ✓ *Starts working (to work) immediately*
- ✓ *Starts killing immediately!*
- ✓ *Kills on Contact (above and below the ground/surface)*
- ✓ *Kills in hours*

Residual claims:

- ✓ Long lasting! [up to] 3 month control [Guaranteed]
- ✓ Kills insects on contact [keeps on killing] for [up to] 3 months
- ✓ Keeps [on] killing for [up to] 3 months
- ✓ One (1) Application kills/controls insects/bugs/pests for the entire season*
- ✓ One (1) application provides season long control*
- ✓ Season Long Control*
- ✓ *Apply once and you're done (for the season)*

- ✓ Works two ways: (1) Kills (and) (2) Prevents
- ✓ Works two (2) ways:
 - Kills- starts working immediately [illustration]
 - Prevents- (one application) provides season long control* [illustration]

Sub-surface claims:

- ✓ Controls/Kills [above and/& below the [lawn] surface
- ✓ Kills insects in/and under the lawn

Insects Killed Claims:

- ✓ Kills over 50 insects
- ✓ Kills lawn-damaging insects
- ✓ Broad spectrum
- ✓ Multi-Purpose Insect killer
- ✓ Controls chinch bugs, sod webworms (lawn moth), fleas, ticks (including ticks that may transmit Lyme Disease), and other lawn pests listed on the back

Usage site claims:

- ✓ Use on residential lawns, home perimeters, ornamental & flower gardens
- ✓ Use around outside of homes, porches and/& patios
- ✓ For use on lawns, ornamentals, home perimeter/foundation and flower beds/gardens

Other Claims:

- ✓ Easy-To-Use
- ✓ Odor-free
- ✓ Easy to apply
- ✓ Not harmful to lawn grasses
- ✓ Guaranteed kill or your money back
- ✓ Specially formulated for lawns