

279-3343

03-04-2010

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

MAR - 4 2010

Ms. Nancy Hilton
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Subject: Label Notification(s) for Pesticide Registration Notice 98-10

Dear Ms. Hilton:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice PRN 98-10 dated February 8, 2010 for the following product(s):

F6131 G Insecticide (EPA Reg. No. 279-3343)

The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that the label change(s) requested falls within the scope of PRN-98-10. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact Regina Foushee'-Smith at 703-605-0780.

Sincerely,

A handwritten signature in black ink, appearing to read "Richard J. Gebken".

Richard J. Gebken
Product Manager
Insecticide Branch
Registration Division (7505P)

12



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 279-3343	2. EPA Product Manager Richard Gebken	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) F6132 G Insecticide	PM# 10	
5. Name and Address of Applicant (Include ZIP Code) FMC Corporation 1735 Market Street Phila., PA 19103 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: NOTIFICATION EPA Reg. No. _____ MAR 4 2010 Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification - Alternate Brand Name

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nancy Hilton	Title Product Registrations Manager	Telephone No. (Include Area Code) 215-299-6753
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registrations Manager	
4. Typed Name Nancy Hilton	5. Date February 8, 2010	

FMC Agricultural Products

3
7

FMC Corporation
1735 Market Street
Philadelphia, PA 19103

215.299.6000 Phone
215.299.6468 Fax

www.fmc.com

February 8, 2010

FEDERAL EXPRESS

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
One Potomac Yard - Room S-4900
2777 South Crystal Drive
Arlington, VA 22202-4501

Ms. Linda Arrington
Ombudsperson
Registration Division (7504P)

Dear Ms. Arrington:

Subject: Chemical Number/Name: 129064/zeta-cypermethrin
Chemical Number/Name: 128825/bifenthrin

F6132 G Insecticide (EPA Reg. No. 279-3343)

Alternate Brand Name

FMC is submitting by notification an alternate brand name of "Talstar XTRA Granular Insecticide" for the subject product.

Enclosed please find the following documents:

1. EPA form – 8570-1 for the subject product
2. One copy of the label for the alternate brand name product.

We trust that the enclosed information is adequate for processing this request. If you have any questions pertaining to this application, please call.

Sincerely,



Nancy J. Hilton
Product Registrations Manager
(215) 299-6753
nancy.hilton@fmc.com

FMC

NOTIFICATION

MAR 4 2010

4
7

Talstar XTRA Granular Insecticide

For use to control pests, as listed, outdoors, including areas surrounding residential, institutional, public, commercial and industrial buildings.

For use on lawns and landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

EPA Reg. No. 279-3343

EPA Est. 279-XXX

Active Ingredients:	By Wt.
Zeta-Cypermethrin*	0.05%
Bifenthrin**	0.20%
Other Ingredients:	99.75%
	100.0%

* *Cis/trans* isomer ratio: Max 75% (\pm) *cis*, and Min. 25% (\pm) *trans*
 ** *Cis* isomers 97% minimum; *trans* isomers 3% maximum.
 U.S. Patent No. 4,238,505 & other US patents pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

See other panels for additional precautionary information.



FMC Corporation
 1735 Market Street
 Agricultural Products Group
 Philadelphia PA 19103

Net Contents:

February 8, 2010 (Alt.Brand Name)

GROUP 3 INSECTICIDES

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.	
NOTE TO PHYSICIAN	
This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.	
For information regarding the use of this product call 1-(800)-321-1FMC (1362).	

PRECAUTIONARY STATEMENTS

Hazards to Human (and Domestic Animals)

CAUTION
 Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

STORAGE AND DISPOSAL

Prohibitions:
 Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage:
 Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. Carefully open containers. After partial use, close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spill. Call FMC: (800) 331-3148.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Product Disposal:
 Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Paper and plastic bags: Non-refillable. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration or by burning. Do not burn unless allowed by state or local authorities, if burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale of other commercial use, or for commercial seed production, or for research purposes.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Also for use as a perimeter or spot treatment to control pests outdoors, including areas surrounding residential, institutional, public, commercial and industrial buildings.

Do not apply more than 0.25 lb. active ingredient (ai.) per acre (2.3 lbs granules per 1000 sq. ft.) per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day-care facilities).

May be applied at up to 0.50 lb. ai. per acre (4.6 lbs granules per 1000 sq. ft.) per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.5 lb. ai. per acre (4.6 lbs per 1000 sq. ft.) per year.

Resistance Management Recommendations. Talstar XTRA Granular Insecticide contains Group 3 Insecticides. Insects with acquired resistance to Group 3 Insecticides may eventually dominate the insect population if Group 3 Insecticides are used repeatedly in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Talstar XTRA Granular Insecticide or other Group 3 Insecticides.

Lawn Application: Broadcast with suitable application equipment to ensure uniform coverage over the treatment area.

Perimeter Applications: Apply to ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Applications: Apply to soil areas, ornamental gardens, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas.

Applic. Rate: Pounds Per 1,000 sq. ft.	Band Width	Pounds of Talstar XTRA Granular for Perimeter Applications								
		Linear Distance (Feet) Around Structure								
		100	150	200	250	300	400	500	1,000	
2.3	5 feet	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7	
2.3	10 feet	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7	
3.0	5 feet	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2	
3.0	10 feet	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9	
3.5	5 feet	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5	
3.5	10 feet	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5	
4.6	5 feet	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4	
4.6	10 feet	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4	

Note: The amounts of Talstar XTRA Granular listed in the above table are based on the total area being treated. This area is determined by multiplying the linear distance by the band width and then adding to that value the area of a circle whose radius is equal to the band width (to account for the area treated at the four corners of a rectangular structure).

APPLICATION RATES

The application rates listed in the following table will provide control of the respective pests under typical conditions.

Pest	Usage per Application		
	Lb Talstar XTRA Granular per 1000 ft ²	Lb Talstar XTRA Granular per Acre	Lb Total Active Ingredient per Acre
Annual Bluegrass Weevil (<i>Hyperodes</i>) (adults)	1.15 to 2.3 Lbs/1000 sq ft	50 to 100 Lbs per acre	0.125 to 0.25 Lbs AI/Acre
Armyworms			
Billbugs (adults)			
Black Turfgrass Ataenius (adults)			
Cutworms			
Leafhoppers			
Mealybugs			
Sod Webworms			
Ants (except Carpenter and Pharoah ants)	2.3 to 4.6 Lbs/1000 sq ft	100 to 200 Lbs per Acre	0.25 to 0.50 Lbs AI/Acre
Imported Fire Ants (adults)			
Crane Flies (larvae)			
Fleas (larvae & adults)			
Chinch Bugs			
Millipedes			
Mole Crickets (nymphs & adults)			
Pillbugs			
Sowbugs			
Ticks			

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Talstar XTRA Granular if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

SPECIFIC PEST INSTRUCTIONS

For optimum control of particular pests, follow recommended procedures outlined below:

Armyworms, Cutworms, and Sod Webworms: To ensure optimum control, irrigate the treated area with water immediately after application to activate (release from the granule) the insecticide.

Annual Bluegrass Weevil (*Hyperodes*) adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

Billbug adults: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

Black Turfgrass Ataenius adults: Applications should be made during May and July to control the first and second generation of black turfgrass Ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).

Chinch Bugs: Chinch bugs infest the base of grass plants and are often found in the thatch layer. Irrigate the treated area with up to 0.25 inches of water immediately after application to activate (release from the granule) the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher applications rates may be required to control populations that contain both nymphs and adults during the summer.

Crane Flies: Treatments should be made to control early to mid-season larvae (approximately August - February) as they feed on plant crowns. Treatments made to late-season larvae (approximately March - April) may only provide suppression. Consult your local extensions agent for specific recommendations for your area.

Fleas (adults and larvae): Control is provided for up to seven days. Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release from the granule) the insecticide.

Imported Fire ants: Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound treatments that will eliminate existing colonies. If the soil is not moist, then it is important to irrigate before or soon after application. Broadcast treatments should apply no more than 0.50 lbs ai per acre (4.6 lbs/1000 square feet). Mounds should be treated by applying 1/2 cup of Talstar XTRA Granular per mound and then drenching the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. Treat three feet out around the mound. For best results, apply in cool weather (65 - 80° F) or in early morning or late evening hours. This treatment will kill the queen within 24 hours of application. Application of at least 125 lbs/acre (2.8 lbs granules per 1000 sq. ft.) of Talstar XTRA Granular Insecticide will provide four months residual activity for control of foraging Imported Fire Ants and newly mated fly-in Queens.

Mole Cricket adults: Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

Mole Cricket nymphs: Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not

moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever): Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Do not allow public use of treated areas during application.

Deer ticks (*Ixodes* sp.) have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in mid to late-spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

SPOT TREATMENT APPLICATIONS

Mound Ant Treatment, including Fire Ants: Apply 1/2 cup of Talstar XTRA Granular Insecticide per mound and then drench the mound with 1 to 2 gallons of water. The mound should be drenched with sufficient force to break its apex and allow the insecticide to flow into the ant tunnels. Treat three feet out around the mound. For best results, apply in cool weather (65 to 80°F) or in early morning or late evening hours. Treat new mounds as they appear. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS

(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

Broadcast and Spot Applications: For the control of actively foraging ants (including Imported Fire Ants) broadcast uniformly 2.3 to 4.6 lbs Talstar XTRA Granular Insecticide per 1000 sq. ft.

Pad Mounted Electrical Sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 1.0 to 1.5 Tablespoons of Talstar XTRA Granular Insecticide per 4 sq. ft. of mounting pad. Where existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application. **Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides.**

Note: Electrical transformers appear to be more attractive to ants during the cooler months. For best results treat in the Fall or early Spring.

2
7

SPREADER EQUIPMENT SETTINGS

Broadcast application should be performed with a calibrated commercial broadcast spreader to evenly distribute the granules. Approximate settings for several models are provided in the table below.

Insert spreader setting table here (optional)

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

Conditions of Sale and Limitation of Warranty and Liability:

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and, to the extent permitted by applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent permitted by applicable law, buyer assumes the risk of any such use.

To the extent permitted by applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC – Trademark of FMC Corporation

© 2010 FMC Corporation All rights reserved.