

279-3352

09/03/2008

4/11



**U.S. ENVIRONMENTAL PROTECTION  
AGENCY**

**Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460**

EPA Reg. Number:

279-3352

Date of Issuance:

SEP 3 2008

SEP 3 2008

Term of Issuance:

Conditional – Expiration Date  
of June 1, 2010

Name of Pesticide Product:

F7120 Turf and IVM Herbicide

## NOTICE OF PESTICIDE:

- Registration  
 Reregistration  
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

FMC Corporation  
1735 Market Street  
Philadelphia, PA 19103

**Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.**

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

The Conditional Registration of this product has the same outstanding data requirements as product with EPA Reg. No. 279-3334.

2. Make the following label changes before shipping product:

Add the batch code to product label.

Remove the EPA Registration No. 279-3334 and replace with EPA Registration No. 279-3352 on the product label.

Under "PRECAUTIONARY STATEMENTS", add "wear protective eyewear" after "Applicators and other handlers must wear:".

If the only intended use sites are non-crop sites as listed under "GENERAL INFORMATION", the "Agricultural Use Requirements" box can be removed. Also remove "...but not limited to:" before listed sites in the first paragraph.

In the table "F7120 Turf and IVM Herbicide USE PER ACRE PER 12 MONTH PERIOD\*", add "Non Crop" before "Dormant Turf\*\*\*\*", "Other" before "Non Crop", and "Non Crop" before "Annual Weed Burndown". Also add "Not for use on turf grown for commercial purposes." after table.

Under "GENERAL INFORMATION", second paragraph, the last sentence does not appear to be correct – "Perennials weeds arising from unattached underground rhizomes or rootstocks or plants receiving foliar application of F7120 Turf and IVM Herbicide will not be affected by the spray and will continue to grow." If the present wording is not the intended statement, correct sentence. Alternatively, remove the sentence.

Under "APPLICATION EQUIPMENT AND TECHNIQUES", change "...combine the recommended rates of ...." to "...combine the specified rates of ....", and add "Apply this product by ground application only." If the intent is to allow aerial application, then add the "Aerial Application" and "SPRAY DRIFT REDUCTION ADVISORY (21)" paragraphs on the label of the product with EPA Reg. No. 279-3334. Under the sub-paragraph "Application Equipment", the last sentence of the second paragraph starting with "For flat fan and cone nozzles and with hand directed mist blowers, etc." appears to contradict the requirement in the first sentence that says "Use Coarse (Large droplets) Sprays only." Remove the last sentence related to the use of mist blowers.

The table for "F7120 Turf and IVM Herbicide USE PER ACRE PER 12 MONTH PERIOD\*", contains "(TVC, IVM)" after the use rates for "Non Crop". Either define "TVC" and "IVM" as intended here, or remove.

Submit one copy of the revised final printed label for the record before shipping this product.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Please contact Phil Errico at 703-305-6663/ [errico.philip@epa.gov](mailto:errico.philip@epa.gov) for assistance in this matter.

Signature of Approving Official:

James A. Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)



Date:

SEP 3 2008

3/11

# F7120 Turf and IVM Herbicide

For Use on Non-crop Sites and Dormant  
Turf Applications

<b>EPA Reg. No. 279-3334</b>	<b>EPA Est.</b>
<b>Active Ingredient:</b>	<b>By Wt.</b>
Sulfentrazone* .....	5.70%
Glyphosate IPA** .....	41.48%
<b>Other Ingredients:</b> .....	<u>52.82%</u>
<b>Total:</b>	100.0%

\*F7120 Turf and IVM Herbicide contains 0.56 pounds per US gallon of the active ingredient Sulfentrazone.

\*\* F7120 Turf and IVM Herbicide contains 4.04 pounds per US gallon of the active ingredient Glyphosate, in the form of isopropylamine salt, (3.00 pounds per gallon of glyphosate acid).

Contains Petroleum Distillates

U.S. Patent Pending

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
SEP 3 2008**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

279-3352



**FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia, PA 19103**

4/11

<b>FIRST AID</b>	
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
<b>For Information Regarding the Use of this Product Call 1-800-321-1FMC(1362)</b>	

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**Caution**

Causes moderate eye irritation. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves made of waterproof material such as polyethylene or polyvinyl chloride, and shoes plus socks.**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:**

**Users should:**

Wash hands before using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This pesticide is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

**Groundwater Advisory**

5/11

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Do not use on coarse soils classified as sand which have less than 1% organic matter.**

**Surface Water Advisory**

This product can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

**Physical/Chemical Hazards**

Combustible. Do not use or store near heat or open flame.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, chemical resistant gloves, and shoes plus socks.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

6/11

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

### Pesticide storage

Store product in original container only, away from other pesticides, fertilizer, food or feed.

Store in a cool dry place and avoid excess heat.

Do not store below 32°F degrees.

### In Case of Spill

Avoid contact. Isolate areas and keep out animals and unprotected persons.

### To Confine Spills.

Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a large holding container. Identify contents per required hazardous waste labeling regulations.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### Container Disposal

**Metal or Plastic Containers** - Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: **(For containers greater than 5 gallons)** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **(For containers 5 gallons or less)** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

**Returnable/Refillable Containers** - Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. ¶ To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## GENERAL INFORMATION

F7120 Turf and IVM Herbicide is a liquid suspension formulation that mixes readily with water for control of many herbaceous weeds when applied as a foliar spray in non-crop situations or as a dormant turf application. F7120 Turf and IVM Herbicide may be used to control the listed weeds in terrestrial noncrop sites (such as those listed below). F7120 Turf and IVM Herbicide may be used to trim-and-edge around objects and landscape features in non-crop sites or as a broadcast treatment to remove vegetation from areas including but not limited to:

- Airports Petroleum Tank Farms
- Pipeline, Power, Telephone & Utility Rights-of-Way
- Golf Courses
- Driveways
- Highways & Roadsides Pumping Installations
- Railroads
- Schools
- Storage Areas
- Electrical substations
- Fencerows
- Edge of landscape beds
- Industrial Plant Sites
- Lumberyards
- Parking Areas
- Pavement cracks

Parks  
Residential, commercial & public landscapes  
Hardscapes (like brick walks, patios)  
Stone -covered yards

F7120 Turf and IVM Herbicide enters the root system by moving through the plant from the point of foliar contact. Symptoms, including rapid browning of the foliage, are apparent on most susceptible annual weeds within a one to two days following application. Rapid browning of the leaves is followed by complete necrosis of visible above ground portion of the plant and deterioration of underground plant parts follows. Refer to the "Weeds Controlled" section of this label for specific weeds controlled by F7120 Turf and IVM Herbicide. Treatments made to weeds approaching maturity or at late growth stages will result in the best control of most perennial weeds. Perennials weeds arising from unattached underground rhizomes or rootstocks or plants receiving foliar application of F7120 Turf and IVM Herbicide will not be affected by the spray and will continue to grow.

Use the higher rate of F7120 Turf and IVM Herbicide per acre within the recommended rate range when undesirable vegetation is heavy or dense. Reduced weed control may occur during the following situations:

1. When weed growth are under stress from disease, insects or drought.
- 2.. When weeds are covered with a thick layer of dust or grass clipping.
3. When rainfall or irrigation occurs within 3 hours after application. An additional application may be needed.

F7120 Turf and IVM Herbicide does not provide residual weed control.

**ATTENTION:**  
**TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS, APPLICATORS MUST BE CAUTIOUS TO AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS, OR FRUIT OF CROPS DESIRABLE PLANTS AND TREES. AVOID DRIFT.**

Serious injury or destruction may occur if any amount of F7120 Turf and IVM Herbicide comes in contact with desirable plants. Avoid the following conditions to reduce the likelihood that desirable plants are damaged:

1. Do not allow any amount of F7120 Turf and IVM Herbicide, or the herbicide solution to mist, drip, drift or splash onto desirable vegetation.
2. Avoid application when winds are gusty or in excess of 5 miles per hour.
3. Avoid other conditions including lesser wind velocities, that will allow spray drift to occur.
4. Avoid combinations of pressure and nozzle type that will result in splatter or fine particles, which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

**MIXING AND APPLICATION INSTRUCTIONS:**

Properly maintained and calibrated equipment must be used for effective product application. Avoid spraying desirable plants by carefully directing the spray from hand-gun sprayers or boom equipment. For best results do not mix with water from ponds and ditches that may contain sediment or soil particles.

**Mixing:**

F7120 Turf and IVM Herbicide mixes readily with water. Fill the mixing or spray tank with 1/2 the required amount of water. Add the recommended amount of F7120 Turf and IVM Herbicide and complete filling the tank to the desired level (see "DIRECTIONS FOR USE" and "WEEDS CONTROLLED" sections of this label).

To prevent or minimize foaming of the spray solution that may occur during mixing and application, avoid the use of mechanical agitators, placing the filling hose below the surface of the spray solution, terminate by-pass and return lines at the bottom of the tank and if needed use an approved anti-foam or defoaming agent.

Nozzle screen size or line strainers should be no finer than 50 mesh. Carefully select correct nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**Surfactants:**

When using F7120 Turf and IVM Herbicide for non-crop applications, use a nonionic surfactant labeled for use with herbicides that contains 50 percent or more active ingredient.

8/11

Always read and follow the manufacturer's surfactant label recommendations, cautionary statements and other information for best results.

If colorants or marking dyes are desired, choose those that are approved for use with herbicides. If colorants or marking dyes are used, performance may be reduced, especially at lower rates or dilutions.

Always read and follow the manufacturer's label recommendations, cautionary statements and other information for best results.

## APPLICATION EQUIPMENT AND TECHNIQUES

### Boom Equipment:

When using boom equipment, combine the recommended rates of F7120 Turf and IVM Herbicide in water and apply 30 gallons of dilution per acre as a broadcast spray to control weed species listed in this label unless otherwise specified. See the "WEEDS CONTROLLED" section of this label for specific rates. To ensure best results complete coverage is required. Spray volume should be increased as density of vegetation increases. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

### Application Equipment:

Use Coarse (Large droplets) Sprays Only

Low volume sprayers, such as, knapsack or backpack type sprayers, and high-volume spraying equipment utilizing handguns or other appropriate nozzle configurations may be used to control weeds listed in this label. Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff. This treatment method is most effective in areas where there is a low density of the targeted vegetation and directed sprays are warranted. If a straight stream nozzle is used, start the application at the upper most growing point of the targeted vegetation and spray evenly, contacting a minimum of 50 percent of the foliage, using a back and forth motion until the bottom of the vegetation is reached. For flat fan and cone nozzles and with hand directed mist blowers, mist the application over the foliage of the targeted vegetation.

## F7120 Turf and IVM Herbicide USE PER ACRE PER 12 MONTH PERIOD\*

Use Site	Ounces F7120 Per Acre	Pounds Active F7120** Per Acre	Ounces F7120 Per 1000sqft
Dormant Turf***	8 - 64	0.288 - 2.28	0.18 - 1.47
Non Crop (TVC, IVM)	44.6 - 86.4	1.60 - 3.10	1.02 - 1.98
Annual Weed Burndown	32 - 64	1.14 - 2.28	0.7 - 1.47

\*The total allowed usage per twelve-month period includes all applications made to the field per twelve-month interval. The twelve-month period is considered to begin upon the initial F7120 Turf and IVM Herbicide application.

\*\* Based on total active sulfentrazone and glyphosate IPA

\*\*\* Apply to Dormant Turf for control of winter annuals well in advance of spring greenup. Control of winter annual weeds is best obtained when weeds are small or newly emerged.

Do not Reseed or sod area receiving F7120 application for 60 - 90 Days after application.

## WEEDS CONTROLLED

BROADLEAVES	
Common Name	Scientific Name
Amaranth, livid	<i>Amaranthus lividus</i>
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus powellii</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Amaranth, spleen	<i>Amaranthus dubius</i>
Ammania, purple	<i>Ammania coccinea</i>
Anoda, spurred	<i>Anoda cristata</i>
Bassia, fivehook	<i>Bassia hyssopifolia</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Bittercress	<i>Cardamine hirsuta</i>
Burcucumber	<i>Sicyos anquilatus</i>
Buttercup	<i>Ranunculus spp.</i>
Carolina geranium	<i>Geranium carolinianum</i>
Carpetweed	<i>Mullugo verticillata</i>
Catchweed Bedstraw	<i>Galium aparine</i>
Chervil	<i>Anthriscus cerefolium</i>



9/11

Chickweed, Common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium Strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryeafolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Coreopsis, plains	<i>Coreopsis tinctoria</i>
Corn speedwell	<i>Veronica arvensis</i>
Corn, volunteer	<i>Zea maize</i>
Cotton, volunteer	<i>Gossypium hirsutum</i>
Croton, tropic	<i>Croton glandulosus</i>
Cutleaf evening primrose	<i>Oenothera laciniata</i>
Devilsclaw	<i>Proboscidea louisiana</i>
Dwarf dandelion	<i>Krigia dandelion</i>
Eclipta	<i>Eclipta prostrata</i>
Falseflax, smallseed	<i>Camelina microcarpa</i>
Fiddleneck	<i>Amsinckia menziesii</i>
Field pennycress	<i>Thlaspi arvense</i>
Fleabane hairy	<i>Conyza bonariensis</i>
Fleabane, annual	<i>Erigeron annuus</i>
Fleabane, rough	<i>Erigeron strigosus</i>
Flixweed	<i>Descurainia sophia</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Golden crownbeard	<i>Verbesina encelioides</i>
Groundcherry, clammy (seedling)	<i>Physalis heterophylla</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Groundsel, common	<i>Senecio vulgaris</i>
Hemp sesbania	<i>Sesbania exaltata</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed	<i>Conyza canadensis</i>
Jimsonweed	<i>Datura stramonium</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
London rocket	<i>Sisymbrium irio</i>
Mallow, common	<i>Malva neglecta</i>
Mayweed	<i>Anthemis cotula</i>
Morningglory, Entireleaf	<i>Ipomea hederacea integrusc</i>
Morningglory, ivyleaf	<i>Ipomea hederacea hederacea</i>
Morningglory, Palmleaf	<i>Ipomea Wrightii</i>
Morningglory, pitted	<i>Ipomea lacunosa</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, scarlet	<i>Ipomea hederifolia</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Mustard, black	<i>Brassica nigra</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, tansy	<i>Descurainia pinnata</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Mustard, wild	<i>Brassica kaber</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Prickly lettuce	<i>Lactuca serriola</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Red rice	<i>Oryza punctata</i>
Redmaids	<i>Calandrinia caulescens</i>
Redstem Filaree	<i>Erodium cicutarium</i>
Sheperdspurse	<i>Capsella bursa pastoris</i>
Sicklepod	<i>Senna obtusifolia</i>
Sida, prickly (Teaweed)	<i>Sida spinosa</i>
Smartweed, PA	<i>Polygonum pensylvanicum</i>
Sowthistle, annual	<i>Sonchus asper</i>
Spanishneedles	<i>Bidens bipinnata</i>
Speedwell, purslane	<i>Veronica peregrina</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Spurry, umbrella	<i>Holosteum umbellatum</i>
Sunflower	<i>Helianthus annuus</i>
Swinecress	<i>Coronopus didymus</i>

10/11

Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Virginia pepperweed	<i>Lepidium virginicum</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatos</i>
Yellow rocket	<i>Barbarea vulgaris</i>
<b>GRASSES</b>	
Annual Bluegrass	<i>Poa Annua</i>
Barley	<i>Hordeum vulgare</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Cheat	<i>Bromus secalinus</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Crowfootgrass	<i>Dactyloctenium aegypticum</i>
Cupgrass, woolly	<i>Eriochloa villosa</i>
Dallisgrass	<i>Paspalum dilatatum</i>
Fall panicum	<i>Panicum dichotomiflorum</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, bristly	<i>Setaria verticillata</i>
Foxtail, Carolina	<i>Alopecurus carolinianus</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria lutescens</i>
Goatgrass, jointed	<i>Aegilops cylindrica</i>
Goosegrass	<i>Eleusine indica</i>
Grain sorghum (milo)	<i>Sorghum bicolor</i>
Itchgrass	<i>Rottboellia cochinchinensis</i>
Johnsongrass, seedling	<i>Sorghum halapense</i>
Junglerice	<i>Echinochloa colona</i>
Oats	<i>Avena sativa</i>
Rye	<i>Secale cereale</i>
Ryegrass	<i>Lolium multiflorum</i>
Sandbur, field	<i>Cenchrus spinifex</i>
Sandbur, longspine	<i>Cenchrus longispinus</i>
Shattercane	<i>Sorghum bicolor</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Sprangletop	<i>Leptochloa fusca</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Texas panicum	<i>Panicum texanum</i>
Wheat, volunteer	<i>Triticum aestivum</i>
Wild Buckwheat	<i>Polygonum convolvulus</i>
Wild oats	<i>Avena fatua</i>
Wild proso millet	<i>Panicum miliaceum</i>
Witchgrass	<i>Panicum capillare</i>
<b>SEDGES</b>	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

### Application to reseeded, overseeded or sprigged areas:

Reseeding, overseeding or sprigging of treated areas within one (1) month after application of this product could inhibit the establishment of desirable turfgrasses. Overseeding of bermudagrass with perennial ryegrass at two (2) to four (4) weeks after an application can be done if slight injury to perennial ryegrass can be tolerated.

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**Notice:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and, to the extent permitted by applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

11/11

Seller warrants that this product conforms to the clinical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and, to the extent permitted by applicable law, buyer assumes the risk of any such use.

To the extent permitted by applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.