

279-3411

10/03/2011

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Ms. Jill Holihan  
FMC Corporation  
1735 Market Street  
Philadelphia, PA 19103

OCT - 3 2011

Subject: Label Amendment  
Product Name: F9009-1 WDG Herbicide  
EPA Reg. No. 279-3411  
Application dated: April 25, 2011  
Revised label submitted: September 23, 2011

Dear Ms. Holihan:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. Products shipped after 18 months from the date on this notice or the next printing of the label, whichever occurs first, must bear the new revised label. Amended labeling will supersede all previously accepted ones. Submit one (1) copy of final printed labeling before you release the product for shipment.

If you have questions or concerns regarding this letter, please contact Beth Benbow at (703) 347-8072 or email at [benbow.bethany@epa.gov](mailto:benbow.bethany@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn V. Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

# F9009-1 WDG Herbicide


For Selective Weed Control in Turf Including Residential, Commercial and Institutional Lawns, Athletic Fields, Commercial Sod Farms, and Golf Courses (Fairways and Roughs)

EPA Reg. No. 279-3411

EPA Est. XXX

<b>Active Ingredients:</b>	<b>By Wt.</b>
Sulfentrazone*	60.00%
Metsulfuron-methyl*	6.00%
Other Ingredients:	34.00%
<b>Total:</b>	<b>100.0%</b>

\*F9009-1 WDG Herbicide contains 0.66 pounds of active ingredient per one pound of product.



## KEEP OUT OF REACH OF CHILDREN CAUTION

See other panels for additional precautionary information.

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
For Information Regarding the Use of this Product Call 1-800-321-1FMC(1362)	

### Net Contents:

**FMC**  
**FMC Corporation**  
**Agricultural Products Group**  
**1735 Market Street**  
**Philadelphia, PA 19103**

## PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

### CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

### Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations:

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## Environmental Hazards

This pesticide is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

Groundwater Advisory: The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface Water Advisory: F9009-1 WDG Herbicide can contaminate surface water through spray drift. Under some conditions, F9009-1 WDG Herbicide may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over long-sleeved shirt and long pants.
- chemical-resistant gloves made of any waterproof material.
- shoes plus socks.

## Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### Pesticide Storage

Store product in original container only, away from other pesticides, fertilizer, food or feed.

Store in a cool, dry place and avoid excess heat.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and spills): (800) 424-9300.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**

If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

**Container Disposal**

**Metal or Plastic Containers:** - Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: **(For containers greater than 5 gallons)** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **(For containers 5 gallons or less)** empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available, or reconditioning, or if approved by state and local authorities, puncture and dispose of in sanitary landfill, or by other procedures. Do not cut or weld or cut metal containers.

**Returnable/Refillable containers:** - Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into the application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Turf Use Instructions**

**Product Information**

F9009-1 WDG Herbicide is a selective post-emergence herbicide which controls annual grasses, broadleaf weeds and sedges in established turf areas including, but not limited to, residential, commercial and institutional lawns, athletic fields, commercial sod farms, golf course fairways and golf course roughs.

F9009-1 WDG Herbicide is formulated as a WDG (Water Dispersible Granule) containing 0.66 lb of active ingredient per pound of product. F9009-1 WDG Herbicide is absorbed by shoots, foliage and roots.

A post emergent application of F9009-1 WDG Herbicide is improved when adequate soil moisture is present at application. Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application. If no rainfall or irrigation occurs within 7 days after application of F9009-1 WDG Herbicide in the amount of 0.5 inches, then irrigation of at least 0.5 inches is recommended.

**Mixing and Application Instructions**

**Product handling instructions**

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well, are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

**SPRAY TANK PREPARATION**

It is important that spray equipment is clean and free of existing pesticide deposits before using this product. Follow the spray tank clean out procedures specified on the label of product previously applied before adding F9009-1 WDG Herbicide to the tank.

F9009-1 WDG Herbicide is a water dispersible granule intended for dilution with water. Mix F9009-1 WDG Herbicide thoroughly and continue agitation during application. If F9009-1 WDG Herbicide is left standing for extended period of time in spray tank, re-agitate to assure uniform suspension of product in spray mixture.

**MIXING WITH WATER**

For best results, fill spray tank with one fourth of the volume of clean water needed for the area to be treated. Start the agitation system and add F9009-1 WDG Herbicide to the tank. Make sure F9009-1 WDG Herbicide is thoroughly mixed before application or before adding another product to the spray tank.

**USE OF SURFACTANTS**

**Temporary discoloration of some turf types may result from use of surfactants or adjuvants with F9009-1 WDG Herbicide. High temperatures and high relative humidity may increase the risk of temporary discoloration. Use of surfactants is not recommended.**

**TANK MIXTURES COMPATIBILITY**

F9009-1 WDG Herbicide has been found to be compatible with most herbicides, fungicides, insecticides and growth regulators commonly used in turf and ornamental plant management. However, when preparing a new tank mix conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. **Shake the mixture vigorously and allow it to stand for five to ten minutes.** Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied. Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the recommended amounts of ingredients using the following order: dry granules first, then liquid suspensions

(flowables) second. As the agitation continues and the tank is filled with water add EC products third followed by the addition of water soluble products.

Read and observe mixing instructions of all tank mix partners. Also read each product label for Directions for Use, Precautionary Statements and Restrictions and Limitations. The most restrictive labeling applies in all tank mixtures. No label dosage rate should be exceeded. Tank mixture recommendations are for use only in states where the companion products and application site are registered. In addition, certain states or geographical regions may have established dosage rate limitations. Consult your state Pesticide Control Agency for additional information regarding the maximum use rates.

Use F9009-1 WDG Herbicide spray mixture immediately after mixing. Do not store the mixture.

**Ground Equipment**

Power sprayers: Uniform and accurate spray coverage requires proper calibration, agitation and operation of spray equipment. The use of marker dyes or foams can improve application accuracy. Boom sprayers equipped with appropriate flat fan nozzles, tips and screens are ideal for broadcast applications. Power sprayers fitted with spray wand/gun may also be used for broadcast application after careful calibration by the applicator. Power sprayers fitted with spray wand/gun are suitable for spot treatments.

Hand operated sprayers: Backpack and compression sprayers are appropriate for small turfgrass areas and spot treatments. Wands fitted with a flat fan nozzle tip should be held stationary at the proper height during application. A side to side or swinging arm motion can result in uneven coverage.

Apply this product in a sufficient volume of carrier solution to provide a uniform spray distribution. Spray volumes of 20 – 175 gallons per acre (0.5 to 4.0 gal/1,000 ft<sup>2</sup>) with spray pressures adjusted to 20 – 40 psi are appropriate. Apply the higher spray volumes for dense weed populations.

**Sprayer Equipment Clean-Out**

**After spraying F9009-1 WDG Herbicide and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure:**

1. Drain sprayer tank, hoses, and spray boom and thoroughly rinse the inside of the sprayer tank with clean water to remove sediment and residues. In addition, thoroughly flush sprayer hoses, boom, and nozzles with clean water.
2. Fill the tank 1/2 full with clean water, and add appropriate detergent or ammonia (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom, and nozzles.
3. Drain the sprayer system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately.
4. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State and local regulations and guidelines.

Do not drain or flush equipment on or near desirable trees or plants. Do not contaminate any body of water including irrigation water that may be used on other plants.

**Weed Control in Turfgrass**

Use Precautions for Turf Use

Turfgrass Safety

This product may be used on seeded, sodded or sprigged turfgrass that are well established. First application of this product can be made following the second mowing providing the turfgrass has developed into a uniform stand with a good root system. Turfgrass injury could result from application of this product on turfgrass that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

**Table 1. Tolerant grasses.**

Grass Type*	Single Application Use Rate		
	Ounces per 1000 ft <sup>2</sup>	Ounces per acre	lbs ai/A
<b>Cool Season Grasses</b>			
Bluegrass, Kentucky ( <i>Poa pratensis</i> ) Fine Fescue, ( <i>Festuca rubra</i> ) Fescue, tall ( <i>Festuca arundinacea</i> )	0.075-0.23	3.25-10	0.134-0.413
<b>Warm Season Grasses</b>			
Bermudagrass ( <i>Cynodon dactylon</i> ) & hybrids Centipedegrass ( <i>Eremochloa ophiuroides</i> ) St. Augustine grass ( <i>Stenotaphrum secundatum</i> ) Zoysiagrass ( <i>Zoysia japonica</i> )	0.075-0.23	3.25-10	0.134-0.413

\*F9009-1 WDG Herbicide has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply F9009-1 WDG Herbicide to a small area prior to treatment of larger areas.

If turf discoloration occurs after an application of F9009-1 WDG Herbicide turfgrass will generally recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply F9009-1 WDG Herbicide on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.

**Application to reseeded, overseeded or sprigged areas:**

Reseeding, overseeding or sprigging of treated areas within one (1) month after application of this product could inhibit the establishment of desirable turfgrasses. Overseeding of bermudagrass with perennial ryegrass at six (6) to eight (8) weeks after an application can be done if slight injury to perennial ryegrass can be tolerated. A spring application of F9009-1 WDG Herbicide may aid in overseeded ryegrass transition in Bermudagrass, transition efficacy may be species dependent. Best results are obtained for

reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

**Sod Production:**

This product may be applied to established sod. Allow sod to establish a good root system, a uniform stand and to fill in the exposed edges. It is recommended that sod be established for up to three (3) months before an application of F9009-1 WDG Herbicide. Do not apply this product within three (3) months of harvest.

**Restrictions and Other Use Precautions:**

- Do not apply to golf course putting greens, collars or tees.
- Do not use on turfgrass other than those listed on this label.
- Do not apply to turfgrass under stress.
- Do not apply with surfactants unless previous experience has demonstrated combinations with surfactant to be physically compatible and non-injurious to the grass type in question.
- Do not treat pastures, rangeland, or other areas grazed or harvested for livestock forage or hay.
- Do not apply directly to landscape ornamentals or ornamental beds.
- Do not allow spray drift to contact landscape ornamentals, shrubs and trees.
- Temporary turfgrass discoloration has been observed when trinexapac-ethyl products have been either tank-mixed or applied within 7 days of a F9009-1 WDG Herbicide application. It is recommended that trinexapac-ethyl applications be made 7 days prior to, or after F9009-1 WDG Herbicide application to reduce risk of turfgrass discoloration.
- Do not use clippings as mulch or compost around flowers, ornamentals, trees or in vegetable gardens.
- Do not plant in treated areas with shrubs and trees for one year after application. Beddings should be planted in treated areas for two years following application.

**POSTEMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS**

When used as directed F9009-1 WDG Herbicide will control or suppress the weeds listed in Table 2 when applied alone shortly after weeds have emerged. Do not exceed the application rate specified for the turfgrass species in Table 1.

**Table 2. Weeds Controlled or Suppressed by F9009-1 WDG Herbicide**

Common Name	Scientific Name	Control	Suppress
Bittercress	( <i>Cardamine spp.</i> )	X	
Barnyardgrass	( <i>Echinochloa crusgalli</i> )		X
Black medic	( <i>Medicago lupulina</i> )	X	
Bluegrass, annual	( <i>Poa annua</i> )		X
Buttercups	( <i>Ranunculus spp.</i> )	X	
Buttonweed, Virginia	( <i>Diodia Virginiana L</i> )		X
Carolina geranium	( <i>Geranium carolinianum</i> )	X	
Carpetweed	( <i>Mollugo verticillata</i> )	X	
Chickweed, common	( <i>Stellaria media</i> )	X	
Chickweed, mouseear	( <i>Cerastium vulgatum</i> )		X
Cinquefoil	( <i>Potentilla spp.</i> )		X
Clover	( <i>Trifolium spp.</i> )	X	
Crabgrass (Large and Smooth)	( <i>Digitaria spp.</i> )		X
Creeping Beggarweed	<i>Desmodium canum</i>	X	
Crown Vetch	( <i>Vicia sativa l.</i> )		
Cudweed	( <i>Gnaphalium spp.</i> )		X
Dandelion	( <i>Taraxacum officinale</i> )	X	
Dock, Curly	( <i>Rumex crispus</i> )	X	
Dogfennel	( <i>Eupatorium capillifolium(Lam.) Small</i> )	X	
Dollarweed	( <i>Hydrocotyle spp.</i> )	X	
Doveweed	( <i>Murdannia nudiflora(L.) Brenan</i> )		X
Eclipta	( <i>Eclipta prostrata (L.)</i> )		X
Evening primrose	( <i>Oenothera biennis</i> )		X
Fiddleneck	( <i>Amsinckia spp.</i> )		X
Filaree	( <i>Erodium spp.</i> )		X
Foxtail spp.	( <i>Setaria spp</i> )	X	
Groundsel	( <i>Senecio smallii Britt.</i> )	X	
Goldenrod	( <i>Solidago spp.</i> )	X	
Goosegrass	( <i>Eleusine indica</i> )	X	
Ground ivy	( <i>Glechema hederacea</i> )	X	
Henbit	( <i>Lamium amplexicaule</i> )	X	
Japanese Stiltgrass	( <i>Microstegium vimineum</i> )	X	
Knotweed, prostrate	( <i>Polygonum aviculare</i> )	X	
Kochia	( <i>Kochia scoparia</i> )	X	
Kyllinga, green	( <i>Kyllinga brevifolia Rottb.</i> )	X	
Kyllinga, False green	( <i>Kyllinga gracillima Miq.</i> )	X	
Lambsquarters, common	( <i>Chenopodium album</i> )	X	
Lawn burweed	( <i>Soliva pterosperma</i> )		X
Lespedeza, common	( <i>Lespedeza striata</i> )	X	
London Rocket	( <i>Sisymbrium irio</i> )	X	
Mallow, common	( <i>Malva neglecta</i> )	X	
Miner Lettuce	( <i>Claytonia perfoliata</i> )		
Morningglory	( <i>Ipomea spp.</i> )	X	
Mustard, Tansy	( <i>Descurainia pinnata brachycarpa</i> )	X	
Mustard, Tumble	( <i>Sisymbrium altissimum L.,</i> )	X	
Nutsedge, Purple	( <i>Cyperus rotundus</i> )		X
Nutsedge, Yellow	( <i>Cyperus esculentus</i> )	X	
Parsley piert	( <i>Alchemilla arvensis</i> )	X	

Pigweed, Redroot	( <i>Amaranthus retroflexus</i> )	X	
Pigweed, Smooth	( <i>Amaranthus hybridus</i> )		
Pigweed, Tumble	( <i>Amaranthus albus</i> )	X	
Pineapple weed	( <i>Matricaria matricanoides</i> )		X
Plantain, broadleaf	( <i>Plantago major</i> )	X	
Plantain, buckhorn	( <i>Plantago lanceolata</i> )	X	
Prickly Lettuce	( <i>Lactuca serriola</i> )		X
Puncture weed	( <i>Tribulus terrestris</i> )		X
Purslane, common	( <i>Portulaca oleracea</i> )		X
Pusley, Florida	( <i>Richardia scabra</i> )	X	
Sedge, Globe	( <i>Cyperus croceus Vahl</i> )	X	
Sedge, annual	( <i>Cyperus compreeus L.</i> )	X	
Shepardspurse	( <i>Capsella bursa pastoris</i> )		X
Smartweed, Pennsylvania	( <i>Polygonum pennsylvanicum</i> )	X	
Sorrel, Red	( <i>Rumex acetosella</i> )		X
Speedwell	( <i>Veronica spp.</i> )	X	
Spurge, (annuals)	( <i>Euphorbia spp.</i> )	X	
Spurge, prostrate	( <i>Euphorbia humistrata</i> )	X	
Spurge, spotted	( <i>Euphorbia maculata</i> )	X	
Star of Bethlehem	( <i>Ornithogalum umbellatum</i> )	X	
Violet, wild	( <i>Viola pratensis</i> )	X	
Wild garlic	( <i>Allium vineale</i> )	X	
Wild onion	( <i>Allium canadense</i> )	X	
Woodsorrel, creeping	( <i>Oxalis corniculata</i> )	X	
Woodsorrel, yellow	( <i>Oxalis stricta</i> )	X	

- Two applications of F9009-1 WDG Herbicide 21 days apart may increase control of weeds over a single application; however, do not exceed 10 oz. product per acre per year

**POSTEMERGENCE CONTROL OF ANNUAL AND PERENNIAL SEDGES**

F9009-1 WDG Herbicide will control or suppress sedges (Table 3). Apply the highest rate consistent with the rate needed for turfgrass safety in Table 1.

Good spray coverage is needed for optimum control of sedges. Temporary discoloration of some turfgrass species may result from use of surfactant. Use of surfactants is not recommended.

**Table 3. Sedge species controlled or suppressed by F9009-1 WDG Herbicide**

Common Name	SCIENTIFIC NAME
Kyllinga, green	( <i>Kyllinga brevifolia</i> )
Kyllinga, false green	( <i>Kyllinga gracillima</i> )
Nutsedge, purple	( <i>Cyperus rotundus</i> )
Nutsedge, yellow	( <i>Cyperus esculentus</i> )
Sedge, annual	( <i>Cyperus compreeus L.</i> )
Sedge, globe	( <i>Cyperus globulosus</i> )
Sedge, cylindric	( <i>Cyperus retrorsus</i> )
Sedge, Surinam	( <i>Cyperus surinamensis</i> )
Sedge, Texas	( <i>Cyperus polystachyos</i> )

- Two applications of F9009-1 WDG Herbicide 21 days apart may increase control of weeds over a single application; however, do not exceed 10 oz. product per acre per year

**Conditions of Sale and Limitation of Warranty and Liability:**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC -- Trademark of FMC Corporation

© FMC Corporation 2010  
9/19/11 Amendment