



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 25, 2023

Catherine Rice
Product Registration Manager
FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

Subject: Notification per PRN 98-10 – Add optional marketing claims to sublabels and other non-notification edits
Product Name: F6875-5 0.4SC RTS Herbicide
EPA Registration Number: 279-3427
Application Date: June 15, 2023
Decision Number: 592892

Dear Catherine Rice:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact me at ondish.mindy@epa.gov or at (202)566-2857.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505T)
Office of Pesticide Programs

NOTE to EPA Product Manager: Text in brackets [] is optional.

NOTIFICATION

279-3427

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

08/25/2023

[Version A – CONCENTRATE]

F6875-5 0.4SC RTS

Herbicide

For Preemergence and early Postemergence Control of Crabgrass and selected annual grasses. Selective Preemergence Control of Annual Broadleaf weeds and Preemergence and Postemergence Control of Sedges in Lawns.

EPA Reg. No. 279-3427

EPA Est. No.

Active Ingredients:	By Wt.
Sulfentrazone:	1.36%
Prodiamine:	2.73%
Other Ingredients:	<u>95.91%</u>
	100.0%

Contains 0.36 pounds of active ingredient (0.12 lb ai sulfentrazone and 0.24 lb ai prodiamine) per gallon

Net Contents: _____

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional precautionary information contained inside label booklet.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

NOTE to EPA Product Manager: Text in brackets [] is optional.

FIRST AID	
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-585-7378 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as natural rubber, selection Category A). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to marine/estuarine invertebrates. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRODUCT INFORMATION

Turf Use Instructions

F6875-5 RTS is a selective preemergence and postemergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in residential lawns.

F6875-5 RTS is a flowable formulation (suspension concentrate) containing 0.4 lbs. of active ingredient per gallon. The mode of action of F6875-5 RTS herbicide involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

APPLICATION INSTRUCTIONS

F6875-5 RTS can be applied using a hand trigger sprayer, pump-up sprayer or hose-end sprayer.

See Table 1 for specific use instructions for either cool or warm-season turfgrass varieties. For preemergence applications, spray recommended rate evenly over entire lawn. For postemergence applications, spray weeds directly until well covered. Do not over-apply as this may cause undesirable turf injury.

For pump-up sprayers, mix recommended amount of tablespoons in Table 1 according to desired turfgrass type in 1 gallon of water per 1000 sq ft covered.

Mix and apply only the amount of product needed to cover desired area. Do not store mixed product.

SPECIFIC USE INSTRUCTIONS IN TURFGRASS

Established Turf

This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

When applied as directed under the conditions described, the following established grasses are tolerant to F6875-5 RTS herbicide at the recommended use rates. Do not exceed 8.25 fl oz/1000 sq ft per calendar year.

NOTE to EPA Product Manager: Text in brackets [] is optional.

Table 1. Tolerant grasses.

Grass Type	Recommended Use Rates Do not exceed maximum use rates per turfgrass species in a single application	
	Fluid ounces per 1000 sq ft	Area Covered (Based on 32 fl oz sprayer jar or retail container)
Cool Season Grasses		
Fescue, fine ¹ (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>), Bluegrass, Kentucky (<i>Poa pratensis</i>) Fescue, tall ² (<i>Festuca arundinacea</i>) Cool-season mixtures	2.75 fl oz Followed by 2.75 fl oz 30 - 60 days later (5.5 Tablespoons followed by 5.5 Tablespoons)	11,600 sq ft
Warm Season Grasses		
Bahiagrass (<i>Paspalum notatum</i>) Buffalograss (<i>Buchloe dactyloides</i>) Carpetgrass (<i>Axonopus affinis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kikuyugrass (<i>Pennisetum clandestinum</i>) Seashore Paspalum (<i>Paspalum vaginatum</i>) Zoysiagrass ³ (<i>Zoysia japonica</i>) Bermudagrass ⁴ (<i>Cynodon dactylon</i>) & hybrids	5.5 fl oz (8 Tablespoons)	5,800 sq ft
St. Augustinegrass	4.0 fl oz Followed by 4.0 fl oz 30 - 60 days later (8 Tablespoons followed by 8 Tablespoons)	8,000 sq ft
<ol style="list-style-type: none"> 1 Use of this product on certain varieties of Chewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury. 2 F6875-5 RTS may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate listed per turf type or 8.25 fl oz/1000 sq ft per calendar year, whichever is lower. 3 F6875-5 RTS application may cause temporary discoloration to exposed leaf surfaces on certain cultivars of zoysiagrass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply F6875-5 RTS on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth. 4 Sequential application of F6875-5 RTS may be needed in bermudagrass lawns for season-long control of yellow nutsedge. If sequential application is needed apply no more than 2.75 fl oz/1000 sq ft. 		

F6875-5 RTS has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties and or cultivars have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply F6875-5 RTS to a small area prior to treatment of larger areas.

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds, Grasses and Sedges

F6875-5 RTS herbicide will control or suppress the weeds listed in [Table 2](#) when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1. Do not apply adjuvants or surfactants with post applications of F6875-5 RTS unless tested on small areas prior to broadcast applications.

Table 2. Weeds (Preemergent and Postemergent) Controlled/Suppressed by Applications of F6875-5 RTS

Common Name	Scientific Name	Preemergent	Postemergent	
			Controlled	Suppressed
Barnyardgrass	(<i>Echinochloa crus-galli</i>)	X		
Bittercress	(<i>Cardamine spp.</i>)			X
Black medic	(<i>Medicago lupulina</i>)			X
Bluegrass annual	(<i>Poa annua</i>)	1		
Buttercups	(<i>Ranunculus spp.</i>)			X
Carolina geranium	(<i>Geranium carolinianum</i>)	X	5	
Carpetweed	(<i>Mollugo verticillata</i>)	X		X
Chickweed, common	(<i>Stellaria media</i>)	2	5	
Chickweed, mouseear	(<i>Cerastium vulgatum</i>)			X
Chickweed, mouseear (from seed)	(<i>Cerastium vulgatum</i>)	X		
Cinquefoil	(<i>Potentilla spp.</i>)			X
Clover	(<i>Trifolium spp.</i>)			X
Crabgrass (Large and Smooth)	(<i>Digitaria spp.</i>)	3	6	
Crowfootgrass	(<i>Dactyloctenium aegyptium.</i>)	X		
Cudweed	(<i>Gnaphalium spp.</i>)			X
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)	X		
Dandelion	(<i>Taraxacum officinale</i>)			X
Dock, Curly	(<i>Rumex crispus</i>)			X
Evening primrose	(<i>Oenothera biennis</i>)			X

NOTE to EPA Product Manager: Text in brackets [] is optional.

Fiddleneck	(<i>Amsinckia spp.</i>)			X
Filaree	(<i>Erodium spp.</i>)			X
Foxtails, Annual	(<i>Setaria spp.</i>)	X		
Goldenrod	(<i>Solidago spp.</i>)		5	
Goosegrass	(<i>Eleusine indica</i>)	4	6	
Ground ivy	(<i>Glechema hederacea</i>)		5	
Hairy Bittercress	<i>Cardamine hirsute</i>	X		
Henbit	(<i>Lamium amplexicaule</i>)	2		X
Itchgrass	(<i>Rottboellia exaltata</i>)	X		
Knotweed, prostrate	(<i>Polygonum aviculare</i>)	X		X
Kochia	(<i>Kochia scoparia</i>)	X	5	
Kyllinga, green	(<i>Kyllinga brevifolia</i>)	X		
Kyllinga, false	(<i>Kyllinga gracillima</i>)	X		
Lambsquarters, common	(<i>Chenopodium album</i>)	X		X
Lawn burweed	(<i>Soliva pterosperma</i>)			X
Lespedeza, common	(<i>Lespedeza striata</i>)			X
Lovegrass	(<i>Eragrostis spp.</i>)	X		
Mallow, common	(<i>Malva neglecta</i>)			X
Nutsedge, Yellow	(<i>Cyperus esculentus</i>)	X	5	
Panicum (Texas,Fall)	(<i>Panicum spp.</i>)	X		
Parsley piert	(<i>Alchemilla arvensis</i>)		5	
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)	X	5	
Pigweed, Tumble	(<i>Amaranthus albus</i>)		5	
Pineapple weed	(<i>Matricaria matricarioides</i>)			X
Plantain, buckhorn	(<i>Plantago lanceolata</i>)		5	
Puncture weed	(<i>Tribulus terrestris</i>)			X
Purslane, common	(<i>Portulaca oleracea</i>)	X		X
Pusley, Florida	(<i>Richardia scabra</i>)	X	5	
Redweed	(<i>Melochia corchorifolia</i>)			X
Rocket, London	(<i>Sisymbrium irio</i>)			X
Sedge, globe	(<i>Cyperus globulosus</i>)	X		
Sedge, cylindrical	(<i>Cyperus retrorsus</i>)	X		
Sedge, Surinam	(<i>Cyperus surinamensis</i>)	X		
Sedge, Texas	(<i>Cyperus polystachyos</i>)	X		
Shephardspurse	(<i>Capsella bursa-pastoris</i>)	2		
Signalgrass, broadleaf	(<i>Brachiaria playphylla</i>)	X		
Smartweed, Pennsylvania	(<i>Polygonum pennsylvanicum</i>)		5	
Sorrel, Red	(<i>Rumex acetosella</i>)			X
Speedwell	(<i>Veronica spp.</i>)	X	5	
Sprangletop	(<i>Leptochola spp.</i>)	X		
Spurge, annuals	(<i>Euphorbia spp.</i>)		5	
Spurge, prostrate	(<i>Euphorbia humistrata</i>)	X	5	
Spurge, spotted	(<i>Euphorbia maculata</i>)		5	
Star of Bethlehem	(<i>Ornithogalum umbellatum</i>)		5	
Velvetleaf	(<i>Abutilon theophrasti</i>)			X
Violet, wild	(<i>Viola pratincola</i>)			X
Wild garlic	(<i>Allium vineale</i>)			X
Wild onion	(<i>Allium canadense</i>)			X
Witchgrass	(<i>Panicum capillare L.</i>)	X		
Woodsorrel, creeping	(<i>Oxalis corniculata</i>)			X
Woodsorrel, yellow	(<i>Oxalis stricta</i>)		5	
Woodsorrel, yellow (from Seed)	(<i>Oxalis stricta</i>)	X		

- 1 Poa annua is a winter annual. Applications of F6875-5 RTS should be made in August or September to established, non overseeded turf before Poa Annuu germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings.
- 2 Apply F6875-5 RTS in late summer, fall or winter before weed germinates for preemergent control.
- 3 **Crabgrass control in cool season grasses from fall application:** Areas where the ground is subject to winter freezes, F6875-5 RTS may be applied in the fall at rates listed in Table 1 (8-24 fluid ounces per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.
- 4 In areas of extended growing season and heavy pressure, preemergent weed control will be enhanced by second application of F6875-5 RTS 60-90 days after initial application
- 5 Weeds are suppressed at lower label rates (<5.5 fl oz/1000 sq ft). For optimum control apply rates of at least 5.5 fl oz per 1000 sq ft in a single application. Do not exceed the application rate specified for the turf species in Table 1.
- 6 F6875-5 RTS herbicide controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.

Application to Reseeded, Overseeded or Sprigged Areas

Reseeding, overseeding or sprigging of treated areas within three to six (3-6) months after application of this product could inhibit the establishment of desirable grasses. [See Table 3 for exact intervals from application].

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

NOTE to EPA Product Manager: Text in brackets [] is optional.

Table 3. Overseeding Intervals after F6875-5 RTS Applications

Ounces Product/1000 sq ft	Total Amount of F6875-5 RTS Applied		Interval (Months Before Overseeding)	
	North	South	Transition	South
2.75	3		3	3
4.0	3		3	3
5.5	4		4	4
8.25	6		5	5

Other Use Precautions

- Do not use on turfgrasses other than those listed on this label.
- Do not graze or feed livestock, forage cut from areas treated with F6875-5 RTS herbicide.
- Do not apply directly to landscape ornamentals or ornamental beds.
- Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.
- Do not apply with adjuvants or surfactants.

STORAGE AND DISPOSAL

Pesticide Storage: Store and transport in upright position. Store only in original container, in a cool dry place inaccessible to children or pets.
Pesticide Disposal: *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain
Container Disposal: *If empty* – Non-refillable container. Do not reuse or refill this container. Offer for recycling, if possible.

WARRANTY

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents) call National Pesticide Information Center at 1-800-858-7378.

{Note to Reviewer: The following are optional advertising and/or marketing phrases that may be present on promotional materials and container labels}

- Tough Weed Killer
- Kills even the toughest weeds like crabgrass, dandelion and chickweed
- Kills Yellow Nutsedge, Crabgrass, Dandelions and other listed weeds
- Kills broadleaf and grassy weeds
- Starts working on contact
- Works through the shoots and roots of the weed
- Rainproof – won't wash away
- Rainfast in hours
- Provides excellent cool weather performance
- Performs in both cool and warm weather
- For use on Warm Season and Cool Season grasses
- Kills the weeds, not the lawn
- Water – Based
- 1 quart treats up to 11,600 sq ft
- This container covers up to 11,600 sq ft
- Won't harm lawn grasses when used as directed
- All-in-one weed control for lawns
- Provides Pre-emergent and Post-emergent weed control
- Kills weeds before they become a problem
- Kills the toughest weeds, down to the root
- Lawn – Friendly formula, when used as directed
- Weed Deadly – Lawn Friendly
- For home lawns only
- For home lawn care
- For a beautiful, weed-free lawn
- Helps keep lawns weed-free
- Economical concentrate
- Controls dandelions before they emerge [from seed]
- Kill & Prevent Weeds [in your lawn]

Non-FIFRA claims
 Bonus size 20%

NOTE to EPA Product Manager: Text in brackets [] is optional.

[Version B – FOR READY-TO-SPRAY PACKAGING]

F6875-5 0.4SC RTS

Herbicide

For Preemergence and early Postemergence Control of Crabgrass and selected annual grasses. Selective Preemergence Control of Annual Broadleaf weeds and Preemergence and Postemergence Control of Sedges in Lawns.

EPA Reg. No. 279-3427

EPA Est. No.

Active Ingredient:	By Wt.
Sulfentrazone:	1.36%
Prodiamine:	2.73%
Other Ingredients:	<u>95.91%</u>
	100.0%

Contains 0.36 pounds of active ingredient (0.12 lb ai sulfentrazone and 0.24 lb ai prodiamine) per gallon

Net Contents: _____

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary information contained inside label booklet.

FMC
FMC Corporation
2727 Walnut Street
Philadelphia, PA 19104

NOTE to EPA Product Manager: Text in brackets [] is optional.

FIRST AID	
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-585-7378 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as natural rubber, selection Category A). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to marine/estuarine invertebrates. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRODUCT INFORMATION

Turf Use Instructions

F6875-5 RTS is a selective preemergence and postemergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in residential lawns.

F6875-5 RTS is a flowable formulation (suspension concentrate) containing 0.4 lbs. of active ingredient per gallon. The mode of action of F6875-5 RTS herbicide involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

APPLICATION INSTRUCTIONS

FOUR STEPS TO APPLY

1. **Measure** the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. **TO BEGIN** spraying: Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the **OFF** position. Turn on water at faucet. Point (Aim) nozzle toward lawn and turn the control valve to the **ON** position. Water will automatically mix with the product. Spray evenly over the measured area.
3. **SPRAY:** Spray evenly over the measured area. Walk at a steady pace. This 32 fl oz container will treat 11,600 sq ft for Cool Season Grasses, 5,800 sq ft for Warm Season Grasses and 8,000 sq ft for St. Augustinegrass.
4. **TO STOP** spraying: Turn control valve to the **OFF** position. Turn off water at faucet. To relieve (discharge) pressure, turn control valve to **ON** position and wait until water stops spraying before removing sprayer from hose.

See Table 1 for specific use instructions for either cool or warm-season turfgrass varieties. For preemergence applications, spray recommended rate evenly over entire lawn. For postemergence applications, spray weeds directly until well covered. Do not over-apply as this may cause undesirable turf injury.

SPECIFIC USE INSTRUCTIONS IN TURFGRASS

Established Turf

This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury

NOTE to EPA Product Manager: Text in brackets [] is optional.

may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

When applied as directed under the conditions described, the following established grasses are tolerant to F6875-5 RTS herbicide at the recommended use rates. Do not exceed 8.25 fl oz/1000 sq ft per calendar year.

Table 1. Tolerant grasses.

Grass Type	Recommended Use Rates Do not exceed maximum use rates per turfgrass species in a single application	
	Fluid ounces per 1000 sq ft	Area Covered (Based on 32 fl oz sprayer jar or retail container)
Cool Season Grasses		
Fescue, fine ¹ (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>), Bluegrass, Kentucky (<i>Poa pratensis</i>) Fescue, tall ² (<i>Festuca arundinacea</i>) Cool-season mixtures	2.75 fl oz Followed by 2.75 fl oz 30 – 60 days later (5.5 Tablespoons followed by 5.5 Tablespoons)	11,600 sq ft
Warm Season Grasses		
Bahiagrass (<i>Paspalum notatum</i>) Buffalograss (<i>Buchloe dactyloides</i>) Carpetgrass (<i>Axonopus affinis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kikuyugrass (<i>Pennisetum clandestinum</i>) Seashore Paspalum (<i>Paspalum vaginatum</i>) Zoysiagrass ³ (<i>Zoysia japonica</i>) Bermudagrass ⁴ (<i>Cynodon dactylon</i>) & hybrids	5.5 fl oz (8 Tablespoons)	5,800 sq ft
St. Augustinegrass	4.0 fl oz Followed by 4.0 fl oz 30 – 60 days later (8 Tablespoons followed by 8 Tablespoons)	8,000 sq ft
<p>8 Use of this product on certain varieties of Chewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury.</p> <p>2 F6875-5 RTS may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate listed per turf type or 8.25 fl oz/1000 sq ft per calendar year, whichever is lower.</p> <p>3 F6875-5 RTS application may cause temporary discoloration to exposed leaf surfaces on certain cultivars of zoysiagrass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply F6875-5 RTS on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.</p> <p>4 Sequential application of F6875-5 RTS may be needed in bermudagrass lawns for season-long control of yellow nutsedge. If sequential application is needed apply no more than 2.75 fl oz/1000 sq ft.</p>		

F6875-5 RTS has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties and or cultivars have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply F6875-5 RTS to a small area prior to treatment of larger areas.

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds, Grasses and Sedges

F6875-5 RTS herbicide will control or suppress the weeds listed in **Table 2** when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1. Do not apply adjuvants or surfactants with post applications of F6875-5 RTS unless tested on small areas prior to broadcast applications.

Table 2. Weeds (Preemergent and Postemergent) Controlled/Suppressed by Applications of F6875-5 RTS

Common Name	Scientific Name	Preemergent	Postemergent	
			Controlled	Suppressed
Barnyardgrass	(<i>Echinochloa crus-galli</i>)	X		
Bittercress	(<i>Cardamine spp.</i>)			X
Black medic	(<i>Medicago lupulina</i>)			X
Bluegrass annual	(<i>Poa annua</i>)	1		
Buttercups	(<i>Ranunculus spp.</i>)			X
Carolina geranium	(<i>Geranium carolinianum</i>)	X	5	
Carpetweed	(<i>Mollugo 8orniculate8</i>)	X		X
Chickweed, common	(<i>Stellaria media</i>)	2	5	
Chickweed, mousear	(<i>Cerastium vulgatum</i>)			X
Chickweed,mousear (from seed)	(<i>Cerastium vulgatum</i>)	X		
Cinquefoil	(<i>Potentilla spp.</i>)			X
Clover	(<i>Trifolium spp.</i>)			X
Crabgrass (Large and Smooth)	(<i>Digitaria spp.</i>)	3	6	
Crowfootgrass	(<i>Dactyloctenium aegyptium.</i>)	X		
Cudweed	(<i>Gnaphalium spp.</i>)			X
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)	X		

NOTE to EPA Product Manager: Text in brackets [] is optional.

Dandelion	(<i>Taraxacum officinale</i>)			X
Dock, Curly	(<i>Rumex crispus</i>)			X
Evening primrose	(<i>Oenothera biennis</i>)			X
Fiddleneck	(<i>Amsinckia spp.</i>)			X
Filaree	(<i>Erodium spp.</i>)			X
Foxtails, Annual	(<i>Setaria spp.</i>)	X		
Goldenrod	(<i>Solidago spp.</i>)		5	
Goosegrass	(<i>Eleusine indica</i>)	4	6	
Ground ivy	(<i>Glechoma hederacea</i>)		5	
Hairy Bittercress	<i>Cardamine hirsute</i>	X		
Henbit	(<i>Lamium amplexicaule</i>)	2		X
Itchgrass	(<i>Rottboellia exaltata</i>)	X		
Knotweed, prostrate	(<i>Polygonum aviculare</i>)	X		X
Kochia	(<i>Kochia scoparia</i>)	X	5	
Kyllinga, green	(<i>Kyllinga brevifolia</i>)	X		
Kyllinga, false	(<i>Kyllinga gracillima</i>)	X		
Lambsquarters, common	(<i>Chenopodium album</i>)	X		X
Lawn burweed	(<i>Soliva pterosperma</i>)			X
Lespedeza, common	(<i>Lespedeza striata</i>)			X
Lovegrass	(<i>Eragrostis spp.</i>)	X		
Mallow, common	(<i>Malva neglecta</i>)			X
Nutsedge, Yellow	(<i>Cyperus esculentus</i>)	X	5	
Panicum (Texas,Fall)	(<i>Panicum spp.</i>)	X		
Parsley piert	(<i>Alchemilla arvensis</i>)		5	
Pigweed, Redroot	(<i>Amaranthus retroflexus</i>)	X	5	
Pigweed, Tumble	(<i>Amaranthus albus</i>)		5	
Pineapple weed	(<i>Matricaria matricarioides</i>)			X
Plantain, buckhorn	(<i>Plantago lanceolata</i>)		5	
Puncture weed	(<i>Tribulus terrestris</i>)			X
Purslane, common	(<i>Portulaca oleracea</i>)	X		X
Pusley, Florida	(<i>Richardia scabra</i>)	X	5	
Redweed	(<i>Melochia corchorifolia</i>)			X
Rocket, London	(<i>Sisymbrium irio</i>)			X
Sedge, globe	(<i>Cyperus globulosus</i>)	X		
Sedge, cylindrical	(<i>Cyperus retrorsus</i>)	X		
Sedge, Surinam	(<i>Cyperus surinamensis</i>)	X		
Sedge, Texas	(<i>Cyperus polystachyos</i>)	X		
Shephardspurse	(<i>Capsella bursa-pastoris</i>)	2		
Signalgrass, broadleaf	(<i>Brachiaria playphylla</i>)	X		
Smartweed, Pennsylvania	(<i>Polygonum pensylvanicum</i>)		5	
Sorrel, Red	(<i>Rumex acetosella</i>)			X
Speedwell	(<i>Veronica spp.</i>)	X	5	
Sprangletop	(<i>Leptochloa spp.</i>)	X		
Spurge, annuals	(<i>Euphorbia spp.</i>)		5	
Spurge, prostrate	(<i>Euphorbia humistrata</i>)	X	5	
Spurge, spotted	(<i>Euphorbia 9ornicul</i>)		5	
Star of Bethlehem	(<i>Ornithogalum umbellatum</i>)		5	
Velvetleaf	(<i>Abutilon theophrasti</i>)			X
Violet, wild	(<i>Viola 9orniculat</i>)			X
Wild garlic	(<i>Allium vineale</i>)			X
Wild onion	(<i>Allium canadense</i>)			X
Witchgrass	(<i>Panicum capillare L.</i>)	X		
Woodsorrel, creeping	(<i>Oxalis 9orniculata</i>)			X
Woodsorrel, yellow	(<i>Oxalis stricta</i>)		5	
Woodsorrel, yellow (from Seed)	(<i>Oxalis stricta</i>)	X		

- 1 Poa annua is a winter annual. Applications of F6875-5 RTS should be made in August or September to established, non overseeded turf before Poa Annuu germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings.
- 2 Apply F6875-5 RTS in late summer, fall or winter before weed germinates for preemergent control.
- 3 Crabgrass control in cool season grasses from fall application: Areas where the ground is subject to winter freezes, F6875-5 RTS may be applied in the fall at rates listed in Table 1 (8-24 fluid ounces per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.
- 4 In areas of extended growing season and heavy pressure, preemergent weed control will be enhanced by second application of F6875-5 RTS 60-90 days after initial application
- 5 Weeds are suppressed at lower label rates (<5.5 fl oz/1000 sq ft). For optimum control apply rates of at least 5.5 fl oz per 1000 sq ft in a single application. Do not exceed the application rate specified for the turf species in Table 1.
- 6 F6875-5 RTS controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.

Application to Reseeded, Overseeded or Sprigged Areas

Reseeding, overseeding or sprigging of treated areas within three to six (3-6) months after application of this product could inhibit the establishment of desirable grasses. [See Table 3 for exact intervals from application].

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

NOTE to EPA Product Manager: Text in brackets [] is optional.

Table 3. Overseeding Intervals after F6875-5 RTS Applications

Total Amount of F6875-5 RTS Applied		Interval (Months Before Overseeding)	
Ounces Product/1000 sq ft	North	Transition	South
2.75	3	3	3
4.0	3	3	3
5.5	4	4	4
8.25	6	5	5

Other Use Precautions

- Do not use on turfgrasses other than those listed on this label.
- Do not graze or feed livestock, forage cut from areas treated with F6875-5 RTS herbicide.
- Do not apply directly to landscape ornamentals or ornamental beds.
- Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.
- Do not apply with adjuvants or surfactants.

STORAGE AND DISPOSAL

Pesticide Storage: Store and transport in upright position. Store only in original container, in a cool dry place inaccessible to children or pets.
Pesticide Disposal: *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain
Container Disposal: *If empty* – Non-refillable container. Do not reuse or refill this container. Offer for recycling, if possible.

WARRANTY

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents) call National Pesticide Information Center at 1-800-858-7378.

{Note to reviewer: The following are optional advertising and/or marketing phrases that may be present on promotional materials and container labels.}

- Tough Weed Killer
- Kills even the toughest weeds like crabgrass, dandelion and chickweed
- Kills Yellow Nutsedge, Crabgrass, Dandelions and other listed weeds
- Kills broadleaf and grassy weeds
- Starts working on contact
- Works through the shoots and roots of the weed
- Rainproof – won't wash away
- Rainfast in hours
- Provides excellent cool weather performance
- Performs in both cool and warm weather
- For use on Warm Season and Cool Season grasses
- Kills the weeds, not the lawn
- Water – Based
- 1 quart treats up to 11,600 sq ft
- This container covers up to 11,600 sq ft
- Won't harm lawn grasses when used as directed
- All-in-one weed control for lawns
- Provides Pre-emergent and Post-emergent weed control
- Kills weeds before they become a problem
- Kills the toughest weeds, down to the root
- Lawn – Friendly formula, when used as directed
- Weed Deadly – Lawn Friendly
- For home lawns only
- For home lawn care
- For a beautiful, weed-free lawn
- Helps keep lawns weed-free

- Controls dandelions before they emerge [from seed]
- Kill & Prevent Weeds [in your lawn]
- Ready to Spray
- Ready to Use

Non-FIFRA claims
Bonus size 20%