



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 21, 2018

Patricia Flis
FMC
2929 Walnut St.
Philadelphia, PA 19104

Subject: Label Amendment – for use rate and application interval table corrections
Product Name: Xonerate 2SC Herbicide
EPA Registration Number: 279-3621
Application Date: 12/29/2017
Decision Number: 538607

Dear Ms. Flis:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Autumn Metzger at 703-305-5314, or Metzger.autumn@epa.gov.

Page 2 of 2
EPA Reg. No. 279-3621
Decision No. 538607

Sincerely,

A handwritten signature in cursive script, appearing to read "Grant Rowland".

Grant Rowland
Acting Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

ACCEPTED

05/21/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 279-3621

AMICARBAZONE

GROUP

5

HERBICIDE

XONERATE 2SC HERBICIDE

For Selective Control of Weeds in Turf on Golf Courses, Sod Farms, Residential and Commercial Turf Sites, Park and Recreation Areas, Athletic Fields, School Grounds and other Turf Areas and Conifers in Nurseries and Field Plantings (including Christmas Trees)

EPA Reg. No. 279-3621

EPA Est. No.

Active Ingredient:

By Wt.

Amicarbazone:4-amino-N-(1, 1-dimethylethyl)-4,5-dihydro-3-(1-methylethyl)-5-oxo-1 H-1,2,4-triazole-1-carboxamide.....	23.25%
Other Ingredients:.....	76.75%
Total:	100.00%

XONERATE 2SC HERBICIDE contains 2.0 pounds per gallon of active ingredient Amicarbazone.

Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique à usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

[See back panel for additional precautionary statements]

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> •Call poison control center or doctor immediately for treatment advice. •Have person sip glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person.
If on skin:	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15 to 20 minutes. •Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with warm water for 15 to 20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye •Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
No specific antidote is available. Treat the patient symptomatically.	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information. For chemical emergencies (spill, leak, fire or exposure) call 1-800-424-9300.	
For Information Regarding the Use of This Product, Call: 1-800-321-1FMC (1362)	

Sold By: [Manufactured For:] [Distributed By:]

Net Contents: _____ gal/fl oz



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinylchloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR §170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL CHEMICAL HAZARDS

Do not mix or allow this product to come into contact with oxidizing agents. Hazardous chemical reaction may occur.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment wash waters or rinsate. Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not allow sprays to drift onto adjacent desirable plants. Drift or runoff may adversely affect non-target plants. Xonerate 2SC Herbicide has properties which may result in ground water contamination. Use of Xonerate 2SC Herbicide in areas where soils are permeable and groundwater is near the surface should be avoided.

To prevent damage to crops and other desirable plants, read and follow all directions and precautions on this label before using.

GROUND WATER ADVISORY

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Amicarbazone and its major degradates are known to leach through soil into ground water under certain conditions as a result of label use.

SURFACE WATER ADVISORY

This product may contaminate surface water via runoff of rain water or drift of spray in wind. This product is classified as having a high potential for reaching surface water via runoff several months or more after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Amicarbazone and its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States or local tribes may have in effect additional requirements regarding wellhead setbacks and operational containment. State, Local, or Tribal regulations may be more restrictive than those listed on this product label. The most restrictive requirements must be followed.

This product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours following application.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear coveralls, chemical-resistant gloves (made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber (includes natural rubber blends and laminates) \geq 14 mils, polyethylene, polyvinylchloride (PVC) \geq 14 mils, or viton \geq 14 mils), and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

Read these entire **DIRECTIONS FOR USE** and **WARRANTY AND DISCLAIMER STATEMENT** before using this product.

PRODUCT INFORMATION

Xonerate 2SC Herbicide is a selective, postemergence herbicide that may be used for the control of certain weeds in established turfgrasses. Xonerate 2SC Herbicide may be applied postemergence to tolerant turfgrasses at rates of 3 to 29 fl oz/A. [Xonerate 2SC Herbicide also has preemergence activity on certain weed species at rates of 14.5 to 29.0 fl oz Xonerate 2SC Herbicide/A]. See additional information for specific turfgrass species in section titled *Timing and Rates of Application*.

Xonerate 2SC Herbicide is formulated as a 23.25% suspension concentrate (SC) herbicide.

Apply Xonerate 2SC Herbicide either alone, in tank mix combination, sequentially prior to or after additional herbicides, except as labeled. When tank mixing, always observe all precautionary statements and limitations on labeling of all products.

Water is the recommended liquid carrier for Xonerate 2SC Herbicide.

Resistance-Management Recommendations

For resistance management, Xonerate 2SC Herbicide is a Group 5 herbicide. Any weed population may contain or develop plants naturally resistant to Xonerate 2SC and other Group 5 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Xonerate 2SC or other Group 5 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 1-800-321-1FMC (1362) [or visit www.fmcprosolutions.com]. You can also contact your pesticide distributor or university extension specialist to report resistance.

Read the entire DIRECTIONS FOR USE before using Xonerate 2SC Herbicide.

USE RESTRICTIONS

- Do not apply this product using aerial application equipment.
- Do not apply this product if the soil pH is greater than 7.4.
- Do not mix, load or clean spray equipment within 50 feet of any wells or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Do not apply within 50 feet of well-heads or the above mentioned aquatic systems.
- Do not apply more than 29 fl oz/A (fluid ounces per acre) of Xonerate 2SC Herbicide per season.
- Do not apply more than 29 fl oz/A of Xonerate 2SC Herbicide per single application.
- Do not allow this chemical to drift onto other crops or non-target plants.
- Do not apply this product through any type of irrigation system. Do not apply to turf under moisture stress. If necessary irrigate before application to eliminate moisture stress.

USE PRECAUTIONS

In order to avoid damaging turfgrasses:

- Do not apply Xonerate 2SC Herbicide where annual bluegrass is maintained as a desirable turf.
- Use on turfgrass species other than those listed in the Cool-Season Turfgrasses and Warm-Season Turfgrasses tables may result in unacceptable injury.
- Do not apply Xonerate 2SC Herbicide to newly established (within six months) turfgrasses, unless otherwise directed on this label.
- Do not apply Xonerate 2SC Herbicide less than four weeks before cutting or lifting sod.
- Do not replant any crop other than turfgrasses to treated areas for a period of 12 months after the last application.
- Do not apply when turfgrasses are under stress from adverse weather, excess heat, pests, disease, chemicals, mechanical injury or other related stress factors.
- To minimize turf injury, irrigate within 24 hours after application with 0.1 to 0.2 inches water.
- Allow at least 1 week between last application and overseeding with winter grasses.
- Apply to actively-growing weed populations.
- Do not apply to cool-season turf that has received an application of products that contain mefludide within the previous 3 months.

LATERAL MOVEMENT

Under certain conditions, Xonerate 2SC may have the potential to move laterally onto sensitive grasses. If high amounts of rainfall are predicted prior to application, it is advised to leave at least a 5 ft barrier between the sensitive species and edge of treated area to ensure an adequate buffer is established if lateral movement occurs.

SPRAY DRIFT MANAGEMENT

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states or tribes have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the application equipment must be set to apply medium to course droplets (i.e., ASAE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrow spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc., in order to minimize drift and optimize coverage and control.

Wind

Avoid making applications when spray particles may be carried by air currents to nontarget areas. Do not spray if wind is gusty, below 2 mph, or in excess of 10 mph and moving in the direction of adjacent sensitive areas. Local terrain may influence wind patterns; the applicator must be familiar with local conditions and understand how they may impact spray drift.

Sensitive Areas

Areas sensitive to this product are defined as bodies of water (ponds, lakes, rivers, streams, and wetlands), known habitats of threatened or endangered species, and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching sensitive areas.

Temperature Inversions

A surface temperature inversion (i.e., increasing air temperature with increasing altitude) greatly increases the potential for drift. Presence of ground fog is a good indicator of a surface temperature inversion. Do not apply during temperature inversions. Always make applications when there is some air movement to determine the direction and distance of

possible spray drift.

EQUIPMENT

All ground equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 ft above the ground surface.

RECOMMENDED TURFGRASS SITES

Xonerate 2SC Herbicide may be applied to tolerant turf species on golf courses, seed and sod production fields, residential sites (including home lawns), schools, playgrounds, parks, recreational areas, sports fields, common areas and grass roadsides maintained as a lawn.

Turfgrass Tolerance

Apply to turfgrasses established for six months, unless otherwise directed on this label. With the exception of Kentucky bluegrass, cool season turfgrasses established in the fall months can be treated the following spring. Kentucky bluegrass should be established for twelve months before application of Xonerate 2SC Herbicide. The use of Xonerate 2SC Herbicide on turf that is not well-established, or that has been weakened by weather, pests, diseases, chemicals, mechanical injury or other related stress factors may result in adverse turf injury.

This product should only be applied to sites that are composed of the following turfgrass species, unless trial use has indicated that the turf species not listed here is tolerant to Xonerate 2SC Herbicide.

Cool-Season Turfgrasses¹	
Bentgrass, Creeping	<i>Agrostis stolonifera</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, Fine	<i>Festuca spp.</i>
Fescue, Tall	<i>Festuca arundinacea</i>
Ryegrass, Perennial	<i>Lolium perenne</i>
Warm-Season Turfgrasses¹	
Bahiagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss	<i>Buchloe dactyloides</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
Seashore Paspalum	<i>Paspalum vaginatum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>

¹Cool- and warm-season turfgrasses have demonstrated acceptable tolerance to Xonerate 2SC Herbicide. However, not all cultivars have been evaluated.

When applied as directed under the conditions described, established turfgrasses are tolerant to this product. However, not all cultivars have been tested. As there are numerous cultivars of the listed turfgrasses as well as a wide range of clipping heights, end users are advised to determine tolerance on a small trial area on their site prior to application to all the intended turfgrass areas. Observe trial areas for two to three weeks after application. If injury is not considered adverse, then the remaining turf area may be treated. Temporary yellowing of the turf may occur after application. This effect is temporary and the turf will recover in 14 to 21 days.

Mixing Instructions – Liquid Carriers

Xonerate 2SC Herbicide is a suspension concentrate formulation that must be diluted before application. The proper mixing sequence for Xonerate 2SC Herbicide and recommended tank mixtures with the appropriate liquid carrier is as follows:

1. Fill the spray tank or nurse tank 1/4 full with the appropriate liquid carrier or water.
2. Start the recirculation and agitation system and continue throughout mixing and application.
3. If the compatibility test indicates the need for a compatibility agent, add the recommended amount to the spray tank.
4. If tank mixing with wettable powders, dry flowable or water dispersible granule products in water as the carrier, these may be added now.
5. Add the appropriate quantity of Xonerate 2SC Herbicide to the spray tank.
6. If tank mixing with suspension concentrates, add the products to the spray tank.
7. If tank mixing with emulsifiable concentrates or soluble products, add the products to the spray tank.
8. If mixing any additional spray adjuvants in the mixture, add them after all other products have been mixed.

9. Xonerate 2SC Herbicide and all registered mixtures should be kept agitated once mixed and then sprayed out.
10. Do not allow mixtures to stand for prolonged periods of time.

Compatibility

Xonerate 2SC Herbicide can be tank-mixed with other herbicides, pesticides and liquid fertilizers not shown on this label; however, compatibility needs to be determined before mixing. A compatibility test can be conducted with the following method:

1. In a small container (jar with lid), add 0.5 to 1 qt of the spray carrier.
2. Add all ingredients of the tank mix in the same ratio as the anticipated use into spray carrier.
3. Vigorously shake the mixture and allow to stand for 5 to 10 minutes.
4. Rapid precipitation, settling, changes in color, etc. and/or failure to resuspend indicates that the mixture is physically incompatible and should not be applied.
5. If the mixture is physically compatible, the tank mix should be tested to determine safety (no adverse turfgrass injury) before treating large areas. Observe treated areas for 2 to 3 weeks after application. If injury is not considered adverse, then the remaining turf area may be treated.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Sprayer Equipment Clean-out

After applying Xonerate 2SC Herbicide the sprayer must be thoroughly cleaned with the following procedure:

1. Drain the tank completely, then wash out tank, boom and hoses with clean water.
2. Drain the tank again.
3. Fill the tank half-full with water and add ammonia at a dilution rate of 1% v/v (i.e. one gallon household ammonia per 100 gal of rinsate). Completely fill tank with water. Operate the sprayer for 10 minutes to flush hoses, boom and nozzles.
4. Drain the sprayer system. Rinse the tank with clean water and flush the hoses, boom and nozzles. Remove and clean spray tips and screens separately.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State and local guidelines.

To avoid injury or exposure to non-target crops, thoroughly clean all mixing and spray equipment, including pumps, nozzles, lines and screens on an approved rinse pad or on a field site where an approved crop is grown. Do not drain or flush sprayer on or near desirable plants. Do not contaminate any body of water.

SPRAY APPLICATION

Broadcast Treatment

Accurately calibrate the sprayer prior to mixing the herbicide treatments. Apply Xonerate 2SC Herbicide at a minimum total spray volume of 20 gal/A using broadcast boom equipment. Power sprayers equipped with a spray wand/gun may also be used for broadcast. Application must be made at a sufficient spray pressure and volume to provide accurate and uniform application of spray particles to a given area without causing spray drift to non-target areas. If mixed with other labeled herbicides, the spray volume may be no less than the minimum volume recommended by any tank mix product used or 20 gal/A, whichever is greater. Use appropriately-sized mesh screens and in-line strainers. Agitate thoroughly before and during application with either bypass or mechanical agitation. Rinse the sprayer thoroughly with clean water immediately after each use.

Improper spray pattern overlaps produce areas of over-application and may increase the potential for turf damage.

Spot Treatment

Backpack and compression sprayers may be used for small turfgrass areas and spot treatments. Power sprayers equipped with a spray wand/gun may also be used for spot application. Wands fitted with a flat fan nozzle tip should be held stationary at the proper height during application. A side-to-side or swinging arm motion may result in uneven spray coverage. Spray target weed to ensure even coverage. Spraying to wet may increase risk of unwanted turf injury.

Improper spray pattern overlaps produce areas of over-application and may increase the potential for turf damage.

For Spot Applications: Use the Following		
Gallons	Fluid Ounces of Xonerate 2SC	Milliliters of Xonerate 2SC
1	0.07 fl. oz.	2 mL
2	0.14 fl. oz.	4 mL
4	0.28 fl. oz.	8 mL
Note: These rates are equivalent to 3 fl. oz./A		

WEEDS CONTROLLED OR SUPPRESSED

Depending on the use rate, Xonerate 2SC Herbicide will control or suppress the following broadleaf and grass weeds in turf.

WEEDS CONTROLLED OR SUPPRESSED by XONERATE 2SC HERBICIDE	
Alexandergrass	Lovegrass, Indian (suppression)
Betony, Florida (suppression)	Morningglorv spp.
Bittercress, hairy	Mustard, wild
Bluegrass, annual (annual biotypes)	Nightshade, eastern black (suppression)
Bluegrass, annual (perennial biotypes)(suppression)	Panicum, fall
Bluegrass, roughstalk (<i>Poa trivialis</i>) (suppression)	Parsley-piert
Burweed, lawn (spurweed)	Pennycress, field
Carpetweed	Pepperweed, Virginia
Cheat (suppression)	Pigweed spp.
Chickweed, common	Prickly sida (suppression)
Chickweed, sticky	Purslane, common
Clover, hop (suppression)	Radish, wild
Cocklebur, common (suppression)	Ragweed, common
Crabgrass spp. (suppression) ²	Ragweed, giant (suppression)
Creeping beggarweed (suppression)	Shepherd's-purse
Cudweed spp.	Sida, prickly (suppression)
Diamond-flower, old world (suppression)	Smartweed, Pennsylvania
Foxtail sp. (suppression)	Speedwell spp.
Dichondra	Spurge, spotted
Drymary, heartleaf	Tropical Signalgrass ¹
Hemp sesbania (suppression)	Toadflax, oldfield
Henbit	Velvetleaf
Horsenettle (suppression)	Waterhemp, tall
Jimsonweed	Waterhemp, common
Kochia (suppression)	Whitehead broom (suppression)
Lambsquarters, common	Woodsorrel, yellow
¹ For effective tropical signalgrass control make two applications with a tank mix herbicide, such as Dismiss® South	
² Does not control India crabgrass (<i>Digitaria longiflora</i>)	

Timing and Rates of Application
Cool-Season Turfgrasses

Established Cool Season Turf			
Site and Turfgrass Species	Pest	Rate	Application Intervals & Maximum Applications Per Year
Creeping Bentgrass (<i>Agrostis stolonifera</i>) Greens	Annual Bluegrass (<i>Poa annua</i>)	3 fl. oz./A	Reapply at 7-14 day intervals Up to Four Applications
Creeping Bentgrass (<i>Agrostis stolonifera</i>) Tees and Greens Surrounds	Annual Bluegrass (<i>Poa annua</i>)	3 fl. oz./A	Reapply at 14 day intervals Up to Four Applications
Creeping Bentgrass (<i>Agrostis stolonifera</i>) Fairways and Roughs	Annual Bluegrass (<i>Poa annua</i>)	3 fl. oz./A OR	Reapply at 14-21 day intervals Up to Four Applications
		6 fl. oz./A	Reapply at 21 day intervals Up to Two Applications
Kentucky Bluegrass (<i>Poa pratensis</i>)	Annual Bluegrass (<i>Poa annua</i>)	3 fl. oz./A OR	Reapply at 14 day intervals Up to Four Applications
		4 fl. oz./A OR	Reapply at 14 day intervals Up to Three Applications
		6 fl. oz./A	Reapply at 14-21 day intervals Up to Two Applications
Tall Fescue (<i>Festuca arundinacea</i>) Fine Fescue (<i>Festuca spp.</i>) Perennial Ryegrass (<i>Lolium perenne</i>)	Annual Bluegrass (<i>Poa annua</i>)	4 fl. oz./A OR	Reapply at 14 day intervals Up to Three Applications
		6-11 fl. oz./A	Reapply at 14-21 day intervals Up to Two Applications

Annual bluegrass (*Poa annua*) weed control in established creeping bentgrass, Kentucky bluegrass, fine fescues, tall fescue and perennial ryegrass.

Creeping Bentgrass-Putting Greens

Creeping bentgrass ('A1', 'A4', 'G6', 'L93', 'Crenshaw' and 'Penncross') maintained at putting green clipping heights has demonstrated acceptable tolerance to Xonerate 2SC Herbicide. Other creeping bentgrass cultivars may be tolerant; however, the end user is advised to do a tolerance test on a small area before spraying all the intended turfgrass areas. Additionally, all potential tank mixtures with fungicides, insecticides and other herbicides have not been evaluated and are not recommended by FMC Corporation for use on putting greens.

For annual bluegrass (*Poa annua*) control on creeping bentgrass putting greens, apply Xonerate 2SC Herbicide in the spring months, two to four weeks after the resumption of active bentgrass growth, at 3 fl oz/A. Repeat the application at 7 to 14 day intervals for a maximum of 4 applications per year.

- Apply Xonerate 2SC Herbicide at air temperatures ranging from 55 to 80 °F. Do not apply Xonerate 2SC Herbicide to creeping bentgrass when the daytime air temperature is expected to exceed 80°F
- Do not apply Xonerate 2SC Herbicide to creeping bentgrass in the summer or early fall months.

Xonerate 2SC Herbicide is effective in controlling *Poa annua* and results from controlling *Poa annua* infestations may be mistaken as temporary injury or thinning of the creeping bentgrass turf. Xonerate 2SC Herbicide is recommended only for creeping bentgrass areas with 10% or less population density of *Poa annua*. Applications to creeping bentgrass areas with high populations (> 10%) of *Poa annua* can result in a decrease of overall turf appearance due to the elimination of *Poa annua*. Creeping bentgrass will gradually fill in the resultant bare areas. Cultural practices to hasten the growth of creeping bentgrass will aid in the regeneration process. Treated areas may be reseeded at 7 or more days after the last application of Xonerate 2SC Herbicide. Do not retreat reseeded areas until turfgrass has become established,

Creeping Bentgrass-Tees and Greens Surrounds

Creeping bentgrass ('A1', 'A4', 'G6', 'L93', 'Pennway', 'Penncross', and 'Seaside') maintained at tee clipping heights has demonstrated acceptable tolerance to Xonerate 2SC Herbicide. Other creeping bentgrass cultivars may be tolerant; however, the end user is advised to do a tolerance test on a small area before spraying all the intended turfgrass areas. Additionally, all potential tank mixtures with fungicides, insecticides, other herbicides have not been evaluated and should be tested on a small scale basis before widespread use.

For annual bluegrass (*Poa annua*) control on creeping bentgrass tees and greens surrounds, apply Xonerate 2SC Herbicide in the spring months, two to four weeks after the resumption of active bentgrass growth, at 3.0 fl oz /A and repeat the application at 14 day intervals up to a maximum of 4 applications per year.

- Apply Xonerate 2SC Herbicide when air temperatures are ranging from 55 to 80 °F. Do not apply Xonerate 2SC Herbicide to creeping bentgrass when the daytime air temperature is expected to exceed 80°F.
- Do not apply Xonerate 2SC Herbicide to creeping bentgrass in the summer or early fall months.

Xonerate 2SC Herbicide is effective in controlling *Poa annua* and results from controlling *Poa annua* infestations may be mistaken as temporary injury or thinning of the creeping bentgrass turf. Xonerate 2SC Herbicide is recommended only for creeping bentgrass areas with 10% or less population density of *Poa annua*. Applications to creeping bentgrass areas with high populations (> 10%) of *Poa annua* can result in a decrease of overall turf appearance due to the elimination of *Poa annua*. Creeping bentgrass will gradually fill in the resultant bare areas. Cultural practices to hasten the growth of creeping bentgrass will aid in the regeneration process. Treated areas may be reseeded at 7 or more days after the last application of Xonerate 2SC Herbicide. Do not retreat reseeded areas until turfgrass has become established.

Creeping Bentgrass- Fairways and Roughs

Creeping bentgrass ('A1', 'A4', 'G6', 'L93', 'Pennway', 'Penncross' and 'Seaside') has demonstrated acceptable tolerance to Xonerate 2SC Herbicide. Other creeping bentgrass cultivars may be tolerant; however, the end user is advised to do a tolerance test on a small area before spraying all the intended turfgrass areas. Additionally, all potential tank mixtures with fungicides, insecticides, and other herbicides should be tested on a small scale basis before widespread use.

For annual bluegrass (*Poa annua*) control in creeping bentgrass fairways and roughs, apply Xonerate 2SC Herbicide in the spring months, two to four weeks after the resumption of active bentgrass growth. Apply at 3 fl oz /A and repeat the application at 14 to 21 day intervals up to a maximum of 4 applications per year. Alternatively, apply Xonerate 2SC Herbicide at up to 6 fl oz /A and repeat the application at a 21 day interval up to a maximum of 2 applications per year.

- Apply Xonerate 2SC Herbicide at air temperatures ranging from 55 to 80°F. Do not apply Xonerate 2SC Herbicide to creeping bentgrass when the daytime air temperature is expected to exceed 80°F
- Do not apply Xonerate 2SC Herbicide to creeping bentgrass in the summer or early fall months.

Xonerate 2SC Herbicide is effective in controlling *Poa annua* and results from controlling *Poa annua* infestations may be mistaken as temporary injury or thinning of the creeping bentgrass turf. Xonerate 2SC Herbicide is recommended only for creeping bentgrass areas with 10% or less population density of *Poa annua*. Applications to creeping bentgrass areas with high populations (> 10%) of *Poa annua* can result in a decrease of overall turf appearance due to the elimination of *Poa annua*. Creeping bentgrass will gradually fill in the resultant bare areas. Cultural practices to hasten the growth of creeping bentgrass will aid in the regeneration process. Treated areas may be reseeded at 7 or more days after the last application of Xonerate 2SC Herbicide. Do not retreat reseeded areas until turfgrass has become established.

Xonerate 2SC Herbicide may be tank mixed with labeled herbicides such as 2,4-D, MCP, dicamba or two and three-way herbicides that contain mixtures of 2,4-D, MCP, clopyralid, dicamba, fluroxypyr, carfentrazone-ethyl, or triclopyr to enhance control of emerged broadleaf weeds.

Xonerate 2SC Herbicide may also be tank mixed with preemergence herbicides for residual control of crabgrass species and other summer annual weeds.

Fine Fescues, Tall Fescues, and Perennial Ryegrass

For annual bluegrass (*Poa annua*) control in fine fescues, tall fescues and perennial ryegrass, apply Xonerate 2SC Herbicide at 6 to 11 fl oz/A and repeat the application at 14 to 21 day intervals up to a maximum of two applications per year. Alternatively, apply Xonerate 2SC Herbicide at up to 4 fl oz /A and repeat the application at 14 day intervals up to a maximum of 3 applications per year.

- Do not apply Xonerate 2SC Herbicide to fine fescues, tall fescues and perennial ryegrass when the daytime air temperature is expected to exceed 85 °F.

Perennial ryegrass and tall fescues may be seeded at 7 or more days after the last application of Xonerate 2SC Herbicide.

Xonerate 2SC Herbicide may be tank mixed with various recommended herbicides such as 2,4-D, MCP, dicamba or two and three-way herbicides that contain mixtures of 2,4-D, MCP, clopyralid, dicamba, fluroxypyr, carfentrazone-ethyl, or triclopyr to enhance control of emerged broadleaf weeds.

Xonerate 2SC Herbicide may also be tank mixed with preemergence herbicides for residual control of crabgrass species and other summer annual weeds.

Kentucky Bluegrass

For annual bluegrass (*Poa annua*) control in Kentucky bluegrass, apply Xonerate 2SC Herbicide in the spring months. Apply Xonerate 2SC Herbicide at 3 fl oz /A and repeat the application at 14 day intervals up to a maximum of 4 applications per year. Alternatively, apply Xonerate 2SC Herbicide at up to 4 fl oz /A and repeat the application at 14 day intervals up to a maximum of 3 applications per year or, alternatively, apply Xonerate 2SC Herbicide at up to 6 fl oz /A and repeat the application at 14 to 21 day intervals up to a maximum of 2 applications per year.

- Do not apply Xonerate 2SC Herbicide to Kentucky bluegrass when the daytime air temperature is expected to exceed 85 °F.
- **Kentucky Bluegrass should be established for twelve months before application of Xonerate 2SC Herbicide.**

Xonerate 2SC Herbicide may be tank mixed with various recommended herbicides such as 2,4-D, MCP, dicamba or two and three-way herbicides that contain mixtures of 2,4-D, MCP, clopyralid, dicamba, fluroxypyr, carfentrazone-ethyl, or triclopyr to enhance control of emerged broadleaf weeds.

Xonerate 2SC Herbicide may also be tank mixed with preemergence herbicides for residual control of crabgrass species and other summer annual weeds.

Warm Season Turfgrasses

Established Warm Season Turf			
Turfgrass Species	Pest	Rate	Application Intervals
Bahiagrass <i>(Paspalum notatum)</i> Bermudagrass <i>(Cynodon dactylon)</i> Buffalograss <i>(Bouteloua dactyloides)</i> Centipedegrass <i>(Eremochloa ophiuroides)</i> Kikuyugrass <i>(Pennisetum clandestinum)</i> Seashore Paspalum <i>(Paspalum vaginatum)</i> St. Augustinegrass <i>(Stenotaphrum secundatum)</i> Zoysiagrass <i>(Zoysia spp.)</i>	Annual Bluegrass <i>(Poa annua)</i>	9 – 29 fl. oz./A	Reapply at 7-21 day intervals Up to 3 Applications (max. 29 fl oz/A per year)
	Annual Broadleaf Weeds	Tropical Signal Grass	14.5 fl. oz./A
Bermudagrass Tees and Fairways Spring application following Fall overseeding with Perennial Ryegrass (Spring application)	Annual Bluegrass <i>(Poa annua)</i>	6 fl. oz/A	Reapply at 14-21 day intervals Up to 3 Applications
		OR 11 fl. oz/A	Reapply at 14-21 day intervals Up to 2 Applications
Bermudagrass Tees and Fairways Fall application prior to overseeding with Perennial Ryegrass or <i>Poa trivialis</i>	Annual Bluegrass <i>(Poa annua)</i>	14 – 29 fl. oz./A	One application only

Annual bluegrass and broadleaf weed control in established bahiagrass, bermudagrass, buffalograss, centipedegrass, kikuyugrass, seashore paspalum, St. Augustinegrass, tropical signalgrass and zoysiagrass.

For postemergence annual bluegrass and annual broadleaf weed control in tolerant warm-season turf, Xonerate 2SC Herbicide may be applied to warm-season turfgrasses that are dormant, partially dormant or actively-growing. Apply Xonerate 2SC Herbicide at rates of 9 to 29 fl oz/A. When using low rates (9 to 14 fl oz/A), repeat the application at 14 to 21 day intervals, up to a maximum of three applications per year. The maximum rate per season must not exceed 29 fl oz /A.

- St. Augustinegrass: Do not use Xonerate 2SC Herbicide on St. Augustinegrass when air temperatures are greater than 90°F.

Xonerate 2SC Herbicide can be tank mixed with labeled herbicides that contain metsulfuron, 2,4-D, dicamba or mixtures of 2,4-D, MCPP, clopyralid, dicamba, fluroxpyr, carfentrazone-ethyl, or triclopyr to enhance control of emerged broadleaf weeds. If sedge species are present on the site, Xonerate 2SC Herbicide can be tank mixed with herbicides labeled for control of sedges. Consult the tank mix partner label for turfgrass species tolerance, weeds controlled and maximum weed sizes at applications. This product can be tank mixed with Dismiss Turf Herbicide for the control of sedges. Follow the most restrictive labeled timing between Xonerate 2SC Herbicide and any tank mix partner.

Xonerate 2SC Herbicide may be tank mixed with preemergence herbicides for residual control of crabgrass species and other summer annual weeds.

Tropical Signal Grass Control in established bahiagrass, bermudagrass, buffalograss, centipedegrass, kikuyugrass, seashore paspalum, St. Augustinegrass, tropical signalgrass and zoysiagrass. For effective tropical signalgrass control Xonerate 2SC Herbicide may be applied in the spring or fall months. Fall is the best application time to maximize tropical signalgrass control. Initial fall applications typically are applied in September or October, depending on geographic location. Make two applications at 14 day intervals of 14.5 fl oz/A Xonerate 2SC as a tank mix with Dismiss® South. Other tank-mix options for effective tropical signalgrass control would be Tribute™ Total, Revolver® or Celsius®.

- St. Augustinegrass: Do not use Xonerate 2SC Herbicide on St. Augustinegrass in the summer months when air temperatures are greater than 90°F.

Fall Application for Annual bluegrass control in Bermudagrass Tees and Fairways Prior to Overseeding with Perennial Ryegrass or *Poa trivialis* (Roughstalk Bluegrass)

Apply Xonerate 2SC Herbicide at 14 to 29 fl oz/A seven or more days before overseeding bermudagrass with either perennial ryegrass or *Poa trivialis* for control of emerged annual bluegrass.

XONERATE 2SC HERBICIDE APPLIED TO SEEDLING *POA TRIVIALIS* WILL RESULT IN SEVERE TURF INJURY.

Spring Application for Annual bluegrass and broadleaf weed control in Bermudagrass Tees and Fairways following Fall Overseeding with Perennial Ryegrass

To control annual bluegrass and certain winter annual broadleaf weeds in bermudagrass overseeded the previous fall with perennial ryegrass apply Xonerate 2SC Herbicide in early- to mid-spring months, when perennial ryegrass is at least six months after seeding, is actively-growing and has recovered from winter-induced stress. Apply at a rate of 6 fl oz/A and repeat the application at 14 to 21 day intervals, up to a maximum of three applications per year. Alternately, apply at a rate of up to 11 fl oz/A and repeat the application at a 14 to 21 day interval up to a maximum of two applications per year. Do not apply Xonerate 2SC Herbicide to perennial ryegrass when the daytime air temperature is expected to exceed 80 °F.

- **Do not apply Xonerate 2SC Herbicide to perennial ryegrass that has received an application of products that contain mefludide within the previous 3 months unless severe injury and/or removal is acceptable.**

Do not apply Xonerate 2SC Herbicide to emerged *Poa trivialis* unless the objective is to suppress or remove *Poa trivialis*.

Xonerate 2SC Herbicide may be tank mixed with various recommended herbicides such as 2,4-D, MCPP, dicamba, sulfentrazone or two and three-way herbicides that contain mixtures of 2,4-D, MCPP, clopyralid, dicamba, fluroxypyr, carfentrazone-ethyl, or triclopyr to enhance control of emerged broadleaf weeds.

Xonerate 2SC Herbicide may also be tank mixed with preemergence herbicides for residual control of crabgrass species and other summer annual weeds.

Crabgrass Species Control in St. Augustinegrass

Xonerate 2SC Herbicide may be used in a tank-mixture for postemergence applications to reduce populations of blanket crabgrass and other crabgrass species (with the exception of India crabgrass) in established St. Augustinegrass. Apply 8.4 fl oz/A tank-mixed with Celsius® WG Herbicide at a 21 day interval for up to two applications per year. St. Augustinegrass may become slightly discolored by this application, but is transient. Complete recovery will occur within two weeks after the last application, depending upon growing conditions. Do not use Xonerate 2SC Herbicide on St. Augustinegrass when air temperatures are greater than 90°F.

Weed Control in Conifers in Nurseries and Field Plantings (including Christmas trees)

Xonerate 2SC Herbicide can be used over-the-top or as a directed spray under the canopy in conifer nurseries and field plantings (including Christmas trees). Apply Xonerate 2SC Herbicide at 11 to 29 fl oz/A to actively growing weeds listed above. Repeat applications may be made but do not exceed 29 fl oz/A per year.

RESTRICTION: Do not use a mechanically pressurized handgun for application to Christmas trees.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. For help with any spill, leak, fire or exposure involving this material, call day or night (800) 424-9300.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Rigid, nonrefillable in containers small enough to shake (5 gallons or 50 pounds or less) Nonrefillable container.

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or dispose of in a sanitary landfill or by incineration.

Rigid, nonrefillable in containers too large to shake (greater than 5 gallons or 50 pounds) Nonrefillable container.

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available or dispose of in a sanitary landfill.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent allowed by law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

[INSERT TRADEMARK STATEMENT AS APPLICABLE]

FMC – Trademarks of FMC Corporation 04/25/2018