



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

279-3637

Date of Issuance:

5/14/21

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

F9944-74

Name and Address of Registrant (include ZIP Code):

Paula Bodey  
FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Cynthia Giles-Parker, Chief  
Fungicide, Registration Division (7505P)

Date:

5/14/21

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 279-3637.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/30/2018

If you have any questions, please contact Marcel Howard by phone at (703-305-6784), or via email at [howard.marcel@epa.gov](mailto:howard.marcel@epa.gov)

Enclosure – stamped “accepted” label



FLUINDAPYR	Group	7	Fungicide
------------	-------	---	-----------

## F9944-74

{Alternate Brand Name: }

{When an alternate brand name is used, it will also be used in the body of the label.}

For use on Tree Nuts (Crop Group 14-12), Cereal Grains Except Rice (Crop Group 15)  
[almond, walnut, pecans and other tree nuts, corn (field corn, field corn grown for seed,  
popcorn, and sweet corn), grain sorghum, wheat, triticale, and barley].

**ACTIVE INGREDIENT:**

Fluindapyr 42.4%

**OTHER INGREDIENTS:** 57.6%

**TOTAL:** 100.0%

Contains 4 pounds per gallon of the active ingredient fluindapyr  
Suspension concentrate.

Keep Out of Reach of Children  
**CAUTION / PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en  
detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL TOLL FREE, DAY OR NIGHT, 1-800- 331-3148
---

– OR –

For MEDICAL EMERGENCY 1-866-331-3148	For SPILLS CHEMTREC 1-800-424-9300
---	--

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

**Note to PM**, the following bracketed statements are individually optional depending on the packaging configuration and whether a booklet label design is used:

- a.[See First Aid statement on back [label] [panel] [of booklet]].
- b.[See additional precautionary statements and Directions for Use in booklet.]
- c.[Read the entire label before using this product.]
- d.[See First Aid, Precautionary Statements, and Directions for Use on individual packages.]

EPA Reg. No. 279-

EPA Est. No. \_\_\_\_\_

Net Contents: \_\_\_\_\_

Sold by:  
FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

[ *placeholder* is a registered trademark of FMC]

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Caution.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing Harmful if inhaled. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes, and chemical resistant gloves.

#### FIRST AID

**If swallowed:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**If on skin:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**If inhaled:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

**If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Hotline Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

**Note to Physician:** No specific antidote. Treat symptomatically.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and all other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves of one of the following types; Barrier Laminate, Butyl Rubber  $\geq$  14 mils, Nitrite Rubber  $\geq$  14 mils, Neoprene Rubber  $\geq$  14 mils, Polyvinyl Chloride (PVC)  $\geq$  14 mils, and Viton  $\geq$  14 mils., and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **ENGINEERING CONTROLS STATEMENT**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS

### **USER SAFETY RECOMMENDATIONS**

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product.

Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinseate.

This product is not acutely toxic to pollinators; however, chronic exposure to the product through pollen and nectar may cause risk to pollinators. Protect pollinators by following label directions intended to limit exposure.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

[Note to Reviewer: Below is optional state specific language.]

[Optional language for CA: In California, this product may only be applied to fields that are not irrigated, or are irrigated using pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems.]

[Optional language for NY: Not for sale, distribution, or use in Nassau or Suffolk Counties of New York State.]

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is located in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves, and shoes plus socks.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store unused product in original container in a cool, dry, secure area.

**Pesticide Disposal:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or

Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling:**

**Nonrefillable container.** [equal to or less than 5 gallons]

Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times.

**Nonrefillable container.** [greater than 5 gallons] [and less than 260 gallons]

Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse as follows:

Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

*{Note to PM, IBC's may be nonrefillable or refillable, the container handling section will state either Nonrefillable or Refillable.}*

**Nonrefillable container.** {or}

**Refillable container.**

**Bottom discharge IBC.**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously



pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

## PRODUCT DIRECTIONS

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**Not for use in greenhouses.**

**Crop Rotation:** Fields treated with F9944-74 may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Tree Nuts (Crop Group 14-12), Cereal Grains Except Rice (Crop Group 15)	Immediately
All other crops	12 months

**Compatibility Jar Test:** F9944-74 is compatible with most products, however not all combinations have been tested. Use the following compatibility jar test to ensure physical compatibility.

Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add wettable powders and water dispersible granular products first, next liquid flowables, then emulsifiable concentrates, and last liquid soluble products. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. Use tank mix combinations on a small number of plants before treating larger areas. When tank mixing, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

**Use of Adjuvants:** An adjuvant may be used with F9944-74 unless otherwise specified in the crop use directions. F9944-74 is compatible with most products, however not all have been tested. Use the compatibility jar test to ensure physical compatibility.

**Tank Mixing:** F9944-74 may be tank mixed with other fungicides, herbicides, insecticides, and/or other additives unless prohibited on the label of the tank mix partner. Although F9944-74 is compatible with most products, not all combinations have been tested. Use the compatibility jar test to ensure physical compatibility. Before applying any tank mixture not specifically recommended on this label, the crop safety of the target crop should be confirmed by applying the mixture to a small area of the target crop in accordance to the label instructions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use on all product labels involved in tank mixing. Users must follow the most restrictive

directions for use and precautionary statements of each product in the tank mixture.

**Spray Equipment/Volume:**

F9944-74 should be applied in sufficient water volume to ensure thorough coverage as a foliar application for good disease control. Ground, air-blast, or aerial equipment may be used, so long as adequate spray coverage is achieved, unless prohibited in crop directions for use. For ground application, apply a minimum of 10 gallons of spray solution per acre.

For aerial application, apply a minimum of 10 gallons of spray solution per acre for tree nut crops, and a minimum of 2 gallons of spray solution per acre for all other crops.

**Resistance Management:** F9944-73 contains fluindapyr, a member of FRAC group 7 which inhibits succinate dehydrogenase (SDHI).

Resistant fungal strains may be selected when fungicides with the same mode of action (i.e. in the same FRAC group) are used repeatedly on the same pathogens within a single growing season. To reduce selection pressure and maintain good performance of F9944-74, do not exceed the number of applications specified in the crop use directions for each individual crop or crop group. F9944-74 should be used preventively (i.e. before disease occurs). For assistance on a particular crop and disease control situation, consult your local agricultural dealer, consultant, applicator, or state extension personnel for specific practices or recommendations in your area.

**Resistance Management Advisory:** In addition to the above requirements for the use of F9944-74, additional resistance management techniques are recommended to maintain the longevity of F9944-74 and other fungicides in FRAC group 7.

1. Use integrated pest management techniques to reduce overall disease pressure.
2. If tank mixed with a partner from a different FRAC group, use each partner at a rate that gives effective disease control when used individually.
3. Do not make more than two consecutive applications of fungicides in FRAC group 7. Rotate to products that have alternate modes of action (i.e. are **not** included in FRAC group 7).

F9944-74 contains the active ingredient fluindapyr which is a Group 7 fungicide based on the mode of action classification system. Repeated use of the same group of fungicides for a targeted disease may lead to the selection of resistant strains of fungi and result in reduced disease control.

To maintain performance of F9944-74 and other fungicides in the same groups, tank mix or rotate with a different fungicide group for good disease resistance management. For assistance on a particular crop and disease control situation, consult your local agricultural dealer, consultant, applicator, or state extension

personnel for specific practices or recommendations in your area.

**High Disease Pressure:** Where a range is given, use the higher rate of F9944-74 and/or reduced spray interval when disease pressure is high and/or conditions are favorable for disease development.

#### **RATE EQUIVALENCY TABLE**

Product rate fl oz/A 4 lb/gal	Equivalent fluindapyr Active Ingredient Rate (lb ai/A)
2.5	0.08
3.6	0.11
4.3	0.13
4.8	0.15
7.1	0.22

## **MANDATORY SPRAY DRIFT MANAGEMENT**

### **Airblast Applications:**

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- Do not apply during temperature inversions.

### **Ground Boom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy and have minimal bounce.
- Applicators are required to use a Medium to Coarse droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

### **Boomless Ground Applications:**

- Applicators are required to use a Medium to Coarse droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.
- Do not apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a

greater spray volume is needed, consider using a nozzle with a higher flow rate.

- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **BOOMLESS GROUND APPLICATIONS**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **HANDHELD TECHNOLOGY APPLICATIONS:**

- Take precautions to minimize spray drift.

## CROP USE DIRECTIONS

### Use Directions for Almond

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot <i>(Alternaria alternata)</i>  Scab <i>(Venturia carpophila)</i>  Shot Hole <i>(Wilsonomyces carpophilus)</i>	3.5 to 4.8	Apply preventatively after petal fall or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7-14
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not apply on almond until after petal fall.</li> <li>• Do not apply by handheld sprayer.</li> <li>• No single application may exceed 4.8 fl oz of product/A.</li> <li>• Do not apply more than 3 applications per year.</li> <li>• Do not apply more than 14.4 fl oz of product/A per year.</li> <li>• Do not apply more than 0.45 lb ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval: Do not apply within 30 days of harvest.</li> </ul>			

**Use Directions for Filbert (hazelnut), Pecan, Walnut (black and English) and Other Tree Nuts Crop Group 14-12. See below for list of crops.**

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Botryosphaeria blight <i>(Botryosphaeria dothidea)</i>  Eastern Filbert Blight <i>(Anisogramma anomala)</i>  Pecan Scab <i>(Fusicladium effusum)</i>	3.5 to 4.8	For Pecan Scab control, start application at bud break  For all other disease apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7 - 14

**List of Tree Nuts:** African nut-tree, Almond, Beechnut, Brazil nut, Brazilian pine, Bunya, Bur oak, Butternut, Cajou nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika nut, Ginkgo, Guiana chestnut, Hazelnut, Heartnut, Hickory nut, Japanese horse-chestnut, Macadamia nut, Mongongo nut, Monkey-pot, Monkey puzzle nut, Okari nut, Pachira nut, Peach palm nut, Pecan, Pequi, Pili nut, Pine nut, Pistachio, Sapucaia nut, Tropical almond, Walnut, black, Walnut, English, Yellowhorn, cultivars, varieties, and/or hybrids of these.

**Restrictions:**

- Do not apply by handheld sprayer.
- No single application may exceed 4.8 fl oz of product/A.
- Do not apply more than 3 applications per year.
- Do not apply more than 14.4 fl oz of product/A per year.
- Do not apply more than 0.45 lb ai of fluindapyr/A per year
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval: Do not apply within 30 days of harvest.



### Use Directions for Corn (Field Corn, Field Corn Grown for Seed, Sweet Corn, and Popcorn)

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
<p>Eyespot (<i>Kabatiella zea</i>)</p> <p>Gray leaf spot (<i>Cercospora zea-maydis</i>)</p> <p>Southern rust (<i>Puccinia polysora</i>)</p> <p>Common rust (<i>Puccinia sorghi</i>)</p> <p>Northern corn leaf blight (<i>Exserohilum turcicum</i>)</p> <p>Southern corn leaf blight (<i>Bipolaris maydis</i>)</p>	<p>2.5 to 4.3</p>	<p>Apply by aerial or ground application in a protective spray schedule or when weather conditions are favorable for disease development. Apply from onset of disease up through R4 growth stage on corn. A repeat application may be made 10-14 days later. When disease pressure is high, use the higher rate and shorter interval.</p> <p>For management of fungal pathogen resistance development, do not make more than 2 sequential applications per year before alternating to another fungicide from a FRAC group different from groups 3, 7, and 11.</p>	<p>10 to 14</p>
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply to sweet corn by mechanically pressurized handgun.</li> <li>No single application may exceed 4.3 fl oz of product/A.</li> <li>Do not apply more than 2 applications per year.</li> <li>Do not apply more than 8.6 fl oz of product/A per year.</li> <li>Do not apply more than 0.27 lb ai of fluindapyr/A per year.</li> <li>Do not use an adjuvant after the V8 stage and prior to the VT stage of corn. An adjuvant may be used at any other growth stage.</li> <li>Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed, and for hand harvesting and detasselling sweet corn is 14 days</li> </ul>			

The REI for all other activities is 12 hours.

- Preharvest Interval: PHI is 7 days of harvest for forage, and 30 days for stover and grain of field corn and popcorn. PHI for sweet corn is 14 days.

**Use Directions for Cereal Grains Except Rice (crop group 15) see below for list of crops. [Wheat (Spring and Winter), Triticale, and Barley]**

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Leaf rust ( <i>Puccinia triticina</i> , <i>P. hordei</i> )  Stripe rust ( <i>Puccinia striiformis</i> )  Stagonospora leaf / glume blotch ( <i>Parastagonospora nodorum</i> )  Tan spot ( <i>Pyrenophora tritici-repentis</i> )  Septoria leaf blotch ( <i>Zymoseptoria tritici</i> )  Stem rust ( <i>Puccinia graminis</i> )  Powdery mildew ( <i>Blumeria graminis</i> )	2.5 to 4.3	Apply F9944-74(Ground or Aerial) in a protective spray schedule or when weather conditions are favorable for disease development.  Apply lower rates early season.  Application of high rate at flag leaf ligule emergence protects the upper foliage during critical grain fill period.	10
<b>List of crops:</b> Barley, Buckwheat, Millet, pearl, Millet, proso, Oats, Rye, Teosinte, Triticale, Wheat			
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• No single application may exceed 4.3 fl oz of product/A.</li> <li>• Do not apply more than 2 applications per year.</li> <li>• Do not apply more than 8.6 fl oz of product/A per year.</li> <li>• Do not apply more than 0.27 lb ai of fluindapyr/A per year.</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval: PHI is 7 days prior to harvest of forage, 14 days prior to harvest of hay, and 30 days prior to harvest of grain/straw.</li> </ul>			

### Use Directions for Grain Sorghum

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Rust, Common ( <i>Puccinia purpurea</i> )	2.5 to 4.3	Apply by ground or aerial application.  Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	10
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• No single application may exceed 4.3 fl oz. of product/A.</li> <li>• Do not apply more than 2 applications per year.</li> <li>• Do not apply more than 8.6 fl oz of product/A per year.</li> <li>• Do not apply more than 0.27 lb ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval: Do not apply within 7 days of harvest for forage and 30 days of harvest for stover or grain.</li> </ul>			

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. To the extent consistent with applicable law, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

[*placeholder* are registered trademarks of FMC]

4-12-2021

[Label code 4120]