



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

November 13, 2024

Faith Kee  
Registration Manager  
FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

Subject: PRIA Label Amendment – add soybean  
Product Name: F9944-74  
EPA Registration Number: 279-3637  
Application Date: 8/2/2022  
Case Number: 479234

Dear Faith Kee:

The application referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable under FIFRA section 3(c)(5).

You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. The next label printing of this product must use this labeling unless subsequent changes have been approved. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of

statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Carmen Swinger at [swinger.carmen@epa.gov](mailto:swinger.carmen@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "C. L. Giles-Parker".

Cynthia L. Giles-Parker, Chief  
Fungicide Branch  
Registration Division (7505T)

Enclosure -stamped "accepted" label



FLUINDAPYR	Group	7	Fungicide
------------	-------	---	-----------

## F9944-74

*{Alternate Brand Name: }*

*{When an alternate brand name is used, it will also be used in the body of the label.}*

For use on Tree Nuts (Crop Group 14-12), Cereal Grains Except Rice (Crop Group 15)  
[almond, walnut, pecans and other tree nuts, corn (field corn, field corn grown for seed,  
popcorn, and sweet corn), grain sorghum, soybean, wheat, triticale, and barley].

**ACTIVE INGREDIENT:**

Fluindapyr 42.4%

**OTHER INGREDIENTS:** 57.6%

**TOTAL:** 100.0%

Contains 4 pounds per gallon of the active ingredient fluindapyr  
Suspension concentrate.

Keep Out of Reach of Children  
**CAUTION / PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en  
detalle.

(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

IN CASE OF A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL TOLL FREE, DAY OR NIGHT, 1-800- 331-3148
---

– OR –

For MEDICAL EMERGENCY 1-866-331-3148	For SPILLS CHEMTREC 1-800-424-9300
---	--

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

**Note to PM**, the following bracketed statements are individually optional depending on the packaging configuration and whether a booklet label design is used:

- a.[See First Aid statement on back [label] [panel] [of booklet]].
- b.[See additional precautionary statements and Directions for Use in booklet.]
- c.[Read the entire label before using this product.]
- d.[See First Aid, Precautionary Statements, and Directions for Use on individual packages.]

EPA Reg. No. 279-

EPA Est. No. \_\_\_\_\_

Net Contents: \_\_\_\_\_

Sold by:  
FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

[ *placeholder* is a registered trademark of FMC]

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Caution.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing Harmful if inhaled. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes, and chemical resistant gloves.

### FIRST AID

**If swallowed:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to by a poison control center or doctor.
- **DO NOT** give anything by mouth to an unconscious person.

**If on skin:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**If inhaled:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

**If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### Hotline Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

**Note to Physician:** No specific antidote. Treat symptomatically.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and all other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves of one of the following types; Barrier Laminate, Butyl Rubber  $\geq$  14 mils, Nitrite Rubber  $\geq$  14 mils, Neoprene Rubber  $\geq$  14 mils, Polyvinyl Chloride (PVC)  $\geq$  14 mils, and Viton  $\geq$  14 mils., and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **ENGINEERING CONTROLS STATEMENT**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**For all liquid formulations; for mixers and loaders:** Removable chemical extraction probes (also known as "stingers") used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

This product is not acutely toxic to pollinators; however, chronic exposure to the product through pollen and nectar may cause risk to pollinators. Protect pollinators by following label directions intended to limit exposure.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

[Note to Reviewer: Below is optional state specific language.]

[Optional language for CA: In California, this product may only be applied to fields that are not irrigated, or are irrigated using pressurized irrigation methods including macro-sprinkler, micro-sprinkler, or drip systems.]

[Optional language for NY: Not for sale, distribution, or use in Nassau or Suffolk Counties of New York State.]

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is located in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water, is: coveralls, chemical resistant gloves, and shoes plus socks.

### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store unused product in original container in a cool, dry, secure area.

**Pesticide Disposal:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or

Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling:**

**Nonrefillable container.** [equal to or less than 5 gallons]

**DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times.

**Nonrefillable container.** [greater than 5 gallons] [and less than 260 gallons]

**DO NOT** reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse as follows:

Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

*{Note to PM, IBC's may be nonrefillable or refillable, the container handling section will state either Nonrefillable or Refillable.}*

**Nonrefillable container.** {or}

**Refillable container.**

**Bottom discharge IBC.**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while



pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

## PRODUCT DIRECTIONS

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**Not for use in greenhouses.**

**Crop Rotation:** Fields treated with F9944-74 may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Tree Nuts (Crop Group 14-12), Cereal Grains Except Rice (Crop Group 15), soybean	Immediately
All other crops	12 months

**Compatibility Jar Test:** F9944-74 is compatible with most products, however not all combinations have been tested. Use the following compatibility jar test to ensure physical compatibility.

Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add wettable powders and water dispersible granular products first, next liquid flowables, then emulsifiable concentrates, and last liquid soluble products. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. Use tank mix combinations on a small number of plants before treating larger areas. When tank mixing, follow more restrictive labeling of any tank mix partner. **DO NOT** tank mix with any product that contains a prohibition on tank mixing.

**Use of Adjuvants:** An adjuvant may be used with F9944-74 unless otherwise specified in the crop use directions. F9944-74 is compatible with most products, however not all have been tested. Use the compatibility jar test to ensure physical compatibility.

**Tank Mixing:** Not all crops within a crop group or varieties, cultivars or hybrids of those crops have been tested for crop safety. F9944-74 may be tank mixed with other fungicides, herbicides, insecticides, and/or other additives unless prohibited on the label of the tank mix partner. Although F9944-74 is compatible with most products, not all combinations have been tested. Use the compatibility jar test to ensure physical compatibility. Before applying F9944-74 or any tank mixture not specifically

specified on this label, the crop safety of the target crop must be confirmed by applying the mixture to a small area of the target crop in accordance to the label instructions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Spray Equipment/Volume:**

F9944-74 must be applied in sufficient water volume to ensure thorough coverage as a foliar application for good disease control. Ground, air-blast, or aerial equipment may be used, so long as adequate spray coverage is achieved, unless prohibited in crop directions for use. For ground application, apply a minimum of 10 gallons of spray solution per acre.

For aerial application, apply a minimum of 10 gallons of spray solution per acre for tree nut crops, and a minimum of 2 gallons of spray solution per acre for all other crops.

**Resistance Management:** F9944-73 contains fluindapyr, a member of FRAC group 7 which inhibits succinate dehydrogenase (SDHI).

Resistant fungal strains may be selected when fungicides with the same mode of action (i.e. in the same FRAC group) are used repeatedly on the same pathogens within a single growing season. To reduce selection pressure and maintain good performance of F9944-74, DO NOT exceed the number of applications specified in the crop use directions for each individual crop or crop group. F9944-74 must be used preventively (i.e. before disease occurs). For assistance on a particular crop and disease control situation, consult your local agricultural dealer, consultant, applicator, or state extension personnel for specific practices or guidance in your area.

**Resistance Management Advisory:** In addition to the above requirements for the use of F9944-74, additional resistance management techniques are advised to maintain the longevity of F9944-74 and other fungicides in FRAC group 7.

1. Use integrated pest management techniques to reduce overall disease pressure.
2. If tank mixed with a partner from a different FRAC group, use each partner at a rate that gives effective disease control when used individually.
3. **DO NOT** make more than two consecutive applications of fungicides in FRAC group 7. Rotate to products that have alternate modes of action (i.e. are **not** included in FRAC group 7).

F9944-74 contains the active ingredient fluindapyr which is a Group 7 fungicide based on the mode of action classification system. Repeated use of the same group of fungicides for a targeted disease may lead to the selection of resistant strains of fungi and result in reduced disease control.

To maintain performance of F9944-74 and other fungicides in the same groups, tank mix or rotate with a different fungicide group for good disease resistance management. For assistance on a particular crop and disease control situation, consult your local agricultural dealer, consultant, applicator, or state extension personnel for specific practices or guidance in your area.

**High Disease Pressure:** Where a range is given, use the higher rate of F9944-74 and/or reduced spray interval when disease pressure is high and/or conditions are favorable for disease development.

## **MANDATORY SPRAY DRIFT MANAGEMENT**

### **Airblast Applications:**

- Sprays must be directed into the canopy.
- DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- DO NOT** apply during temperature inversions.

### **Ground Boom Applications:**

- User must only apply with the release height specified by the manufacturer, but no more than 4 feet above the ground or crop canopy and have minimal bounce.
- Applicators are required to use a Medium to Coarse droplet size (ASABE S572.1).
- DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- DO NOT** apply during temperature inversions.

### **Boomless Ground Applications:**

- Applicators are required to use a Medium to Coarse droplet size (ASABE S572.1) for all applications.
- DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom must remain level with the crop and have minimal bounce.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## **WIND**

Drift potential increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **BOOMLESS GROUND APPLICATIONS**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

## **HANDHELD TECHNOLOGY APPLICATIONS:**

- Take precautions to minimize spray drift.

## **RATE EQUIVALENCY TABLE**

<b>Product rate fl oz/A 4 lb/gal</b>	<b>Equivalent Fluindapyr Active Ingredient Rate (lb ai/A)</b>
<b>2.5</b>	<b>0.08</b>
<b>3.5</b>	<b>0.11</b>
<b>4.3</b>	<b>0.13</b>
<b>4.8</b>	<b>0.15</b>
<b>5.7</b>	<b>0.18</b>
<b>7.1</b>	<b>0.22</b>

## CROP USE DIRECTIONS

### Use Directions for Almond

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
<p>Alternaria Leaf Spot (<i>Alternaria alternata</i>)</p> <p>Brown rot blossom blight (<i>Monilinia spp.</i>)</p> <p>Scab (<i>Venturia carpophila</i>)</p> <p>Shot Hole (<i>Wilsonomyces carpophilus</i>)</p>	3.5 to 4.8	<p>For brown rot blossom blight, begin applications at pink bud stage (about 5% bloom). Repeat application at full bloom (about 80% bloom) if conditions are conducive for disease development. When conditions are less favorable for disease, a single application at 20-40% bloom may provide sufficient control.</p> <p>For all other diseases, apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rates and shorter interval when disease pressure is severe.</p>
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply by handheld sprayer.</li> <li>• No single application may exceed 4.8 fl oz of product (0.15 lb fluindapyr)/A.</li> <li>• Minimum retreatment interval (RTI) is 7 days.</li> <li>• <b>DO NOT</b> apply more than 3 applications per year.</li> <li>• <b>DO NOT</b> apply more than 14.4 fl oz of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.45 lb ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval (PHI) is 30 days.</li> </ul>		

**Use Directions for Filbert (hazelnut), Pecan, Walnut (black and English) and Other Tree Nuts Crop Group 14-12. See below for list of crops.**

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
Botryosphaeria blight ( <i>Botryosphaeria dothidea</i> )  Eastern Filbert Blight ( <i>Anisogramma anomala</i> )  Pecan Scab ( <i>Fusicladium effusum</i> )	3.5 to 4.8	For pecan scab initiate applications at bud break  For all other disease apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rates and shorter interval when disease pressure is severe.
<b>List of crops:</b> African nut-tree, Almond, Beechnut, Brazil nut, Brazilian pine, Bunya, Bur oak, Butternut, Cajou nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika nut, Ginkgo, Guiana chestnut, Hazelnut, Heartnut, Hickory nut, Japanese horse-chestnut, Macadamia nut, Mongongo nut, Monkey-pot, Monkey puzzle nut, Okari nut, Pachira nut, Peach palm nut, Pecan, Pequi, Pili nut, Pine nut, Pistachio, Sapucaia nut, Tropical almond, Walnut, black, Walnut, English, Yellowhorn, cultivars, varieties, and/or hybrids of these.		
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply by handheld sprayer.</li> <li>• No single application may exceed 4.8 fl oz of product (0.15 lb fluindapyr)/A.</li> <li>• Minimum retreatment interval (RTI) is 7 days.</li> <li>• <b>DO NOT</b> apply more than 3 applications per year.</li> <li>• <b>DO NOT</b> apply more than 14.4 fl oz of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.45 lb ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval (PHI) is 30 days</li> </ul>		

## Use Directions for Corn (Field Corn, Field Corn Grown for Seed, Sweet Corn, and Popcorn)

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
<p>Common rust (<i>Puccinia sorghi</i>)</p> <p>Eyespot (<i>Kabatiella zeae</i>)</p> <p>Gray leaf spot (<i>Cercospora zeae-maydis</i>)</p> <p>Northern corn leaf blight (<i>Exserohilum turcicum</i>)</p> <p>Southern corn leaf blight (<i>Bipolaris maydis</i>)</p> <p>Southern rust (<i>Puccinia polysora</i>)</p>	<p>2.5 to 4.3</p>	<p>Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Apply no later than growth stage R4 (early dough stage). Use higher rates and shorter interval when disease pressure is severe.</p>
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply to sweet corn by mechanically pressurized handgun.</li> <li>• No single application may exceed 4.3 fl oz of product (0.13 lb fluindapyr)/A.</li> <li>• Minimum retreatment interval (RTI) is 10 days.</li> <li>• <b>DO NOT</b> apply more than 2 applications per year.</li> <li>• <b>DO NOT</b> apply more than 8.6 fl oz of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.27 lb ai of fluindapyr/A per year.</li> <li>• <b>DO NOT</b> use an adjuvant after the V8 stage and prior to the VT stage of corn. An adjuvant may be used at any other growth stage.</li> <li>• Restricted Entry Interval (REI) is 14 days for detasselling corn grown for seed, and for hand harvesting sweet corn. The REI for all other activities is 12 hours.</li> <li>• Preharvest Interval (PHI) is 7 days for forage, 14 days for sweet corn, and 30 days for stover and grain.</li> </ul>		



**Use Directions for Cereal Grains Except Rice (crop group 15) see below for list of crops. [Wheat (Spring and Winter), Triticale, and Barley]**

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
<p>Leaf rust (<i>Puccinia triticina</i>, <i>P. hordei</i>)</p> <p>Powdery mildew (<i>Blumeria graminis</i>)</p> <p>Septoria leaf blotch (<i>Zymoseptoria tritici</i>)</p> <p>Stagonospora leaf / glume blotch (<i>Parastagonospora nodorum</i>)</p> <p>Stem rust (<i>Puccinia graminis</i>)</p> <p>Stripe rust (<i>Puccinia striiformis</i>)</p> <p>Tan spot (<i>Pyrenophora tritici-repentis</i>)</p> <p>Tan spot (<i>Pyrenophora tritici-repentis</i>)</p>	<p>2.5 to 4.3</p>	<p>Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.</p> <p>Apply lower rates in the early season.</p> <p>Application of high rate at flag leaf ligule emergence protects the upper foliage during critical grain fill period.</p>
<p><b>List of crops:</b> Barley, Buckwheat, Millet, pearl, Millet, proso, Oats, Rye, Teosinte, Triticale, Wheat cultivars, varieties, and/or hybrids of these.</p>		
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• No single application may exceed 4.3 fl oz of product (0.13 lb fluindapyr) /A.</li> <li>• Minimum retreatment interval (RTI) is 10 days.</li> <li>• <b>DO NOT</b> apply more than 2 applications per year.</li> <li>• <b>DO NOT</b> apply more than 8.6 fl oz of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.27 lb ai of fluindapyr/A per year.</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval(PHI) is 7 days for forage, 14 days for hay, and 30 days for straw or grain.</li> </ul>		

## Use Directions for Grain Sorghum

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
Rust, Common ( <i>Puccinia purpurea</i> )	2.5 to 4.3	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rates and shorter interval when disease pressure is severe.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• No single application may exceed 4.3 fl oz. of product (0.13 lb fluindapyr) /A.</li> <li>• Minimum retreatment interval (RTI) is 10 days.</li> <li>• <b>DO NOT</b> apply more than 2 applications per year.</li> <li>• <b>DO NOT</b> apply more than 8.6 fl oz of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.27 lb ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• Preharvest Interval (PHI) is 7 days for forage and 30 days for stover or grain.</li> </ul>		

## Use Directions for Soybean

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
Anthracnose ( <i>Colletotrichum truncatum</i> )  Brown spot ( <i>Septoria glycines</i> )  Cercospora leaf blight and purple seed stain( <i>Cercospora kikuchii</i> )  Frogeye leaf spot ( <i>Cercospora sojina</i> )  Powdery mildew ( <i>Erysiphe diffusa</i> )  Rust ( <i>Phakopsora pachyrhizi</i> )  Target spot ( <i>Corynespora cassiicola</i> )  White mold ( <i>Sclerotinia sclerotiorum</i> )	2.5 to 3.5	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rate and shorter interval when disease pressure is severe.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• No single application may exceed 3.5 fl. oz. of product (0.11 lb fluindapyr)/A.</li> <li>• Minimum retreatment interval (RTI) is 14 days.</li> <li>• <b>DO NOT</b> apply more than 2 foliar applications per year.</li> <li>• <b>DO NOT</b> apply more than 7 fl. oz. of product/A per year.</li> <li>• <b>DO NOT</b> apply more than 0.224 lb. ai of fluindapyr/A per year</li> <li>• Restricted Entry Interval (REI) is 12 hours.</li> <li>• <b>DO NOT</b> feed forage or hay to animals or permit animals to graze.</li> <li>• Preharvest Interval (PHI) is 7 days for forage or hay and 21 days for seed.</li> </ul>		

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. To the extent consistent with applicable law, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product.


To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

[placeholder are registered trademarks of FMC]

Label Code D-4714 072622032624

## SUPPLEMENTAL LABELING



### F9944-74

EPA Reg. No. 279-3637

FLUINDAPYR	Group	7	Fungicide
------------	-------	---	-----------

For soybean as a rotational crop.

*THIS SUPPLEMENTAL LABEL EXPIRES ON May 31, 2027  
AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE*

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Read the label affixed to the container for F9944-74 before applying. Use of F9944-74 according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the F9944-74 container.

**Crop Rotation:** Fields treated with F9944-74 may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Tree Nuts (Crop Group 14-12), Cereal Grains Except Rice (Crop Group 15), soybean	Immediately
All other crops	12 months

## SUPPLEMENTAL LABELING

### CROP USE DIRECTIONS

#### Use Directions for SOYBEAN

[Not for use in California]

Diseases	Product Rate (fl oz/A)	Application Directions
Anthracnose ( <i>Colletotrichum truncatum</i> )  Brown spot ( <i>Septoria glycines</i> )  Cercospora leaf blight and purple seed stain( <i>Cercospora kikuchii</i> )  Frogeye leaf spot ( <i>Cercospora sojina</i> )  Powdery mildew ( <i>Erysiphe diffusa</i> )  Rust ( <i>Phakopsora pachyrhizi</i> )  Target spot ( <i>Corynespora cassiicola</i> )  White mold ( <i>Sclerotinia sclerotiorum</i> )	2.5 to 3.5	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rate and shorter interval when disease pressure is severe.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>No single application may exceed 3.5 fl. oz. of product (0.11 lb fluindapyr)/A.</li> <li>Minimum retreatment interval (RTI) is 14 days.</li> <li><b>DO NOT</b> apply more than 2 foliar applications per year.</li> <li><b>DO NOT</b> apply more than 7 fl. oz. of product/A per year.</li> <li><b>DO NOT</b> apply more than 0.224 lb. ai of fluindapyr/A per year</li> <li>Restricted Entry Interval (REI) is 12 hours.</li> <li><b>DO NOT</b> feed forage or hay to animals or permit animals to graze.</li> <li>Preharvest Interval (PHI) is 7 days for forage or hay and 21 days for seed.</li> </ul>		