



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 1, 2021

Edward Bockrath
Product Registration Manager
FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Subject: Registration Review Label Mitigation for Thifensulfuron, Tribenuron, and
Metsulfuron
Product Name: Agility SG Herbicide (With TotalSol soluble granules)
EPA Registration Number: 279-9608 formerly 352-751
Application Date: 12/18/2017
Decision Number: 569774, 569775, 569776

Dear Mr. Bockrath:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the Sulfonylurea (SU) Herbicides Interim Decision, and has concluded that your submission is acceptable. The agency also completed review of your amended label referred to above, submitted in connection with registration under FIFRA, as amended, and has determined the label is also acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2
EPA Reg. No. 279-9608
Decision No. 569774, 569775, 569776

If you have any questions about this letter, please contact Jaclyn Pyne by phone at 703-347-0445, or via email at pyne.jaclyn@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a long horizontal flourish extending to the right.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure



Dicamba	Group	4	Herbicide
Metsulfuron methyl	Group	2	Herbicide
Tribenuron methyl	Group	2	Herbicide
Thifensulfuron methyl	Group	2	Herbicide

For Use on Wheat, Barley, Triticale and Fallow.

AGILITY® SG herbicide is a soluble granule for selective postemergence weed control.

Active Ingredients	By Weight
Sodium salt of dicamba (3,6-dichloro-2-Methoxybenzoic acid)*	63.6%
Thifensulfuron methyl Methyl 3-[[[(4-methoxy-6-methyl-1,3,5--triazin-2-yl) amino]carbonyl]amino]-sulfonyl]-2-thiophenecarboxylate	4.7%
Tribenuron methyl Methyl 2-[[[N-(4-methoxy-6-methyl-1,3,5--triazin-2-yl)methylamino]carbonyl]-amino]sulfonyl]benzoate	2.4%
Metsulfuron methyl Methyl 2-[[[(4-methoxy-6-methyl -1,3,5--triazin-2-yl)amino]carbonyl] amino]-sulfonyl]benzoate	1.9%
Other Ingredients	27.4%
TOTAL	100.0%

*This product contains 57.8% 3,6-dichloro-2-Methoxybenzoic acid (dicamba)

Contains 0.636 lb Sodium salt of dicamba* per pound
 Contains 0.047 lb Thifensulfuron Methyl per pound
 Contains 0.024 lb Tribenuron Methyl per pound
 Contains 0.019 lb Metsulfuron Methyl per pound

EPA Est. No. _____

Nonrefillable Container

Net: _____ OR

Refillable Container

Net: _____

EPA Reg. No. 279-9608

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.


**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 WARNING**

Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Sold By



FMC Corporation
 2929 Walnut Street
 Philadelphia, PA 19104



PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are made of any waterproof material.

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves (except for pilots).
- Shoes plus socks.
- Protective eyewear.

See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6).

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Apply this product only as directed on the label.

Groundwater Advisory

This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

Ground and Surface Water Protection

Point source contamination: To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below. Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.

Movement by surface runoff or through soil: Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the product information section of this label.

Movement by water erosion of treated soil: Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Avoid storage of pesticides near well sites.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGILITY® SG herbicide (with TotalSol® soluble granules) – referred to below as AGILITY SG herbicide - must be used only in accordance with instructions on this label or in supplemental FMC publications.

FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specified by FMC.

AGILITY SG herbicide is for use on wheat, barley, triticale and fallow in most states, check with your state extension service or Department of Agriculture before use, to be certain AGILITY SG herbicide is registered in your state. AGILITY SG herbicide is not registered for use in Alamosa, Conejos, Costilla, Rio Grande, and Saquache counties of Colorado unless use is directed otherwise by supplemental labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated areas if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over short-sleeve shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material.
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

PRODUCT INFORMATION

AGILITY SG herbicide is a water soluble granule that is used for selective postemergence weed control in wheat (including durum), barley, triticale and fallow.

The best control is obtained when AGILITY SG herbicide is applied to young, actively growing weeds. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

AGILITY SG herbicide is noncorrosive, nonflammable, and does not freeze. AGILITY SG herbicide should be mixed and completely dissolved in water and applied as a uniform broadcast spray (See Tank Mixtures and Mixing Instructions sections for use with Liquid Nitrogen Fertilizer Solutions).

RESTRICTIONS

- **DO NOT use AGILITY SG herbicide plus Malathion or Lorsban** as crop injury may result.
- DO NOT harvest sooner than 45 days after the last application of AGILITY SG herbicide.
- AGILITY SG herbicide is only registered on wheat, barley, triticale and fallow. DO NOT use on any other crop.
- DO NOT apply to wheat, barley or triticale undersown with legumes and grasses, because injury to the forages will result.
- DO NOT treat irrigation ditches or water used for crop irrigation or domestic uses.
- DO NOT apply this product through any type of irrigation system.

PRECAUTIONS

- Varieties of wheat (including durum), barley and triticale may differ in their response to various herbicides. FMC recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AGILITY SG herbicide to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after AGILITY SG herbicide application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix AGILITY SG herbicide with 2,4-D and apply after the crop is in the tillering stage of growth.
- Application to wheat, barley or triticale that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, may result in crop injury. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- For ground applications applied when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced. The addition of 2,4-D or MCPA may improve weed control under these conditions.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.

Sensitive Crop Precautions

AGILITY SG herbicide may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems, or foliage. These plants are most sensitive to AGILITY SG herbicide during their development or growing stage.

Directions to avoid herbicide drift near sensitive crops:

- DO NOT make applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. DO NOT spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and only sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays (volume median diameter of 400 microns or more) to avoid potential herbicide drift. Select nozzles that are designed to produce minimal amounts of fine spray particles (less than 200 microns). Examples of nozzles designed to produce coarse sprays via ground applications are Delavan® Raindrops, Spraying Systems XR (excluding 110° tips) flat fans, Turbo Teejets®, Turbo Floodjets®, or large capacity flood nozzles such as D10, TK10, or greater capacity tips.
- Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gallons per acre (for ground broadcast applications), unless otherwise required by the manufacturer of drift-reducing nozzles. Consult your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- DO NOT apply AGILITY SG herbicide adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85° F as drift is more likely to occur.
- Agriculturally approved drift-reducing additives may be used.
 - Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley or triticale.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

AGILITY SG herbicide is absorbed through the roots, shoots, and foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to three weeks after postemergence application to weeds (2 to 5 weeks for wild garlic), susceptible plants are controlled. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed.

AGILITY SG herbicide will provide up to 4 to 6 weeks of residual weed control. Susceptible weeds may germinate and emerge a few days after postemergence applications, but growth then ceases and leaves become chlorotic 3- 5 days after emergence. Death of leaf tissue and growing point will follow in some species, while others may remain green but stunted and noncompetitive.

AGILITY SG herbicide provides the best control of weeds in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not provide satisfactory control. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of AGILITY SG herbicide may be less effective on weeds stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, weeds hardened-off by drought stress are less susceptible to AGILITY SG herbicide.

WEED RESISTANCE MANAGEMENT

AGILITY SG herbicide, which contains the active ingredients Dicamba, Thifensulfuron methyl, Tribenuron methyl and Metsulfuron methyl is a group 2 and 4 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices.

Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance. The continued effectiveness of this product depends on the successful implementation of a weed resistance management program. To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of AGILITY SG herbicide for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.
- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple sites of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 2 and 4 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 2 and 4 herbicides.
- Avoid making more than two applications of AGILITY SG herbicide and any other Group 2 and 4 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.

- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

WEEDS CONTROLLED

AGILITY SG herbicide effectively controls the following weeds when used according to label directions:

Annual knawel	Kochia*
Annual sowthistle	Ladysthumb
Black mustard	Lambsquarter (common, slimleaf)
Black nightshade	London rocket
Blue/Purple mustard *	Marshelder
Broadleaf dock	Mayweed, chamomile
Buckwheat (tartary, wild)	Miners lettuce
Bur buttercup (testiculate)	Narrowleaf lambsquarters
Bushy wallflower/ Treacle mustard	Nightflowering catchfly
Canada thistle *	Pennsylvania smartweed
Carolina geranium	Pigweed (prostrate, redroot, rough, smooth, tumble)
Clasping pepperweed	Pineappleweed
Coast fiddleneck (tarweed)	Plains coreopsis
Common buckwheat	Prickly lettuce*
Common chickweed	Redmaids
Common cocklebur	Russian thistle*
Common mallow	Scentless chamomile / mayweed
Common purslane	Shepherd's-purse
Common radish	Smallflower buttercup
Common ragweed	Smallseed falseflax
Common sunflower	Smartweed (green, ladysthumb, pale)
Conical catchfly	Snow speedwell
Corn chamomile	Sticky chickweed
Corn cockle	Stinking mayweed / dogfennel
Corn gromwell *	Swinecress
Corn spurry	Tansymustard
Cowcockle	Tarweed fiddleneck
Cress (mouse-ear)	Tumble/ Jim Hill mustard
Curly dock	Velvetleaf
Cutleaf eveningprimrose	Volunteer lentils
False chamomile	Volunteer peas
Field chickweed	Volunteer sunflower
Field pennycress (fanweed)	Waterpod
Filaree (redstem, Texas)	Wild buckwheat
Flixweed *	Wild chamomile
Groundsel (common)	Wild garlic *
Henbit	Wild mustard
Knawel (German moss)	Wild radish *
Knotweed (prostrate)	

WEEDS PARTIALLY CONTROLLED**

AGILITY SG herbicide partially controls the following weeds when used according to label directions:

Broadleaf plantain
 Catchweed bedstraw
 Common dandelion
 Common yarrow
 Field bindweed

Mallow (little)
 Nightshade (cutleaf, hairy, silverleafs)
 Sowthistle (annual) *
 Tall waterhemp
 Vetch* (common, hairy)

* See the Specific Weed Problems section of this label for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest labeled rate of AGILITY SG herbicide and include a tank mix partner such as 2,4-D, MCPA, or bromoxynil (such as Buctril® herbicide, Bronate® herbicide or Bronate Advanced™ herbicide). Refer to the Tank Mixtures section of this label.

RATE CONVERSION CHART FOR AGILITY SG HERBICIDE

Ounces of AGILITY SG herbicide	Pounds of AGILITY SG herbicide	Active Ingredient	Pounds of Active Ingredient/A
1.6	0.1	Dicamba	0.0636 (Sodium salt of dicamba) 0.0578 (3,6-dichloro-2-Methoxybenzoic acid)
		Thifensulfuron methyl	0.0047
		Tribenuron methyl	0.0024
		Metsulfuron methyl	0.0019
2.4	0.15	Dicamba	0.0954 (Sodium salt of dicamba) 0.0867 (3,6-dichloro-2-Methoxybenzoic acid)
		Thifensulfuron methyl	0.0071
		Tribenuron methyl	0.0036
		Metsulfuron methyl	0.0029
3.2	0.2	Dicamba	0.1272 (Sodium salt of dicamba) 0.1156 (3,6-dichloro-2-Methoxybenzoic acid)
		Thifensulfuron methyl	0.0094
		Tribenuron methyl	0.0048
		Metsulfuron methyl	0.0038

LABELLED USES

AGILITY SG herbicide provides selective postemergence control of certain broadleaf weeds in wheat, barley, triticale, and fallow.

Fallow

Application and Use Rate Information	Use Rates (oz of AGILITY SG herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>AGILITY SG herbicide can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply AGILITY SG herbicide as a broadcast or spot treatment to emerged and actively growing weeds.</p> <p>Apply 1.6 to 3.2 oz/A of AGILITY SG herbicide to fallow fields (postharvest, fallow, crop stubble, set-aside). See SPRAY ADJUVANTS for proper adjuvant rates.</p>	1.6 to 3.2	Thifensulfuron methyl	0.0047 to 0.0094
		Tribenuron methyl	0.0024 to 0.0048
		Metsulfuron methyl	0.0019 to 0.0038
		Dicamba	0.0636 to 0.1272 (Sodium salt of dicamba) 0.0578 to 0.1156 (3,6-dichloro-2-methoxybenzoic acid)

RESTRICTIONS in Fallow:

- AGILITY SG herbicide is only registered for use on wheat, barley, triticale and fallow. DO NOT use on any other crop.
- DO NOT use less than 1.6 oz/A AGILITY SG herbicide.
- DO NOT apply more than 3.2 oz/A of AGILITY SG herbicide in a single application (maximum active ingredient per single application is 0.0094 lb/A thifensulfuron methyl, 0.0047 lb/A tribenuron methyl, 0.0038 lb/A metsulfuron methyl, and 0.0636 lb/A Sodium salt of dicamba).
- DO NOT apply more than 3.2 oz/A of AGILITY SG herbicide per year (maximum active ingredient per year of product is 0.0094 lb/A thifensulfuron methyl, 0.0047 lb/A tribenuron methyl, 0.0038 lb/A metsulfuron methyl, and 0.0636 lb/A Sodium salt of dicamba).
- DO NOT exceed two applications of AGILITY SG herbicide per year in Fallow when using reduced application rates.
- If using products that contain dicamba as a sequential treatment with AGILITY SG herbicide, DO NOT exceed 16.8 ounces acid equivalent dicamba per acre per year in fallow.
- DO NOT exceed 2.0 pounds acid equivalent dicamba per acre per year.

TANK MIXTURES IN FALLOW

AGILITY SG herbicide can be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY SG herbicide.

Wheat, Barley, and Triticale

Application and Use Rate Information	Use Rates (oz of AGILITY SG herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>Apply AGILITY SG herbicide at the rate of 1.6 to 3.2 oz/A to wheat, barley and triticale. See SPRAY ADJUVANTS for proper adjuvant rates.</p> <p>Use 3.2 oz/A of AGILITY SG herbicide for heavy infestation of the weeds listed under Weeds Partially Controlled when application timing and environmental conditions are marginal (refer to Biological Activity and Environmental Conditions section of this label for best performance).</p> <p>Use 1.6 to 2.4 oz/A of AGILITY SG herbicide for light infestation of the weeds listed under Weeds Controlled. Conditions at application should be optimum for effective treatment of these weeds.</p> <p>Note: See Tank Mix Section for additional info on required combinations when used at less than 3.2 oz/A.</p>	1.6 to 3.2	Thifensulfuron methyl	0.0047 to 0.0094
		Tribenuron methyl	0.0024 to 0.0048
		Metsulfuron methyl	0.0019 to 0.0038
		Dicamba	0.0636 to 0.1272 (Sodium salt of dicamba) 0.0578 to 0.1156 (3,6-dichloro-2-methoxybenzoic acid)

RESTRICTIONS in Wheat, Barley and Triticale:

- AGILITY SG herbicide is only registered for use on wheat, barley, triticale and fallow. DO NOT use on any other crop.
- DO NOT use less than 1.6 oz/A AGILITY SG herbicide.
- DO NOT apply more than 3.2 oz/A of AGILITY SG herbicide in a single application (maximum active ingredient per single application is 0.0094 lb/A thifensulfuron methyl, 0.0047 lb/A tribenuron methyl, 0.0038 lb/A metsulfuron methyl, and 0.0636 lb/A Sodium salt of dicamba).
- DO NOT apply more than 3.2 oz/A of AGILITY SG herbicide per year (maximum active ingredient per year of product is 0.0094 lb/A thifensulfuron methyl, 0.0047 lb/A tribenuron methyl, 0.0038 lb/A metsulfuron methyl, and 0.0636 lb/A Sodium salt of dicamba).
- DO NOT exceed two applications of AGILITY SG herbicide per year in Wheat, Barley and Triticale when using reduced application rates.
- If using products that contain dicamba as a sequential treatment with AGILITY SG herbicide, DO NOT exceed the following amounts of acid equivalent dicamba per acre per year: 8.4 ounces in wheat, 5.6 ounces in barley, and 2.1 ounces in triticale.
- DO NOT exceed 2.0 pounds acid equivalent dicamba per acre per year.
- PHI is 7 days for forage, 30 days for hay, and 45 days for wheat, barley and triticale.

APPLICATION TIMING – Wheat, Barley and Triticale

Fall-seeded Wheat, Barley, and Triticale

Make applications after the crop is in the 2-leaf stage but before the jointing stage.

Spring-seeded Wheat

Make applications after the crop is in the 2-leaf stage but before it exceeds the 6-leaf stage. Early developing wheat varieties (such as TAM 107, Madison, or Wakefield) must receive application between early tillering and the jointing stage (stagger planting dates with these varieties to be certain that the applications occur before jointing). Durum and Wampum variety must receive application after tillering but before it exceeds the 6-leaf stage and should be made in combination with 2,4-D. See Tank Mix Section for additional information.

Spring-seeded Barley

Make applications after the crop is in the 2-leaf stage but before it exceeds the 4-leaf stage.

DO NOT tank mix AGILITY SG herbicide with 2,4-D in early season applications on spring-seeded barley.

Spring-seeded Triticale

Make applications after the crop is in the 2-leaf stage but before it exceeds the 6-leaf stage.

Additional Timing Information

Weed control may be reduced if rainfall or snowfall occurs soon after application. Six hours of dry weather are needed to allow AGILITY SG herbicide to be sufficiently absorbed by weed foliage.

If applied to irrigated wheat, barley or triticale the first post-treatment irrigation should be delayed for at least 6 hours after treatment and should not exceed 1 inch of water.

Applications of AGILITY SG herbicide to stressed crops may cause crop injury. To reduce the potential of crop injury, tank mix AGILITY SG herbicide with 2,4-D (ester formulations perform best-see TANK MIXTURES for more information) and apply after the crop is in the tillering stage of growth.

TANK MIXTURES

AGILITY SG herbicide may be tank mixed with other suitable registered herbicides to control weeds listed as partially controlled, weeds resistant to AGILITY SG herbicide or weeds not listed under Weeds Controlled. Read and follow all manufacturers label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY SG herbicide. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AGILITY SG herbicide can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley and triticale.

With 2,4-D (amine or ester) or MCPA (amine or ester)

AGILITY SG herbicide can be tank mixed with 2,4-D and MCPA (preferably ester formulations) herbicides for use on wheat, barley, triticale and fallow. For best results, add 2,4-D or MCPA herbicides to the tank at 1/8 to 3/8 lb active ingredient per acre. In tank mixes containing 1/8 lb active ingredient 2,4-D or MCPA per acre, add 1 to 2 pt of non-ionic surfactant per 100 gal of spray solution; in tank mixes containing 1/4 to 3/8 lb active ingredient 2,4-D or MCPA per acre, add 1 pt of non-ionic surfactant per 100 gal of spray solution. Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels.

Always mix AGILITY SG herbicide in water prior to adding 2,4-D or MCPA and add the surfactant last. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

With Bromoxynil containing products (including Buctril® herbicide and "Bronate®" branded herbicides)

AGILITY SG herbicide may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, triticale or fallow. For best results, add bromoxynil-containing herbicides to the tank at 3 to 6 oz/A active ingredient (such as Bronate® herbicide at 3/4 - 1 1/2 pt per acre).

Tank mixes of AGILITY SG herbicide plus Bromoxynil may result in reduced control of Canada thistle.

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling.

With Stinger® herbicide or Curtail® herbicide

AGILITY SG herbicide can be tank mixed with Stinger® or Curtail® herbicide for improved control of weeds in wheat, barley and triticale.

Refer to the Stinger® herbicide and Curtail® herbicide labels for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. DO NOT use the tank mix if any restrictions on the Stinger® herbicide or Curtail® herbicide labels conflict with instructions on the AGILITY SG herbicide label.

With Puma® 1EC herbicide

AGILITY SG herbicide at 1.6 oz/A can be tank mixed with Puma® 1EC herbicide for use in wheat and barley for control of green foxtail, foxtail millets, volunteer corn, and light infestations of broadleaf weeds on WEEDS CONTROLLED list. Refer to the Puma® 1EC herbicide label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the Puma® 1EC herbicide label conflict with instructions on the AGILITY SG herbicide label.

With Discover® NG herbicide

AGILITY SG herbicide at 1.6 -2.4 oz/A can be tank mixed with Discover® NG herbicide for improved control of weeds in wheat. Refer to the Discover® NG herbicide label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. DO NOT use the tank mix if any restrictions on the Discover® NG herbicide label conflict with instructions on the AGILITY SG herbicide label.

With "Everest®" branded herbicides

AGILITY SG herbicide can be tank mixed with "Everest®" branded herbicide for improved control of weeds in wheat. Refer to the "Everest®" branded product label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Everest®" branded product label conflict with instructions on the AGILITY SG herbicide label.

With Other Herbicides

AGILITY SG herbicide may be tank mixed with other suitable registered cereal or fallow herbicides to control weeds listed as suppressed, weeds resistant to AGILITY SG herbicide, or weeds not listed under Weeds Controlled. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY SG herbicide. Tank mixes of AGILITY SG herbicide plus metribuzin may result in reduced control of wild garlic.

DO NOT tank mix AGILITY SG herbicide with Hoelon® 3EC herbicide, because grass control may be reduced.

DO NOT tank mix AGILITY SG herbicide with ALLY® XP herbicide, ALLY® EXTRA herbicide, ALLY® EXTRA SG herbicide or other products containing metsulfuron methyl.

DO NOT tank mix AGILITY SG herbicide with other dicamba containing products (such as Clarity® herbicide or Banvel® herbicide) for use on cereal crops.

With Fungicides

AGILITY SG herbicide may be tank mixed or used sequentially with fungicides registered for use on cereal crops.

With Insecticides

RESTRICTIONS

- DO NOT apply AGILITY SG herbicide within 60 days of crop emergence where an organophosphate insecticide has been applied as an infurrow treatment, because crop injury may result.
- DO NOT use AGILITY SG herbicide plus Malathion or Lorsban as crop injury may result.

PRECAUTIONS

AGILITY SG herbicide may be tank mixed or used sequentially with insecticides registered for use on cereal crops. However, under certain conditions (drought stress, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of AGILITY SG herbicide with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

With Liquid Nitrogen Fertilizer Solution

Liquid nitrogen fertilizer solutions (e.g., 28-0-0, 32-0-0) may be used as a carrier in place of water. Run a tank mix compatibility test before mixing AGILITY SG herbicide in fertilizer solution.

AGILITY SG herbicide must first be dissolved with water and then added to liquid nitrogen solutions. Ensure that the agitator is running while the AGILITY SG herbicide is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pt -1 qt per 100 gal of spray solution (0.06 -0.25% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. If 2,4-D or MCPA is included with AGILITY SG herbicide and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Additional surfactant may not be needed when using AGILITY SG herbicide in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions. Consult your agricultural dealer, consultant, field advisor, or FMC representative for specific instructions before adding an adjuvant to these tank mixtures.

Note: In certain areas east of the Mississippi river unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant, field advisor, or FMC representative for specific instructions before using nitrogen fertilizer carrier solutions.

DO NOT use low rates of liquid fertilizer as a substitute for a surfactant.

DO NOT use with liquid fertilizer solutions with a pH less than 3.0.

SPECIFIC WEED PROBLEMS

Note: Thorough spray coverage of all weed species listed below is very important.

Blue Mustard: For best results, use 2.4-3.2 oz/A and apply AGILITY SG herbicide in tank mixtures with 2,4-D or MCPA postemergence to mustards, but before bloom (refer to Tank Mixtures section of this label for additional details).

Flixweed: For best results, use 2.4-3.2 oz/A and apply AGILITY SG herbicide in tank mixtures with 2,4-D or MCPA postemergence, but before bloom (refer to Tank Mixtures section of this label for additional details).

Canada Thistle: For best results, use 3.2 oz/A and apply AGILITY SG herbicide plus 2,4-D, or MCPA (refer to Tank Mixtures for additional details) in the spring after the majority of thistles have emerged and are small (rosette stage to 6" elongating stems) and actively growing. The application will inhibit the ability of emerged thistles to compete with the crop.

Corn Gromwell: For best results, use 2.4-3.2 oz/A and apply AGILITY SG herbicide when weeds are actively growing, are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing 2,4-D, MCPA, or bromoxynil containing products (including Buctril® herbicide, Bronate® herbicide, or Bronate Advanced™ herbicide) with AGILITY SG herbicide usually improves results (refer to Tank Mixtures section of this label for additional details).

Kochia, Prickly Lettuce, and Russian Thistle: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use AGILITY SG herbicide at 2.4 to 3.2 oz/A in a tank mix with 2,4-D and apply in the spring when weeds are less than 2" tall or wide and are actively growing (refer to Tank Mixtures section of this label for additional details).

Vetch (common and hairy): For best results, use 3.2 oz/A and apply AGILITY SG herbicide when vetch is less than 6" in length. For severe infestations of vetch, or when vetch is greater than 6" in length, use AGILITY SG herbicide in combination with 2,4-D, or MCPA (refer to Tank Mixtures section of this label for additional details).

Wild garlic: For best results, use 3.2 oz/A and apply AGILITY SG herbicide when wild garlic plants are less than 12" tall with 2" to 4" of new growth. Plants hardened-off by cold weather and/or drought stress may be more difficult to control. Thorough spray coverage of all garlic plants is essential. Typical symptoms of dying garlic plants may not be noticeable for 2 to 5 weeks. Control will be improved by using AGILITY SG herbicide in combination with 2,4-D or MCPA (refer to Tank Mixtures section of this label for additional details).

Wild radish: For best results, use 3.2 oz/A applied in the fall to wild radish rosettes less than 6" in diameter and before plants harden-off. Alternatively, AGILITY SG herbicide can be applied in the spring for control of wild radish. Control will be improved by using AGILITY SG herbicide in combination with 2,4-D or MCPA (refer to Tank Mixtures section of this label for additional details) when wild radish rosettes are less than 6" in diameter. Applications made later than 30 days after weed emergence, either in the fall or spring, will result in partial control.

PRODUCT MEASUREMENT

AGILITY SG herbicide can be measured using the AGILITY SG herbicide volumetric measuring cylinder included in the case. The degree of accuracy of this cylinder varies by $\pm 7.5\%$. For more precise measurement, use scales calibrated in ounces.

SPRAY ADJUVANTS

Always include a spray adjuvant with applications of AGILITY SG herbicide. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used. Antifoaming agents may be needed.

Consult your Ag dealer or applicator, local FMC fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with AGILITY SG herbicide, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40CFR 1001).

Nonionic Surfactant (NIS)

- Apply 0.125 to 0.50% volume/volume (1 pt to 4 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.
- See the Tank Mixtures section of this label for additional information.

Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Only use adjuvants that contain COC or MSO in fallow applications. Do not use for postemergence applications in cereals.
- Apply at 1% volume/volume (1 gal per 100 gal spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.
- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by FMC product management. Consult separate FMC technical bulletins for detailed information before using adjuvant types not specified on this label.

Ammonium Nitrogen Fertilizer

• Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.

Application and Spray Volumes

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. **HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING.**

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 2 to 10 gallons of diluted spray per treated acre when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

AGILITY SG herbicide should not be applied during periods of gusty wind or when wind is in excess of 15 mph as uneven spray coverage may occur.

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).

For flood nozzles on 30" spacing, use flood nozzles no larger than TK10 (or the equivalent), a pressure of at least 30 psi and a spray volume of at least 10 GPA only. For 40" nozzle spacing, use at least 13 GPA; for 60" spacing use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

"Raindrop RA" nozzles are not recommended for AGILITY SG herbicide applications, because weed control performance may be reduced.

Use screens that are 50-mesh or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, Washington, or Utah.

When applying AGILITY SG herbicide by air in areas near sensitive crops, use solid-stream nozzles oriented straight back. Adjust swath to avoid spray drift damage to downwind sensitive crops and/or use ground equipment to treat border edge of field. See the **Spray Drift Management** section of this label.

For aerial application in Washington, follow the directions in the Spray Drift Management Section of this label and the following Washington state restrictions:

- Applications of AGILITY SG herbicide must be made in equipment that meets the most restrictive Washington Agricultural Codes (WAC) for the prevention of herbicide drift for the respective county.
- DO NOT apply in equipment that does not meet these WAC standards.

SEQUENTIAL APPLICATIONS

AGILITY SG herbicide can be applied either before or after applications of other products registered for use in wheat, barley, triticale or fallow. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these in sequence with AGILITY SG herbicide. If those instructions conflict with this label, do not use that product in sequence with AGILITY SG herbicide. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

- DO NOT use any metsulfuron methyl-containing product as a sequential treatment with AGILITY SG herbicide.
- If using HARMONY® EXTRA XP herbicide as a sequential treatment with AGILITY SG herbicide, DO NOT exceed 0.7 oz/A of HARMONY® EXTRA XP herbicide per year.

- If using HARMONY® EXTRA SG herbicide (with TotalSol® soluble granules) as a sequential treatment with AGILITY SG herbicide, DO NOT exceed 1.0 oz/A of HARMONY® EXTRA SG herbicide per year.
- If using EXPRESS® XP herbicide as a sequential treatment with AGILITY SG herbicide, DO NOT exceed 0.25 oz/A of EXPRESS® XP herbicide per year.
- If using EXPRESS® herbicide (with TotalSol® soluble granules) as a sequential treatment with AGILITY SG herbicide, DO NOT exceed 0.375 oz/A of EXPRESS® herbicide per year.
- If using products that contain dicamba as a sequential treatment with AGILITY SG herbicide, do not exceed the following amounts of acid equivalent dicamba per acre per season: 8.4 ounces in wheat, 5.6 ounces in barley, 2.1 ounces in triticale, and 16.8 ounces in fallow. Do not exceed 2.0 pounds acid equivalent dicamba per acre per year.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of AGILITY SG herbicide.
3. Continue agitation until the AGILITY SG herbicide is fully dissolved, at least 5 minutes.
4. Once the AGILITY SG herbicide is fully dissolved, maintain agitation and continue filling tank with water. AGILITY SG herbicide should be thoroughly dissolved with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of surfactant. Always add surfactant last. Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability of AGILITY SG herbicide.
6. Dispersed tank mix partners can settle if the tank mixture is not continually agitated.
7. Apply AGILITY SG herbicide spray mixture within 24 hours of mixing to avoid product degradation.
8. If AGILITY SG herbicide and a tank mix partner are to be applied in multiple loads, fully dissolve the AGILITY SG herbicide in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the AGILITY SG herbicide.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's instructions for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

DO NOT make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Sprayer Precaution

The spray equipment must be cleaned before AGILITY SG herbicide is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of AGILITY SG herbicide are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING AGILITY SG HERBICIDE AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY OR TRITICALE

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of AGILITY SG herbicide as follows:

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Fill the tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight. Flush the

boom well and empty the sprayer. Completely drain the sump.

4. Remove the nozzles and screens and clean separately in a bucket containing water.

The rinsate solution may be applied to the crop(s) specified on this label. Do not exceed the maximum-labeled use rate. If cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

Notes:

1. Always start with a clean spray tank.

2. Steam-cleaning aerial spray tanks is recommended to facilitate the removal of any caked deposits.

3. When AGILITY SG herbicide is tank mixed with other pesticides, all cleanout procedures for each product should be examined and the most rigorous procedure should be followed.

4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.

- If using products that contain dicamba as a sequential treatment with AGILITY SG herbicide, do not exceed the following amounts of acid equivalent dicamba per acre per season: 8.4 ounces in wheat, 5.6 ounces in barley, 2.1 ounces in triticale, and 16.8 ounces in fallow.

CROP ROTATION

Before using AGILITY SG herbicide carefully consider your crop rotation plans and options. For rotational flexibility, DO NOT treat all of your acres at the same time.

MINIMUM ROTATIONAL INTERVALS

Minimum rotation intervals* are determined by the rate of breakdown of AGILITY SG herbicide applied. AGILITY SG herbicide breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase AGILITY SG herbicide breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow AGILITY SG herbicide breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting. **Minimum rotation intervals must be extended 1 crop season if drought conditions prevail after application and before the rotational crop is planted.**

Soil pH Limitations

AGILITY SG herbicide should not be used on soils having a pH above 7.9, because extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, AGILITY SG herbicide could remain in the soil for 34 months or more, injuring wheat, barley or triticale. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of AGILITY SG herbicide.

Checking Soil pH

Before using AGILITY SG herbicide, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0” to 4” samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

FIELD BIOASSAY

A field bioassay is necessary if crops other than wheat, barley or those listed on this label are to be planted on land previously treated with AGILITY SG herbicide. To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with AGILITY SG herbicide. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips.

If a field bioassay is planned, check with your local FMC representative for information detailing field bioassay procedure.

GRAZING

DO NOT graze treated forage within 7 days after application. DO NOT feed forage from treated areas to livestock within 7 days after application. DO NOT feed hay from treated areas to livestock within 30 days of application. Harvested straw may be used for bedding and/or feed. DO NOT harvest grain within 45 days after application.

CROP ROTATION TABLES

See the following tables for specific crop rotation guidance based on AGILITY SG herbicide use rate, location, soil pH, cumulative precipitation, and crop.

All Areas - Following Use of AGILITY SG herbicide at 1.6 to 3.2 Oz/A

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Winter wheat, spring wheat and Triticale	7.9 or lower	No restrictions	1
Durum wheat, barley, spring/winter oat	7.9 or lower	No restrictions	10

Rotation Intervals For Crops in Non-Irrigated Land Following Use of AGILITY SG Herbicide at 1.6 to 3.2 Oz/A on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Colorado	Statewide	Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Generally N of I-70	Field corn	7.9 or lower	15	12
	Statewide	BOLT® technology soybeans STS® Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
Idaho	Southern Idaho	Flax, Safflower	7.9 or lower	No restrictions	22
	Statewide	Peas, Lentils, Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
Kansas	Statewide	BOLT® technology soybeans STS® Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Central and Western Kansas (West of the Flint Hills)	Field corn	7.9 or lower	15	12
	Western Kansas W. of Hwy. 183	Soybeans	7.5 or lower	22	22
			7.6-7.9	33	34
	Central Kansas; generally E. of Hwy. 183 and W. of the Flintheads	Soybeans	7.9 or lower	15	12

Continued on next page

Rotation Intervals For Crops in Non-Irrigated Land (continued)

Following Use of AGILITY SG herbicide at 1.6 to 3.2 Oz/A on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Montana	Statewide	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22
		Alfalfa (hay only)	7.6-7.9	No restrictions	34
			7.5 or lower	No restrictions	22
		Flax, Safflower	7.9 or lower	No restrictions	22
Nebraska	Statewide	BOLT® technology soybeans STS® Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Generally W. of Hwy. 77 and E. of the Panhandle	Field corn	7.9 or lower	15	12
		Soybeans	7.5 or lower	22	22
			7.6-7.9	33	34
New Mexico	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower	7.9 or lower	No restrictions	22
	Eastern New Mexico	Cotton (dryland only)	7.9 or lower	30	22
North Dakota	W. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower	7.9 or lower	22	22
	E. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower	7.9 or lower	34	34
Oklahoma	Statewide	BOLT® technology soybeans STS® Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
	Panhandle	Cotton (dryland only)	7.9 or lower	30	22
		E. of the Panhandle	Cotton (dryland only)	7.9 or lower	25

Continued on next page

Rotation Intervals For Crops in Non-Irrigated Land (continued)

Following Use of AGILITY SG herbicide at 1.6 to 3.2 Oz/A on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Oregon	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
South Dakota	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22
	S. of Hwy. 212 & E. of the Missouri River, & S. of Hwy. 34 & W. of Missouri River	Grain sorghum, Proso millet	7.9 or lower	13	12
	Generally E. of Missouri River & S. of Hwy. 14, & W. of Missouri River	Field corn	7.9 or lower	15	12
Texas	Statewide	BOLT® technology soybeans STS® Soybeans	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Panhandle	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	30	22
	N. Central Texas*	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	25	14
	* The counties of N. Central Texas are: Archer, Baylor, Bell, Bosque, Bowie, Callahan, Camp, Cass, Clay, Collin, Cooke, Coryell, Dallas, Delta, Denton, Eastland, Ellis, Falls, Fannin, Foard, Franklin, Grayson, Hardeman, Haskell, Hill, Hood, Hopkins, Hunt, Jack, Johnson, Kaufman, Knox, Lamar, Limestone, McLennan, Milam, Montague, Morris, Nafarro, Palo Pinto, Parker, Rains, Red River, Robertson, Rockwall, Shackelford, Somervell, Stephens, Tarrant, Throckmorton, Titus, Upshur, Van Zandt, Wilbarger, Wichita, Williamson, Wise, Wood, Young.				
Utah	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22
Continued on next page					

Rotation Intervals For Crops in Non-Irrigated Land (continued)

Following Use of AGILITY SG herbicide at 1.6 to 3.2 Oz/A on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Washington	Statewide	Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
		Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
Wyoming	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22
	Southern Wyoming	Grain sorg Proso millet	7.9 or lower	No restrictions	10
	Southern Wyoming (Goshen, Laramie, and Platte counties only)	Field corn	7.9 or lower	15	12
	Northern Wyoming	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22

Rotation Intervals for crops not covered above - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

Rotation Intervals For Crops in Non-Irrigated Land

Following Use of AGILITY SG herbicide up to 2.4 Oz/A on Wheat, Barley, Triticale or Fallow in the states of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas and Wyoming

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Sunflower	7.9 or lower	No restrictions	10

Rotation Intervals for crops not covered above (up to 2.4 oz/A) - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

Rotation Intervals For Crops in Non-Irrigated Land

Following Use of AGILITY SG herbicide at 2.4 to 3.2 Oz/A on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Colorado Idaho Kansas Montana Nebraska New Mexico Oklahoma South Dakota Texas Utah Wyoming	Statewide	Sunflower	7.9 or lower	No restrictions	22
North Dakota	W. of Hwy. 1	Sunflower	7.9 or lower	22	22
	E. of Hwy. 1	Sunflower	7.9 or lower	34	34

Rotation Intervals for crops not covered above (2.4 to 3.2 oz/A) - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

Rotation Intervals For Crops in Non-Irrigated Land

Following Use of AGILITY SG herbicide at 1.6 Oz/A on Wheat, Barley Triticale or Fallow

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Sorghum, Grain	7.9 or lower	No restrictions	4
Cotton	7.9 or lower	No restrictions	10
Safflower	7.9 or lower	No restrictions	10
Peas, Dry /Green	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Lentils	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Alfalfa	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Beans, Dry	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Sunflower	7.9 or lower	No restrictions	10

Rotation Intervals for crops not covered above (1.6 oz/A) - The minimum rotation interval is 22 months with at least 18" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABES572.1).
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

Aerial Applications:

- DO NOT release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABES572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Boom-less Ground Applications

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Sensitive Areas

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

Drift Control Additives

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Council of Producers & Distributors of Agrotechnology (CPDA).

IDENTIFICATION INFORMATION FOR PRODUCTS REFERENCED IN THIS LABEL

REGISTERED PRODUCTS REFERENCED IN THIS LABEL FOR TANK MIXTURES OR MENTIONED FOR OTHER REASONS		
Product Name	Active Ingredient(s)	EPA Registration Number
Discover® NG Herbicide	Clodinafop-propargyl	100-1173
Buctril® Herbicide	Bromoxynil	264-437
Bronate® Herbicide	Bromoxynil + MCPA	264-438
Hoelon® 3EC Herbicide	Diclofop-methyl	264-641
Puma® IEC Herbicide	Fenoxaprop-p-ethyl	264-666
Bronate Advanced™ Herbicide	Bromoxynil + MCPA	264-690
Clarity® Herbicide	Dicamba	7969-137
Curtail® Herbicide	2,4-D + Clopyralid	62719-48
Stinger® Herbicide	Clopyralid	62719-73
Banvel® Herbicide	Dicamba	66330-276
Everest® 2.0 Herbicide	Flucarbazone-sodium	66330-391
Banvel® 480 Herbicide	Dicamba	66330-421
Everest® 3.0 Herbicide	Flucarbazone-sodium	66330-429
Everest® 3.0 AG	Flucarbazone-sodium	66330-433

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container only.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with AGILITY SG herbicide containing dicamba, thifensulfuron methyl, tribenuron methyl and metsulfuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with AGILITY SG herbicide containing dicamba, thifensulfuron methyl, tribenuron methyl and metsulfuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact FMC at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transportation and Spills) at 1-800-424-9300, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

FMC, Agility®, Ally®, Express®, Harmony®, and TotalSol® are trademarks of FMC Corporation or an affiliate

"Clarity" is a registered trademark of BASF Corporation

"Bronate", "Bronate Advanced", "Buctril", "Hoelon" and "Puma" are registered trademarks of Bayer CropScience

"Raindrop" is a registered trademark of Delavan

"Banvel" and "Everest" are registered trademarks of Arysta LifeScience North America LLC

"Discover" is a registered trademark of Syngenta Crop Protection LLC

"Stinger", "Curtail", and "Lorsban" are registered trademarks of Dow AgroSciences

"BOLT" and "STS" are registered trademarks of Corteva Agriscience United States

D - 4030 120919

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and, to the extent permitted by applicable law, Buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.