

SPECIMEN LABEL
SABADILLA DUST

CONTROL OF INSECTS LISTED ON:

**TOMATOES — TURNIPS — CABBAGE — CANTALOUPES
CUCUMBERS — SQUASH — WATERMELONS**

CONTAINS 20% POWDERED SABADILLA SEED

CONTENTS: 4 LB.

ACCEPTED
Jul 1, 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE ACT OF 1947 AND THE
FEDERAL FOOD, DRUG AND COSMETIC ACT
ED UNDER NO. 299-47. SUBJECT
TO ATTACHED COMMENTS.

CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS: Sabadilla Alkaloids 0.8%
INERT INGREDIENTS 99.2%
100.0%

E. Martin & Sons incorporated

DIRECTIONS

Sabadilla Dust can be used in any type of hand or power duster. Apply a light film of dust to all parts of plants, especially the underside of leaves (approximately 1/2 to 3/4 lb. per thousand square feet.)

Vegetable Plants: Tomatoes, Turnips, Cabbage, Cantaloupes, Cucumbers, Squash, Watermelons, Beans, Corn and Potatoes. For the control of Striped Cucumber Beetle, Squash Bug, Harlequin Bug, Lygus Bug, Leafhoppers, Mexican Bean Beetle, Potato Psyllid, Cabbage Worm, and Chinch Bug. Use 1/2 to 3/4 lbs. per thousand square feet.

For special information for uses in different locations which are consistent with label directions, consult your Agricultural Agent or State Experiment Station.

CAUTION

Sabadilla Dust may be irritating to the mucous membrane of the nose and throat. Avoid inhalation. A mask or cloth over nose and mouth may be used. Avoid contaminating food stuffs. Not for human use.

U.S.D.A. Reg. No. 299-47

Austin, Texas

**CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS.**

ACTIVE INGREDIENTS: Sebadilla Alkaloids
INERT INGREDIENTS

E. J. Martin Sons incorporated

SPECIMEN LABEL

DIRECTIONS

Sabadilla Dust can be used in any type of hand or power duster. Apply a light film of dust to all parts of plants, especially the underside of leaves (approximately 1/2 to 3/4 lb. per thousand square feet.)

Vegetable Plants: Tomatoes, Turnips, Cabbage, Cantaloupes, Cucumbers, Squash, Watermelons, Beans, Corn and Potatoes. For the control of Striped Cucumber Beetle, Squash Bug, Harlequin Bug, Lygus Bug, Leafhoppers, Mexican Bean Beetle, Potato Psyllid, Cabbage Worm, and Chinch Bug. Use 1/2 to 3/4 lbs. per thousand square feet.

For special information for uses in different locations which are consistent with label directions, consult your Agricultural Agent or State Experiment Station.

CAUTION

Sabadilla Dust may be irritating to the mucous membrane of the nose and throat. Avoid inhalation. A mask or cloth over nose and mouth may be used. Avoid contaminating food stuffs. Not for human use.

U.S.D.A. Reg. No. 299-47

Austin, Texas