

ACCEPTED

299-77

BACK RUBBING OIL: 1/2 gallon SUPER STOCK TOX mixed with 9 1/2 gallons kerosene or diesel fuel oil makes 10 gallons Back Rubbing Oil. To aid in controlling Lice and Hornflies on Beef Cattle, charge burlap roll with solution so it is wet but not to the point of dripping. Repeat charging as necessary. Do not permit animals access to back rubbers within 30 days of slaughter, nor to cattle under 6 months of age.

CAUTION: SUPER STOCK TOX forms a stable emulsion with water, but some settling may occur, especially in hard water. If the mixture is allowed to set for a period of time, it should be thoroughly agitated before starting to spray.

DO NOT USE ON DOGS, CATS OR FOWL.

Do not treat animals under three months of age. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not use on dairy goats. Do not apply to animals within 60 days of slaughter by dipping and within 30 days by spraying. Avoid contamination of food and feed stuff.

WARNING: SUPER STOCK TOX is toxic and may be absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid inhaling spray mist. At first signs of nasal irritation, stop spraying and seek fresh air. In case of skin irritation, wash surface with soap and water.

Toxic to fish and wildlife, do not contaminate streams, lakes or ponds. Destroy empty containers, spillage and spray tank washings. Do not contaminate water by cleaning of equipment, or disposal of wastes.

May be fatal if swallowed. **ANTIDOTE:** One tablespoon of salt in a glass of warm water to induce vomiting. Repeat until vomit is clear.

U.S.D.A. Reg. No. 299-77



SUPER STOCK TOX

New Improved Formula Containing Lindane. For use as a Livestock Spray to control Lice, Ticks, and Hornflies.

AN EMULSIFIABLE CONCENTRATE

Containing Toxaphene and the Gamma Isomer of BHC which in pure state is called Lindane.

CONTENTS: ONE-HALF GALLON LIQUID

WARNING: Keep Out of Reach of Children.

SEE BACK PANEL FOR ADDITIONAL WARNINGS

ACTIVE INGREDIENTS:

*Toxaphene	62.0%
Gamma Isomer of Benzene Hexachloride from Lindane	2.5%
Petroleum Solvents	29.5%
INERT INGREDIENTS:	6.0%
	<hr/>
	100.0%

*Technical Chlorinated Camphene containing 67-69% Chlorine.

CATTLE: F Gulf Coast L Lice—mix 1/2 Thoroughly S a mist spraye and neck. D necessary.

SHEEP / Sheep Ticks 1/2 gallon SU thoroughly d Scabies—mix thoroughly. C on Dairy Goa

HOGS— SUPER STOC One applicat animals thorc 1/2 gallon SU sows within 2 after.

VATS—S for Lice on S per 800 gal STOCK TOX stirring. Stir longed stanc cause breakd tained. vat s

Austin, Texas

E. J. Martin & Sons incorporated

ACCEPTED
FEDERAL INSECTICIDE
TOXICITY TESTS
DO NOT MIX WITH
OTHER TOXIC
SUBSTANCES
DO NOT USE
ON WET PLANTS
NECESSARY. DO NOT
USE FOR 30 DAYS OF

299-77



SUPER STOCK TOX

New Improved Formula Containing Lindane. For use as a Livestock Spray to control Lice, Ticks, and Hornflies.

AN EMULSIFIABLE CONCENTRATE

Containing Toxaphene and the Gamma Isomer of BHC which in pure state is called Lindane.

CONTENTS: ONE-HALF GALLON LIQUID

WARNING: Keep Out of Reach of Children.

SEE BACK PANEL FOR ADDITIONAL WARNINGS

ACTIVE INGREDIENTS:	
*Toxaphene	62.0%
Gamma Isomer of Benzene Hexachloride from Lindane	2.5%
Petroleum Solvents	29.5%
INERT INGREDIENTS:	
	6.0%
	<hr/>
	100.0%

*Technical Chlorinated Camphene containing 67-69% Chlorine.

DIRECTIONS:

CATTLE: For the control of Lone Star, Cattle, Winter and Gulf Coast Ticks, Hornflies, and Long and Short Nose Cattle Lice—mix 1/2 gallon SUPER STOCK TOX with 100 gallons water. Thoroughly Spray all parts of animal's body. For Ear Ticks use a mist sprayer and apply spray into each ear and to the head and neck. Do not use on Dairy Cows. Repeat application as necessary.

SHEEP AND GOATS—For the control of Fleece Worms, Sheep Ticks (keds), Sheep Lice, Red and Yellow Goat Lice—mix 1/2 gallon SUPER STOCK TOX with 100 gallons water and spray thoroughly during warm weather when wool is short. For Sheep Scabies—mix 1/2 gallon with 70 gallons water and dip sheep thoroughly. One application is generally sufficient. Do not use on Dairy Goats.

HOGS—For the control of lice on hogs, mix 1/2 gallon SUPER STOCK TOX with 100 gallons water and spray thoroughly. One application is usually sufficient. For Sarcoptic Mange, spray animals thoroughly twice at 14 day intervals with a mixture of 1/2 gallon SUPER STOCK TOX in 70 gallons water. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

VATS—SUPER STOCK TOX may be used in vats as a dip for Lice on Sheep & Goats. Mix 1/2 gallon SUPER STOCK TOX per 800 gallons of water. Pour required amount of SUPER STOCK TOX in vat and add water slowly at first with vigorous stirring. Stir contents of vat thoroughly before use. After prolonged standing, fermentation of organic matter in vat may cause breakdown of the emulsion. If good mixture is not obtained, vat should be emptied, cleaned and recharged.

E. J. Martin & Sons incorporated

Austin, Texas