

[ 299-105 ]



# 5% MALATHION DUST

For use on Beef and Dairy Cattle, Hogs, Sheep, Goats, Poultry, also Garden and Ornamental Plants.

JAN 22, 1970

NET CONTENTS: 4 POUNDS

**CAUTION: Keep Out of Reach of Children.**  
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS  
INERT INGREDIENTS

5%  
95%

\*100% methyl dithiophosphate or diethyl malathion concentrate

U.S.P.A. Reg. No. 299-105 0765



**DIRECTIONS FOR USE ON BEEF CATTLE** For the control of horn flies, take a handful (4 tablespoons) and dust over an animal's back and neck. Will protect a cow from hornflies up to 14 days. Do not apply to calves under one month of age.

**DAIRY CATTLE** For control of hornflies on lactating dairy cattle, apply 1 1/2 ounces per head to back and neck of animals. Do not apply within 5 hours of milking or during milking. Repeat application at 10-14 day intervals if needed. For non-lactating dairy cattle, apply dust thoroughly to back and neck of animal. Do not treat non-lactating dairy cattle within 2 weeks of freshening. Repeat application at 10-14 day intervals if needed.

**SHEEP AND GOATS** For the control of ticks, sheep ticks (keds), and lice, use 2 to 4 tablespoons per animal depending on size of animal and amount of hair. Thoroughly cover the entire body of the animal. Do not apply to milk goats. Repeat in two weeks if necessary.

**HOGS** For control of lice on hogs, work 2 to 4 tablespoons of MALATHION DUST into the hair. Thoroughly cover the entire body of the animal. Repeat in two weeks if necessary.

**POULTRY** To control lice on poultry, mix 2 pounds of Malathion 5% Dust with 100 pounds of feed to make a 4% mixture. Then use 2 lbs. per 100 to 150 square feet of floor space, or apply pinches directly to the birds.

### DIRECTIONS FOR USE ON GARDEN AND ORNAMENTAL PLANTS

MARTIN'S 5% MALATHION DUST may be used on Ornamentals, shrubs, and shade trees for control of Aphids, Japanese Beetles, adults, Lace Bugs, Leafhoppers, Mealy Bugs, Fawn-spined Plant Bugs, Thrips, Whitefly, Bagworms, and Spider Mites. MARTIN'S 5% MALATHION DUST can be used on Tomatoes, Peas, Beans, Cucumbers, Squash, Melon, Onions, and Potatoes for the control of Aphids, Spider Mites, Leafhoppers, Mexican Bean Beetle and other insects. Do not apply later than 3 days before harvest on the above vegetables.

**CAUTION** Do not use on calves and other young animals less than one month of age. Avoid contamination of feed, foodstuffs, and watering troughs. Destroy empty containers and waste.

This product contains Malathion which is a cholinesterase inhibitor. Do not use this product simultaneously with other drugs, insecticides, pesticides or chemicals having cholinesterase inhibiting activity nor within a few days before or after treatment with Malathion.

Keep out of any body of water. This product is toxic to fish.

**WARNING** Avoid contact with skin. Avoid breathing of dust. Wash thoroughly.

**KEEP OUT OF REACH OF CHILDREN**

204 W. MAIN ST. NAACOGDOCHES, TEX. 75941

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

**SEE BACK PANEL FOR OTHER CAUTIONS**

ACTIVE INGREDIENT

MALATHION

INERT INGREDIENTS

500 grams (17.6 ounces) per 100 grams (3.5 ounces)

U.S.D.A. Reg. No. 299-105-0765

*C. J. Martin Company*

**DIRECTIONS FOR USE ON BEEF CATTLE:** To control horn flies, take a handful (4 tablespoons) and dust over the back of a cow from horn flies up to 4 days. Do not apply to calves less than one month of age.

**DAIRY CATTLE:** For control of horn flies, apply dust to the back, neck, and head of animals. Do not apply within 24 hours of milking or during milking. Repeat application at 10-14 day intervals if needed. For non-lactating dairy cattle, apply dust thoroughly to back and neck of animal. Do not treat non-lactating dairy cattle within 2 weeks of freshening. Repeat application at 10-14 day intervals if needed.

**SHEEP AND GOATS:** For the control of ticks, sheep ticks (keds) and lice, use 2 to 4 tablespoons per animal depending on size of animal and amount of hair. Thoroughly cover the entire body of the animal. Do not apply to milk goats. Repeat in two weeks if necessary.

**HOGS:** For control of lice on hogs, work 2 to 4 tablespoons of MALATHION DUST into the hair. Thoroughly cover the entire body of the animal. Repeat in two weeks if necessary.

**POULTRY:** To control lice on poultry, mix 2 pounds of Malathion 5% Dust with 1/2 pound talc to make a 4% mixture. Then use 2 lbs. per 100 to 120 square feet of floor space, or apply pinches directly to the birds.

**DIRECTIONS FOR USE ON GARDEN AND ORNAMENTAL PLANTS:**

MARTIN'S 5% MALATHION DUST may be used on Ornamentals, shrubs, and Shade trees for control of Aphids, Japanese Beetles (adults), Lace Bugs, Leafhoppers, Mealy Bugs, Tarnished Plant Bugs, Thrips, Whitefly, Bagworms, and Spider Mites. MARTIN'S 5% MALATHION DUST can be used on Tomatoes, Peas, Beans, Cucumbers, Squash, Melon, Onions, and Potatoes for the control of Aphids, Spider Mites, Leafhoppers, Mexican Bean Beetle and Onion Thrips. Do not apply later than 3 days before harvest on the above vegetables.

**CAUTION:** Do not use on calves and other young animals less than one month of age. Avoid contamination of feed, foodstuffs, and watering troughs. Destroy empty containers and waste.

This product contains Malathion which is a cholinesterase inhibitor. Do not use this product simultaneously with other drugs, insecticides, pesticides or chemicals having cholinesterase inhibiting activity nor within a few days before or after treatment with Malathion.

Keep out of any body of water. This product is toxic to fish.

**WARNING:** Avoid contact with skin. Avoid breathing of dust. Wash thoroughly.

**KEEP OUT OF REACH OF CHILDREN.**

606 W. MAIN ST.

NAZOGDOCHES, TEX. 75962