219	-	//	0
-----	---	----	---

MART'IN'S

50% MALATHION CONCENTRATE

For use on Dogs, Cats, Household pests, **xrtash** vegtables and ornamentals.

04/27/1967

ACCEPTED

APR 2 7 1967

UNDER THE FEDERAL INSECTICIDE FUNCICIDE AND RODENTICIDE ACT FOR SCONOMIC POISON REGISTER-ED UNDER NO. 249-110. SUBJECT

6%

TO ATTACHED COMMENTS.

CONTENTS:

CAUTION: KEEP OUR OR REACH OF CHILDREN. See back panel for other cautions.

Malathion* . . . ACTIVE INGREDIENTS: 50% 44% Aromatic Petroleum Derivative

INERT INGREDIENTS:

*0,0-dimethyl dithiophosphate of diethylmercaptosuccinate DIRECTIONS: ORNAMENTALS: Can be used on Roses, Camellias, Azaleas, Chrysanthemums, Evergreens and many other flowers and shrubs for the control of Aphids, Birch Leaf Minor, Japanese Beetles (Adult), Lace Bugs, Leafhoppers, Mealy Bugs, Tarnished Plant Bugs, Thrips, and Whiteflies; use 1 teaspoonful per gallon of water. For the control of Bagworms, Ten caterpillars, Black Scale Crawlers, Leaf Scale, Oyster Shell Scale, Soft Brown Scale and Spider Mites; use 2 teaspoonfuls per gallon of water. Do not use on certain ferns including Boston, Maidenhair and Pteris, as well as on some species of Crassula.

Can be used on Tomatoes, Potatoes, Peas, Beans, Peppers, **VEOTABLES:** -Cabbage, Broccoli, Cucumbers, Squash and Melons for the control of Aphids, Spider Mites, Leafhoppers, Mexican Bean Beetles and Imported Cabbage Worms, use l teaspoonful per gallon of water. For the control of-Onion Thrips on onions use 2 teaspoonfuls per gallon of water. DO NOT USE later than 3 day before harvest on the above vegetables with the exception of Cabbage and this should not be applied later than 7 days before harvest.

HOUSEHOLD PESTS: For the control of Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites and Earwigs, use 8 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to 1 gallon of water. Apply by means of coarse spray or paint brush to window sills, base boards, drainboards, under sinks, stoves and to other areas frequented by insects.

DOGS AND CATS: For the control of Fleas, use $2\frac{1}{4}$ tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to each gallon of water. For control of Fleas in the animal's quarters, use 12 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE ato each gallon of water. Repeat applications, if necessary, should not be more often than once a week.

FLY CONTROL: For the control of Flies, mix $2\frac{1}{4}$ quarts per 100 gallons of water. Spray on walls, fences, garbage containers and wherever flies congregate and breed. Wet all surfaces thoroughly. The addition of 8 tablespoonfuls of sugar per gallon will increase the effectiveness of the spray.

MARTIN'S MALATHION CONCENTRATE 50% (cont.)

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or prepeated contact with skin. If spilled on skin, remove contaminated clothing and wash the contaminated skin areas thoroughly with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals

NOTE TO PHYSICIANS: Malathion, upon repeated or prolonged careless use, may cause cholinesterase inhibition. Atropine is an antiodote. Keep container closed and vaway from heat and open flame.

NOTICE: SELLER makes no warranty, expressed or implied, concerning the use, storage, or handling of this product, except that it conforms to the chemical description on the label. USER assumes all risks of using or handling, whether in accordance with directions or not.

U.S.D.A. REG. NO. 299-110

MANUFACTURED BY C. J. MARTIN & SONS, INC., AUSTIN, TEXAS