

ACCEPTED

Jul 25, 1968

UNDER THE FEDERAL INSECTICIDE

water. For the control of Bagworms, Tent Caterpillars, Pear Sawflies, Crows, Leaf Scale, Oyster Shell Scale, Soft Scale, Spider Mites; use 2 teaspoonfuls per gallon of water. Do not use on certain ferns, including Boston, Maidenhair and others, as on some species of Crassula.

VEGETABLES: Can be used on Tomatoes, Potatoes, Peas, Beans, Peppers, Cabbage, Broccoli, Cucumbers, Squash and Melons for the control of Aphids, Spider Mites, Leafhoppers, Mexican Bean Beetles and Imported Cabbage Worms; use 1 teaspoonful per gallon of water. For the control of onions, Thrips on onions, use 2 teaspoonfuls per gallon of water. DO NOT USE later than 3 days before harvest on the above vegetables with the exception of Cabbage and this should not be applied later than 7 days before harvest.

HOUSEHOLD PESTS: For the control of Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites and Earwigs, use 8 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to 1 gallon of water. Apply by means of coarse spray or paint brush to window sills, base boards, beneath drainboards, under sinks, stoves and to other areas infested by insects.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. If spilled on the skin, remove contaminated clothing and wash the contaminated skin areas thoroughly with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals. Do not reuse empty containers. Wash thoroughly with water and detergent. Discard in a safe place.

This product contains "Malathion" which is a cholinesterase inhibitor. Do not use this product simultaneously with other drugs, insecticides, pesticide or chemicals having cholinesterase inhibiting activity nor within a few days before or after treatment with "Malathion."

NOTE TO PHYSICIANS: Malathion upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is an antidote. Keep container closed and away from heat and open flame.

This product is toxic to fish. Keep out of lakes, streams, and ponds.

U.S.D.A. Reg. No. 299 110



SPECIAL LABEL

50% MALATHION CONCENTRATE

For use on Beef Cattle, Sheep, Goats, Hogs, Dogs, Cats, Household Pests, Vegetables, Trees and Ornamentals.

Kills: Lice, Ticks, Fleas, Mites, Flies, Mosquitoes, Roaches, Ants and Spiders

CONTENTS:

CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:	
Malathion	50%
Aromatic Petroleum Derivative Solvent	45%
INERT INGREDIENTS:	
	100%

*0.0 d methyl dithiophosphate of diethyl mercaptosuccinate

DIRECTIONS

BEEF CATTLE: For the control of Lice, Ticks and Horn Flies, use 1 quart of Martin's 50% MALATHION CONCENTRATE to 20 gallons of water. Spray animal thoroughly. Repeat at 2 week intervals if necessary. Do not apply to lactating dairy animals.

SHEEP AND GOATS: For the control of Lice, Ticks, and Keds, use 1 quart of Martin's 50% MALATHION CONCENTRATE to 20 gallons of water. Spray animals thoroughly. Repeat application after 2 or 3 weeks if necessary.

HOGS: For the control of Lice, use 1 quart of Martin's 50% MALATHION CONCENTRATE to 20 gallons of water. Apply thoroughly to the animals, pens and litter. Repeat only if necessary.

DOGS AND CATS: For the control of Fleas, use 2 1/4 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to each gallon of water. For control of Fleas in the animal's quarters, use 12 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to each gallon of water. Repeat application, if necessary, should not be more than once a week.

FLY CONTROL: For the control of Flies, mix 2 1/4 quarts per 100 gallons of water. Spray on walls, fences, garbage containers and wherever flies congregate and breed. Wet all surfaces thoroughly. The addition of 8 tablespoonfuls of sugar per gallon will extend the life of the bait residue.

FRUITS: (Apples, Peaches, Pears and Plums) For the control of Spider Mites, Oriental Fruit Moth and Plum Curculio. For the control of Red Spider Mites, Mealy Bugs, Woolly Apherid, Pear Psyllid and Japanese Beetle (Adults); use 1/2 to 2 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to 1 gallon of water. Do not make application on fruits 7 days before harvest.

PECAN TREES: For the control of Pecan Nut Casebearer, Pecan Phylloxera, Pecan Budmoth, Spider Mites and Aphids, use 2 tablespoonfuls of Martin's 50% MALATHION CONCENTRATE to 1 gallon of water.

ORNAMENTALS: Can be used on Roses, Camellias, Azaleas, Chrysanthemums, Evergreens and many other flowers and shrubs for the control of Aphids, Birch Leaf Miner, Japanese Beetles (Adult), Lace Bugs, Leafhoppers, Mealy Bugs, Tarnished Plant Bugs, Thrips, and Whiteflies; use 1 teaspoonful per gallon of

Austin, Texas

E. J. Martin & Sons incorporated

Austin, Texas

299-110

07/25/1968

111