



MARTIN'S

10% SEVIN DUST (R)

Ready to use on:

FLOWERS - VEGETABLES - GARDENS - LAWNS - SHRUBS

CAUTION: Keep out of reach of children

Contents: 2 Pounds

(R) Trademark of Union Carbide Corporation for 1-Naphthyl Methylcarbamate

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl Methylcarbamate).....10.00%

INACTIVE INGREDIENTS:.....90.00%

TOTAL..... 100.00%

DIRECTIONS:

APPLICATION: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES

Corn, Eggplant, Pepper, Tomato: To control Corn earworm, European corn borer, sap beetles, fall army worm, tomato fruitworm, flea beetles, leafhopper, tomato hornworm and stink bugs--apply at the rate of 1 to 2 ounces to 50 feet of row.

Beans: to control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer-- apply at the rate of 1 to 2 ounces per 50 feet of row.

Cucumber, Summer Squash: to control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug, and Flea Beetles--apply at the rate of 1 ounce to 50 feet of row.

Potato: To control Colorado Potato Beetle, Leafhoppers, Flea Beetle, and Lygus Bug -- apply at the rate of 1 to 2 ounces to 50 feet of row.

Applications to above vegetables can be made up to day of harvest.

MARTIN'S 10% Sevin Dust - (cont'd)

LAWNS

To control Leafhoppers, Fall Army Worms, and Sod Webworms-- apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks of necessary.

ORNAMENTALS

To control Apple Aphids, Bagworms, Birch Leaf Miners, Blister Beetles, Box Elder Bugs, Boxwood Leaf Miners, Elm Leaf Aphids, Elm Leaf Beetles, Flea Beetles, Japanese Beetles, Lace Bugs, Leafhoppers, Leaf Rollers, Oak Leaf Miners, Orange Tartrix, Periodical Cicada, Plant Bugs, Psyllids, Roseslugs, Rose Aphids, Scale insects in the crawler stage, Tent Caterpillars, Willow Leaf Beetles, and exposed Thrips--apply thoroughly to the infested plants at the first sign of damage and repeat at 7-day intervals or as necessary for good insect control. To control scale, apply in spring and early summer when scale crawlers are present.

DOGS, CATS

To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens less than four weeks old.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. ~~XXXXXXXXXXXXXXXXXXXXXXXXXXXX~~

WARRANTY

NOTICE: C.J. Martin & Sons, Inc. makes no warranty, expressed or implied, concerning this material, except that it conforms to the chemical description on the label. User assumes all risks of using or handling, whether in accordance with directions or not.