



SPECIAL FORMULATION

10% SEVIN[®] DUST

READY TO USE ON:
**VEGETABLES • GARDENS • LAWNS
POULTRY • DOGS • CATS**

ACCEPTED
Jul 1, 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER 299-117
SUBJECT TO ATTACK

CAUTION: Keep Out of Reach of Children.
SEE BACK PANEL FOR OTHER CAUTIONS.

CONTENTS: 4 POUNDS

[®] Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.
ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate)10%
INERT INGREDIENTS:90%
100%

E. I. Martin & Sons incorporated

DIRECTIONS

APPLICATION: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES

CORN, EGGPLANT, PEPPER, TOMATO: To control Corn Earworm, European Corn Borer, Sap Beetles, Fall Army Worm, Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm and Stink Bugs—apply at the rate of 1 ounce per 50 feet of row. For better control of Corn Earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Reapply immediately after rain.

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer—apply at the rate of 1 ounce per 50 feet of row.

CUCUMBER, SUMMER SQUASH: To control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug and Flea Beetles—apply at the rate of 1 ounce to 50 feet of row. Apply to Cucumber and Summer Squash only when plants are dry.

POTATO: To control Colorado Potato Beetles, Leafhoppers, Flea Beetles, and Lygus Bugs—apply at the rate of 1 ounce to 50 feet of row.

APPLICATIONS TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST

LAWNS

To control Leafhoppers, Fall Army Worms and Sod Webworms—apply at the rate of 1/2 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY

Litter Treatment—Apply 1/2 pound to 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas, and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

HAND APPLICATION TO BIRDS

Use 1/2 pound to treat 100 birds by means of shaker can, or hand duster to control Northern Fowl Mite, Lice and Fleas. Direct dust to vent and fluff areas. Do not contaminate feed or drinking water. The application of this material can not be repeated within 4 weeks of each treatment. Hand application to birds is a supplement to litter treatment for control of these pests. Do not apply to poultry within 7 days of slaughter.

DOGS, CATS

To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Do not treat kittens less than four weeks old. Do not use on nursing puppies. Do not get powder in eyes and mouth of animals during treatment. Do not repeat within 1 week of each treatment.

CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. Reg. No. 299-117

Austin, Texas