

299-117

05/20/1971

1/2

M

MARTIN'S

10% SEVIN[®]
DUST

READY TO USE ON:

VEGETABLES • GARDENS • LAWNS
POULTRY • DOGS • CATS

CAUTION: Keep Out of Reach of Children.
SEE BACK PANEL FOR OTHER CAUTIONS.

NET WEIGHT: 4 POUNDS

® Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) ----- 10%

INERT INGREDIENTS: ----- 90%

100%

606 W. Main Street

C. J. Martin Company

Nacogdoches, Texas 75961

ACCEPTED
MAY 20 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 299-117. SUBJECT
TO ATTACHED COMMENTS.

4-11-51
BB2-bb

DIRECTIONS

APPLICATION: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES

CORN, EGGPLANT, PEPPER, TOMATO: To control Corn Earworms, European Corn Borers, Sap Beetles, Armyworms, Tomato Fruitworms, Flea Beetles, Leafhoppers, Tomato Hornworms, and Stink Bugs---dust the upper and lower leaf surfaces thoroughly. For better control of Corn Earworms, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Reapply immediately after rain.

BEANS: To control Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworms, Lygus Bugs, and Lima Bean Pod Borers---dust the upper and lower leaf surfaces thoroughly.

CUCUMBER, SUMMER SQUASH: To control Pickleworms, Melonworms, Cucumber Beetles, Squash Bugs, and Flea Beetles---dust the upper and lower leaf surfaces thoroughly. Apply to Cucumber and Summer Squash only when plants are dry.

POTATO: To control Colorado Potato Beetles, Leafhoppers, Flea Beetles, and Lygus Bugs---dust the upper and lower leaf surfaces thoroughly.

APPLICATIONS TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST

LAWNS

To control Chinch Bugs, Armyworms, Ants, Fleas, Sod Webworms, Lawn Moths, Millipedes, and Earwigs---apply 1/2 pound to 200 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY

Litter Treatment---apply 1/2 pound to 40 square feet to control Northern Fowl Mites, Chicken Mites, Lice, Fleas, and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

HAND APPLICATION TO BIRDS

Use 1/2 pound to treat 100 birds by means of shaker can, or hand duster to control Northern Fowl Mites, Lice, and Fleas. Direct dust to vent and fluff areas. Do not contaminate feed or drinking water. The application of this material can not be repeated within 4 weeks of each treatment. Hand application to birds is a supplement to litter treatment for control of these pests. Do not apply to poultry within 7 days of slaughter.

DOGS, CATS

To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes, and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Do not treat kittens less than four weeks old. Do not use on nursing puppies. Do not get powder in eyes or mouth of animals during treatment. Do not apply within 1 week of each treatment.

CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

