

DIRECTIONS FOR USE
(Continued)

VEGETABLES:

Corn, Eggplant, Pepper, Tomato: To control Corn Earworms, European Corn Borers, Sap Beetles, Armyworms, Tomato Fruitworms, Flea Beetles, Leafhoppers, Tomato Hornworms, and Stink Bugs - dust the upper and lower leaf surfaces thoroughly. For better control of Corn Earworms, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals. Reapply immediately after rain.

Beans: To control Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworms, Lygus Bugs, and Lima Bean Pod Borers - dust the upper and lower leaf surfaces thoroughly.

Cucumber, Summer Squash: To control Pickleworms, Melonworms, Cucumber Beetles, Squash Bugs, and Flea Beetles - dust the upper and lower leaf surfaces thoroughly. Apply to cucumber and summer squash only when plants are dry.

Potato: To control Colorado Potato Beetles, Leafhoppers, Flea Beetles, and Lygus Bugs - dust the upper and lower leaf surfaces thoroughly.

APPLICATION TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST. SUBSEQUENT PLANTING OF FOOD AND FEED CROPS IN TREATED SOIL MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABELS.

LAWNS: To control Chinch Bugs, Armyworms, Anles, Fleas, Sod Webworms, Lawn Moths, Millipedes, and Earwigs - apply 1/2 pound to 200 sq. ft. For best results apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY: Litter Treatment - apply 1/2 pound to 40 sq. ft. to control Northern Fowl Mites, Chicken Mites, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nest eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

Hand Application To Birds: Use 1/2 pound to treat 100 birds by means of shaker can, or hand duster to control Northern Fowl Mites, Lice, and Fleas. Direct dust to vent and fluff areas. Do not contaminate feed or drinking water. The application of this material can not be repeated within 4 weeks of each treatment. Hand application to birds is a supplement to litter treatment for control of these pests. Do not apply to poultry within 7 days of slaughter.

DOGS, CATS: To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes, and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Do not treat kittens less than four weeks old. Do not use on nursing puppies. Do not get powder in eyes or mouth of animal during treatment. Do not apply within 1 week of each treatment. Do not use on pregnant dogs.

E.P.A. Reg. No. 299-117
E.P.A. Est. No. 299-TX-1




10% SEVIN® DUST

READY TO USE ON:

VEGETABLES • GARDENS • LAWNS
POULTRY • DOGS • CATS

NET WEIGHT:
1 POUND



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel For Other Cautions

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 10%

INERT INGREDIENTS: 90%

©Registered Trademark of Union Carbide for 1-naphthyl N-methylcarbamate

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTE TO PHYSICIAN

Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dust has settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

STORAGE & DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry place until used. Store out of reach of children, pets and animals.

DISPOSAL: Do not reuse empty container. Completely empty can then dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

APPLICATION: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

487121

606 W. Main St.

Cy Martin Company

Nacogdoches, Texas 75963-1089

299-117
10/17/86