

MARTIN'S

5% SEVIN DUST

Ready to use on : Poultry, Dogs, Cats, Vegetables, Flowers, Gardens, Lawns, Shrubs

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back panel for other cautions.

M

CONTENTS

(R) Trademark of Union Carbide Corporation for 1-naphthyl-N-methylcarbamate

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl-N-methylcarbamate) 5%

INERT INGREDIENTS: 95%
100%

DIRECTIONS: Application: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES - Corn, Eggplant, Pepper, Tomato: To control Corn earworm, European corn borer, sap beetles, fall army worm, tomato fruitworm, flea beetles, leafhopper, tomato hornworm and stink bugs - apply at the rate of 1 to 2 ounces per 50 feet of row.

BEAN: to control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer - apply at the rate of 1 to 2 ounces per 50 feet of row.

CUCUMBER, SUMMERSQUASH: to control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug and Flea Beetles - apply at the rate of 1 ounce to 50 feet of row.

POTATO: to control Colorado Potato Beetle, Leafhoppers, Flea Beetles, and Lygus Bugs - apply at the rate of 1 to 2 ounces to 50 feet of row. Applications to above vegetables can be made up to day of harvest.

LAWNS: To control Leafhoppers, Fall Army Worms, and Sod Webworms - apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY - Litter Treatment - Apply 1 pound to 40 square feet to control northern fowl mite, chicken mite, lice, fleas and bed bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

HAND APPLICATION TO BIRDS: Use 1 pound to treat 100 birds by means of a shaker can, or hand duster to control northern fowl mite, lice and fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests. Do not apply within 7 days of slaughter.

DOGS, CATS: To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens less than four weeks old.

MARTIN'S 5% SEVIN DUST (cont.)

CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

WARRANTY - NOTICE: C. J. MARTIN & SONS, INC. make no warranty, expressed or implied concerning this material, except that it conforms to the chemical description on the label. USER assumes all risks of using or handling, whether in accordance with directions or not.

U.S.D.A. REG. NO. 299-114

