

299-121

04/03/69

1/2

MARTIN'S
5% SEVIN (R) DUST

Ready to use on: Beef Cattle-Horses-Sheep-Swine-
Poultry-Dogs-Cats-Vegetables-Gardens-Lawns

CONTENTS:

CAUTION: Keep Out of Reach of Children. See Back Panel For
Other Cautions.

(R) Trademark of Union Carbide Corporation for 1-naphthyl
N-methylcarbamate.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate).....	5.00%
INERT INGREDIENTS.....	95.00%
TOTAL...	<u>100.00%</u>

DIRECTIONS:

Begin application when insect damage is first observed and
repeat on a regular 7 to 10 day schedule or as necessary
for good insect control.

VEGETABLES--Corn, Eggplant, Pepper, Tomato: To control Corn
Earworm, European Corn Borer, Sap Beetles, Fall Army Worm,
Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm
and Stink Bugs--apply at the rate of 1 to 2 ounces per 50
feet of row.

BEAN--To control Mexican Bean Beetle, Bean Leaf Beetle,
Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles,
Corn Earworm, Lygus Bug and Lima Bean Pod Borer--apply at
the rate of 1 to 2 ounces per 50 feet of row.

CUCUMBER, SUMMER SQUASH--To control Pickleworm, Melonworm,
Cucumber Beetles, Squash Bug and Flea Beetles--apply at the
rate of 1 ounce to 50 feet of row.


POTATO--To control Colorado Potato Beetle, Leafhoppers, Flea
Beetles, and Lygus Bugs--apply at the rate of 1 to 2 ounces
to 50 feet of row. Applications to above vegetables can be
made up to 1 day of harvest

LAWNS--To control Leafhoppers, Fall Army Worms, and Sod
Web-worms--apply at the rate of 1 pound to 250 square feet.
For best results, apply after rain or watering and do not
water for at least two days after application. Repeat in
two or three weeks if necessary.

ACCEPTED

APR 3 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
UNDER NO. 299-121 SUBJECT
TO ATTACHED COMMENTS.



SEVIN 5% DUST
Cont'd.....

POULTRY--Litter Treatment--Apply 1 pound to 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within seven days of slaughter.

DOGS, CATS--To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

LIVESTOCK: For the control of horn flies and stable flies on Beef Cattle, Horses, Sheep and Swine, rub 2 ounces of powder per animal into the hair on back and neck. Repeat if necessary, but not more often than once every 4 days. Do not apply within one week of slaughter and do not exceed dosages as given above. Do not use on dairy animals.

CAUTION-HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible Cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. REG. NO. 299-121

MANUFACTURED BY C.J. MARTIN & SONS, INC., AUSTIN, TEXAS - 78702

orm
2-20-69

