



5% SEVIN DUST

Ready to use on: Beef Cattle, Horses, Sheep,
Swine, Poultry, Dogs, Cats,
Vegetables, Gardens, Lawns

CONTENTS: 4 POUNDS

CAUTION: Keep Out of Reach of Children

SEE BACK PANEL FOR OTHER CAUTIONS

® Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS:	95%
	100%

E. I. Martin & Sons incorporated

DIRECTIONS: Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

VEGETABLES--Corn, Eggplant, Pepper, Tomato: To control Corn Earworm, European Corn Borer, Sap Beetles, Fall Army Worm, Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm and Stink Bugs -- apply at the rate of 1 to 2 ounces per 50 feet of row. When used on sweet corn for control of corn earworms, application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

BEAN--To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer -- apply at the rate of 1 to 2 ounces per 50 feet of row.

CUCUMBER, SUMMER SQUASH--To control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug, and Flea Beetles -- apply at the rate of 1 ounce to 50 feet of row.

POTATO--To control Colorado Potato Beetle, Leafhoppers, Flea Beetles, and Lygus Bugs -- apply at the rate of 1 to 2 ounces to 50 feet of row. Applications to above vegetables can be made up to day of harvest.

LAWNS--To control Leafhoppers, Fall Army Worms, and Sod Web-worms -- apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY--Litter Treatment -- Apply 1 pound to 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within seven days of slaughter.

DOGS, CATS--To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

LIVESTOCK: For the control of hornflies and suppression of stable flies on Beef Cattle, Horses, Sheep and Swine, rub 2 ounces of powder per animal into the hair on back and neck. Stable fly treatment of legs of animals is supplemental to premise treatment for stable fly control. Repeat if necessary, but not more often than once every 4 days. Do not apply within one week of slaughter and do not exceed dosages as given above. Do not use on dairy animals.

CAUTION--HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible Cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. REG. NO. 299-121

96124

Austin, Texas

ACCEPTED

AUG 19 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
UNDER NO. 299-121

