

**POULTRY:** Litter Treatment -- Apply 1 pound to 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

**DOGS, CATS:** To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

**HORSES:** For the control of horn-flies and suppression of stable flies, rub 2 ounces of powder per animal into the hair on back and neck. Stable fly treatment of legs of animals is supplemental to premise treatment for stable fly control. Repeat if necessary, but not more often than once every 4 days.

**CAUTION -- HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate to severe cholinesterase inhibitor. Atropine is an antidote.

E. P. A. Reg. No. 299-121

17019



**5%**

# SEVIN DUST


**READY TO USE ON:**

VEGETABLES - GARDENS - LAWNS

POULTRY - HORSES

DOGS - CATS

NET WEIGHT: 1 POUND



**Keep Out of Reach of Children.**

**SEE BACK PANEL FOR OTHER CAUTIONS.**

© Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate) ..... 5%

INERT INGREDIENTS ..... 95%

**DIRECTIONS**

Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

**VEGETABLES**

**CORN, EGGPLANT, PEPPER, TOMATO:** To control Corn Earworm, European Corn Borer, Sap Beetles, Fall Army Worm, Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm and Stink Bugs -- Dust upper and lower leaf surfaces thoroughly. When used on Sweet Corn for control of Corn Earworm, application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

**BEANS:** To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer -- Dust upper and lower leaf surfaces thoroughly.

**CUCUMBER, SUMMER SQUASH:** To control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug, and Flea Beetles -- Dust upper and lower leaf surfaces thoroughly.

**POTATO:** To control Colorado Potato Beetle, Leafhoppers, Flea Beetles, and Lygus Bugs -- Dust upper and lower leaf surfaces thoroughly.

**APPLICATIONS TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST.**

**LAWNS:** To control Chinch Bugs, Leafhoppers, Fall Army Worms, and Sod Webworms -- apply at the rate of 1 pound to 200 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.