

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow drift to blooming crops or weeds if bees are visiting the treatment area.

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dust has settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

STORAGE & DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry place until used. Store out of reach of children, pets and animals.

DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

581010



5% SEVIN DUST

READY TO USE ON:

**VEGETABLES • LAWNS • GARDENS
POULTRY • HORSES • DOGS • CATS**



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel For Other Cautions

NET WEIGHT: 50 POUNDS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%

©Registered Trademark of Union Carbide for 1-naphthyl N-methylcarbamate

**DIRECTIONS FOR USE
(Continued)**

VEGETABLES:

Corn, Eggplant, Pepper, Tomato: To control Corn Earworm-European Corn Borer, Sap Beetles, Fall Army Worm, Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm and Stink Bugs - Dust the upper and lower leaf surfaces thoroughly. When used on Sweet Corn for control of Corn Earworm, application of this product to the tassel region of corn during the silks shed period will seriously reduce bee population.

Beans: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Leafhopper, Flea Beetle, Japanese Beetle, Corn Earworm, Lygus Bug and Lima Bean Pod Borer - Dust the upper and lower leaf surfaces thoroughly.

Cucumber, Summer Squash: To control Pickleworm, Melonworm, Cucumber Beetle, Squash Bug, and Flea Beetle - Dust the upper and lower leaf surfaces thoroughly.

Potato: To control Colorado Potato Beetle, Leafhopper, Flea Beetle, and Lygus Bug - Dust the upper and lower leaf surfaces thoroughly.

APPLICATION TO THE ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST. SUBSEQUENT PLANTINGS OF FOOD AND FEED CROPS IN TREATED SOIL MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABELS.

LAWNS: To control Chinch Bugs, Leafhoppers, Fall Army Worms, and Sod Webworms - apply at the rate of 1 pound to 200 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

POULTRY: Litter Treatment: Apply 1 pounds to 40 square feet to control Northern Fowl Mites, Chicken Mites, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

DOGS, CATS: To control Brown Dog Ticks and Fleas - apply liberally to animal, rub thoroughly into skin and apply to feet, between toes, and on legs. Dust in and around sleeping quarters, all areas where fleas and ticks occur. Repeat weekly if needed. Do not treat puppies or kittens under four weeks old. Do not use on pregnant dogs.

HORSES: For the control of Hornflies and suppression of Stable Flies - rub 2 ounces of powder per animal into the hair on back and neck. Stable Fly treatment of legs of animals is supplemental to premise treatment for stable fly control. Repeat if necessary, but not more often than once every 4 days.

E.P.A. Reg. No. 299-121
E.P.A. Est. No. 299-TX-1

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75963-1089

299-121

10/17/86

1/1