

Insecticide, apply carefully when the insects threaten and repeat every few days until the pests are eliminated.

PRECAUTION: Do not apply to foliage within one week after applying any spray containing Oil Emulsion. Do not apply any spray containing Oil Emulsion within 30 days to foliage dusted or sprayed with sulfur. Sulfur may burn foliage if applied when temperature is high (90°-95°). Do not make application at this time.

Common Chigger Control—For the control of chiggers in the yard, dust Dusting Sulfur over the entire lawn, flower gardens, shrubs and any place the chiggers may be hiding. This should be done a day or two before protection is desired and should be repeated at weekly intervals. The contents of this package will cover from 1,500 to 2,000 square feet. Dust on clothing, especially around the legs when visiting chigger-infested areas.

NOTICE—Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.



WETTABLE DUSTING SULFUR

(93% THROUGH 325 MESH)

CAUTION: Keep Out of Reach of Children

SEE BACK PANEL FOR OTHER CAUTIONS

NET WEIGHT 2 LBS.

ACTIVE INGREDIENTS: Sulfur	93%
INERT INGREDIENTS:	7%
<hr/>	
TOTAL	100%

DIRECTIONS: Recommended for the control of the following: Black Spot and Powdery Mildew on Roses; Powdery Mildew on Chrysanthemums, Calendula, Dahlias, Delphiniums, Phlox, Sweet Peas, Beans, Peas, and Peaches.

For Use as A Liquid Spray For one gallon of spray, mix 3 to 4 tablespoonfuls of Dusting Sulfur with an equal amount of water and stir into a thick paste, then add water slowly, stirring constantly until the gallon of water has been thoroughly combined. Apply with ordinary sprayer. Keep well stirred at all times.

For Use as a Dust—Dusting Sulfur is all ready for use as it comes in the package. This package is furnished with punch-out holes on the top for dusting direct from the package or use a hand duster procurable at your local dealer.

When to Apply—Apply early in the morning or cool of the evening when the air is still. Thoroughly cover all stems, leaves and blossoms of both upper and lower surfaces. As a fungicide, frequent applications should be made early in the season. During the summer and fall applications should be made at 10 day intervals. As an

Austin, Texas

E. J. Martin & Sons incorporated

Austin, Texas