

CAUTION

Harmful if swallowed. Avoid breathing of vapor. May be absorbed through skin. Avoid getting in eyes or on skin. Wash thoroughly after use. In case of human eye contact, flush eyes immediately with water and get prompt medical attention.

NOTE TO PHYSICIAN: Ronnel is a cholinesterase inhibitor. Atropine is antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine. Treat symptomatically.

Avoid contamination of food and feedstuffs.

Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is antidote. Consult veterinarian at first sign of adverse reaction.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

E.P.A. Reg. No. 299-114
E.P.A. Est. No. 299-TX-1



KORLAN LIVESTOCK WOUND SPRAY

KILLS SCREW WORMS - BLOWFLY MAGGOTS
AND SPINOSE EAR TICKS

For Use On
CATTLE, HORSES, SHEEP, GOATS, SWINE

JET STREAM
For
FAST - DIRECT - EASY APPLICATION

NET WEIGHT, 1 POUND

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR OTHER CAUTIONS

* Trademark of the Dow Chemical Company

472111

DIRECTIONS

SHAKE WELL BEFORE USING

FOR CATTLE, HORSES, SHEEP,
GOATS, SWINE

**SCREW WORM AND BLOWFLY
INFESTATIONS:** Wet wound thor-

oughly by spraying directly into and about 1 inch around wound. Use particular care to get spray into deep pockets caused by screw worms. Retreat at 2 week intervals as needed.

**PREVENTION OF SCREW WORM
AND BLOWFLY INFESTATIONS:**

Apply directly to all open or fresh wounds, such as from dehorning, castration, shear cuts, etc. Treat navels of newborn animals. Fresh bleeding wounds may require several treatments. Retreat at 2 week intervals as needed.

**TO KILL SPINOSE EAR TICKS
AND REDUCE INFESTATION:**

Treat the inside of the ear of the animal thoroughly, making sure that all surfaces are completely covered. Repeat application as necessary for ear tick control.

FLEECE-WORM INFESTATIONS:

Spray infested area liberally, working toward the center of the infestation. Shearing or removing the wool from the infested area is not necessary. Retreat at 2 week intervals as needed.

ACTIVE INGREDIENT:

Ronnel* (0,0-Dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate) 2.50%

INERT INGREDIENTS: 97.50%

606 W. MAIN ST.

C. J. Martin Company

NACOGDOCHES, TEXAS 75961