

potato, tomato, and eggplant, use 4 tablespoonfuls per gallon of water for cutworm control.

On cucumbers, melons, pumpkin, and squash, use 1 to 2 tablespoonfuls per gallon of water for pickle worm and melonworm, and 2 tablespoons per gallon of water for cucumber beetles, flea beetles, leafhoppers, and squash bugs. Avoid excessive applications. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. SEVIN injures watermelons in Florida.

Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary unless shorter intervals are specified.

CAUTION:

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Does not control spider mites but is compatible with common miticides.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

CAUTION:

MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take other equally effective precautions. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

U.S.D.A. Reg. No. 299-141

17224



50% SEVIN WETTABLE POWDER

FOR USE ON:

LAWNS, VEGETABLE GARDENS,
FRUIT, TREES, SHRUBS, FLOWERS

NET WEIGHT: 1 POUND

**CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS**

® Trademark of Union Carbide Corporation for 1 naphthyl methylcarbamate

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 50% by wt.

INERT INGREDIENTS 50% by wt.

DIRECTIONS FOR USE

Keep spray mixtures well agitated. Apply for full coverage in normal spray schedules or when insects or their damage appear. Refer to your state recommendations.

LAWN INSECT CONTROL: Use one (1) pound of Martin's 50% SEVIN in 75-100 gallons of water for each 2500 square feet of established lawn area for the control of chinch bugs, fall armyworms, ants, mosquitoes, leafhoppers, cutworms, fleas, millipedes, sod webworms (lawn moths), and earwigs.

In Florida: Use 1 1/2 pounds per 75-100 gallons of water for each 2500 square feet for chinch bug control.

Use garden hose sprayers or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.


ORNAMENTAL INSECT CONTROL: Use 2 tablespoonfuls of Martin's 50% SEVIN per gallon of water when insects or their damage appear. Repeat weekly or as needed. SEVIN injures Boston Ivy. Use on herbaceous and annual plants (carnations, gladioli, zinnia, chrysanthemum, etc.) to control blister beetle, June beetle, leaf roller, flea beetle, mealybug, rose aphid, leafhopper, thrips (exposed), and Japanese beetle.

Use on shrubs, trees and woody plants (rose, lilac, arborvitae, Juniper, Hydrangea, azalea, elm, maple, oak, dogwood, birch, pine, etc.) to control apple aphid, gypsy moth, mealybug, rose aphid, bagworm, Japanese beetle, oakleaf miner, mimosa webworm, June beetle, scale insects, orange tortrix, tent caterpillar, boxwood leaf miner, leafhopper, thrips (exposed), elm leaf aphid, leaf roller, and elm leaf beetle.

FRUIT INSECT CONTROL: Peaches, apricots, and nectarines. Use 2 tablespoonfuls per gallon of water for codling moth, cucumber beetles, European earwig, Japanese beetle, Pandemis moth, June beetles, lesser peach free borer, orange tortrix, oriental fruit moth, peach twig borer, periodical cicada, plum curculio, fruit tree leaf roller, red-banded leaf roller, tarnished plant bug, tussock moth, Lecanium scale, San Jose scale and olive scale. For optimum scale control, apply when crawlers are present. For lesser peach tree borer control, spray limbs and trunk thoroughly, weekly during moth flight.

Do not apply within 1 day of harvest of peaches. Do not apply within 3 days of harvest of apricots and nectarines.

VEGETABLE INSECT CONTROL: Potato, tomato, eggplant and pepper. Use 1 to 2 tablespoonfuls per gallon of water for Colorado potato beetle, flea beetles, and leafhoppers. Use 2 to 4 tablespoonfuls per gallon of water for European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bugs, and stink bugs. On



potato, tomato, and eggplant, use 4 tablespoonfuls per gallon of water for cutworm control.

On cucumbers, melons, pumpkin, and squash, use 1 to 2 tablespoonfuls per gallon of water for pickleworm and melonworm, and 2 tablespoons per gallon of water for cucumber beetles, flea beetles, leafhoppers, and squash bugs. Avoid excessive applications. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. SEVIN injures watermelons in Florida.

Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary unless shorter interval is specified.

CAUTION:

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Does not control spider mites but is compatible with common miticides.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

CAUTION:

MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take other equally effective precautions. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

U.S.D.A. Reg. No. 299-141

17224



50% SEVIN WETTABLE POWDER

FOR USE ON:

LAWNS, VEGETABLE GARDENS,
FRUIT, TREES, SHRUBS, FLOWERS

NET WEIGHT: 1 POUND

**CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS**

® Trademark of Union Carbide Corporation for 1 naphthyl methylcarbamate

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 50% by wt.

INERT INGREDIENTS 50% by wt.

DIRECTIONS FOR USE

Keep spray mixtures well agitated. Apply for full coverage in normal spray schedules or when insects or their damage appear. Refer to your state recommendations.

LAWN INSECT CONTROL: Use one (1) pound of Martin's 50% SEVIN in 75-100 gallons of water for each 2500 square feet of established lawn area for the control of chinch bugs, fall armyworms, ants, mosquitoes, leafhoppers, cutworms, fleas, millipedes, sod webworms (lawn moths), and earwigs.

In Florida: Use 1 1/4 pounds per 75-100 gallons of water for each 2500 square feet for chinch bug control.

Use garden hose sprayers or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

ORNAMENTAL INSECT CONTROL: Use 2 tablespoonfuls of Martin's 50% SEVIN per gallon of water when insects or their damage appear. Repeat weekly or as needed. SEVIN injures Boston Ivy. Use on herbaceous and annual plants (carnations, gladioli, zinnia, chrysanthemum, etc.) to control blister beetle, June beetle, leaf roller, flea beetle, mealybug, rose aphid, leafhopper, thrips (exposed), and Japanese beetle.

Use on shrubs, trees and woody plants (rose, lilac, arborvitae, Juniper, Hydrangea, azalea, elm, maple, oak, dogwood, birch, pine, etc.) to control apple aphid, gypsy moth, mealybug, rose aphid, bagworm, Japanese beetle, oakleaf miner, mimosa webworm, June beetle, scale insects, orange tortrix, tent caterpillar, boxwood leaf miner, leafhopper, thrips (exposed), elm leaf aphid, leaf roller, and elm leaf beetle.

FRUIT INSECT CONTROL: Peaches, apricots, and nectarines: Use 2 tablespoonfuls per gallon of water for codling moth, cucumber beetles, European earwig, Japanese beetle, Pandemis moth, June beetles, Lesser peach tree borer, orange tortrix, oriental fruit moth, peach twig borer, periodical cicada, plum curculio, fruit tree leaf roller, red-banded leaf roller, tarnished plant bug, tussock moth, Lecanium scale, San Jose scale and olive scale. For optimum scale control, apply when crawlers are present. For lesser peach tree borer control, spray limbs and trunk thoroughly, weekly during moth flight.

Do not apply within 1 day of harvest of peaches. Do not apply within 3 days of harvest of apricots and nectarines.

VEGETABLE INSECT CONTROL Potato, tomato, eggplant and pepper. Use 1 to 2 tablespoonfuls per gallon of water for Colorado potato beetle, flea beetles, and leafhoppers. Use 2 to 4 tablespoonfuls per gallon of water for European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bugs, and stink bugs.

ACCEPTED
5-7-71
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO.