



# GARDEN DUST

Contains SEVIN<sup>®</sup>

For Control Of

MANY DESTRUCTIVE INSECT PESTS

IN GARDENS ACCEPTED

NET WEIGHT: 1 POUND

**CAUTION: Keep Out of Reach of Children**

**SEE BACK PANEL FOR OTHER CAUTIONS**

<sup>®</sup> Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate) ..... 5%

INERT INGREDIENTS ..... 95%

**LAWNS** -- To control Leafhoppers, Fall Army Worms, and Sod Webworms -- apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

**CAUTION -- HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and food-stuffs.

**CAUTION -- THIS PRODUCT IS TOXIC TO BEES.** Any use of the product when honey bees are visiting the crop will seriously reduce bee population. Bees are important pollinators and are essential to the production of many crops. Before using this product during such period consult your State Agricultural Extension Service.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. Reg. No. 299-143

17417

**DIRECTIONS**

Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control.

**CAUTION:** For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

**VEGETABLES -- Corn, Eggplant, Pepper, Tomato:** To control Corn Earworm, European Corn Borer, Sap Beetles, Fall Army Worm, Tomato Fruitworm, Flea Beetles, Leafhopper, Tomato Hornworm and Stink Bugs -- Dust upper and lower leaf surfaces thoroughly. When used on Sweet Corn for control of Corn Earworm, application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

**BEANS --** To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer -- Dust upper and lower leaf surfaces thoroughly.

**CUCUMBER, SUMMER SQUASH --** To control Pickleworm, Melonworm, Cucumber Beetles, Leafhoppers, Flea Beetles, and Lygus Bugs -- Dust upper and lower leaf surfaces thoroughly. Applications to above vegetables can be made up to day of harvest.