


DUST INDIVIDUAL FOWLS THIS WAY



Individual dusting is necessary for best results.

TO MAKE A DUST BATH:

Add 1 pound of Martin's Lice Killer to 1 bushel of dust, place this in a wooden box or some other similar container and locate same in a place where fowls can reach it readily. Do not apply within 7 days of slaughter.



CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. Reg. No. 299-144

0758



Martin's

LICE KILLER

Contains SEVIN®

For Use On:

- POULTRY
- BEEF CATTLE
- HORSES
- SWINE
- SHEEP
- DOGS
- CATS

To Control:

- POULTRY LICE
- and
- CERTAIN OTHER INSECTS

CONTENTS: 1 POUND

CAUTION: Keep Out of Reach of Children
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl methylcarbamate)	5%
INERT INGREDIENTS	95%

® Trademark of Union Carbide Corporation for 1-Naphthyl methylcarbamate.

DIRECTIONS

POULTRY:

For poultry lice, chicken mites, and fleas: Shake powder thoroughly into feathers, especially under wings, breast and back of head. Use 1 pound of dust for each 100 birds. Application for chicken mites and fleas is supplemental to litter treatment for these pests. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water.

Litter treatment for lice, chicken mites, fleas, northern fowl mites and bedbugs: Apply 1 pound of dust to 40 square feet of floor area. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

BEEF CATTLE, HORSES, SHEEP, SWINE: For hornflies and suppression of stable flies: Rub 2 ounces of powder per animal into the hair on back and neck. Repeat as necessary but not more often than once every 4 days. Do not apply within one week of slaughter and do not exceed dosages given above. Do not use on dairy animals.

DOGS, CATS:

For brown dog ticks, fleas: Apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

606 W. MAIN ST

C. J. Martin Company

NACOGDOCHES, TEX. 75961

299-144

1-12-1971

42

MS-PA

12117

ACCEPTED

JAN 12 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 299-144 SUBJECT
TO ATTACHED COMMENTS.

2/2