

299-144

7-22-1971

1/3

M
MARTIN'S
ESTABLISHED 1883

LICE KILLER
Contains SEVIN®

For Use On:

POULTRY

HORSES

DOGS

CATS

To Control:

POULTRY LICE

and

CERTAIN OTHER

INSECTS

ACCEPTED
JUL 22 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 299-144 SUBJECT
TO ATTACHED COMMENTS.

CAUTION: Keep Out of Reach of Children;
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl methylcarbamate) 5%
INERT INGREDIENTS 95%

NET WEIGHT: 1 POUND

® Trademark of Union Carbide Corporation for 1-Naphthyl methylcarbamate.

606 W. Main St.

C. J. MARTIN COMPANY

Nacogdoches, Texas 75961

606 W. Main St.

DIRECTIONS

POULTRY:

FOR POULTRY LICE, CHICKEN MITES, AND FLEAS: Shake powder thoroughly into feathers, especially under wings, breast and back of head. Use 1 pound of dust for each 100 birds. Application for chicken mites and fleas is supplemental to litter treatment for these pests. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water.

LITTER TREATMENT FOR LICE, CHICKEN MITES, FLEAS, NORTHERN FOWL MITES AND BEDBUGS: Apply 1 pound of dust to 40 square feet of floor area. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.


DUST INDIVIDUAL FOWLS THIS WAY

Individual dusting is necessary for best results.



TO MAKE A DUST BATH:

For control of northern fowl mite and lice and as a supplement to litter treatment for chicken mite and fleas: Use 2 1/2 lbs. of LICE KILLER per box for each 50 birds. Do not apply within 7 days of slaughter.



HORSES:

FOR HORNFLEES AND SUPPRESSION OF STABLE FLIES: Rub 2 ounces of powder per animal into the hair on back and neck. Repeat as necessary but not more often than once every 4 days.



DOGS, CATS:

FOR BROWN DOG TICKS, FLEAS: Apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

CAUTION: HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

U.S.D.A. REG. NO. 299-144

