

**HORSES.**  
 For hornflies and suppression of stable flies: Rub 2 ounces of powder per animal into the hair on back and neck. Repeat as necessary but not more often than once every 4 days.

**DOGS, CATS:**  
 For brown dog ticks, fleas: Apply liberally to animal, rub thoroughly into skin, apply to feet, between toes and on legs. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat puppies or kittens less than four weeks old.

**CAUTION: HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

E.P.A. Reg. No. 299-144



4362

# LICE KILLER

Contains SEVIN®

For Use On: **ACCEPTED**  
 POULTRY HORSES DOGS CATS  
 To Control: **NOV 19, 1971** POULTRY LICE  
 CERTAIN OTHER INSECTS  
UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT  
 SUBJECT TO THE PROVISIONS OF THE FEDERAL FOOD, DRUG AND COSMETIC ACT  
 299-144  
 COMMENTS.

NET WEIGHT: 1 POUND

**CAUTION: Keep Out of Reach of Children**  
**SEE BACK PANEL FOR OTHER CAUTIONS.**

**ACTIVE INGREDIENTS:**  
 Carbaryl (1-Naphthyl methylcarbamate) . . . . . 5%  
**INERT INGREDIENTS** . . . . . 95%

® Trademark of Union Carbide Corporation for 1-Naphthyl methylcarbamate.

## DIRECTIONS

**POULTRY:**  
 For poultry lice, chicken mites, and fleas: Shake powder thoroughly into feathers, especially under wings, breast and back of head. Use 1 pound of dust for each 100 birds. Application for chicken mites and fleas is supplemental to litter treatment for these pests. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water.

Litter treatment for lice, chicken mites, fleas, northern fowl mites and bedbugs: Apply 1 pound of dust to 40 square feet of floor area. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

DUST INDIVIDUAL FOWLS THIS WAY



Individual dusting is necessary for best results.

### TO MAKE A DUST BATH:

For control of northern fowl mite and lice and as a supplement to litter treatment for chicken mite and fleas: Use 2½ lbs. of LICE KILLER per box for each 50 birds. Do not apply within 7 days of slaughter.

