DIRECTIONS FOR USE

CHICKENS, TURKEYS, DUCKS. GEESE GAME BIRDS PIGEONS

Direct Spray on Birds: For control of Northern F. . Meter Flatter Red Mite, Lice (Chicken Body Louse Menuconthic strates a Shaft Louse, Menopan gallinea, Wing Louse - Laking in appr Fluff Louse, Goniocotes gailinger and Poultry Flease Maria ..... of Super 85P in 5 gallons of water. Use high on the • cages, on litter or slatted floor. Direct spray of the Population Mites and Poultry Fleas is supplemental to spraying the buildings for control of these pests. Repeat of 4 week to sary. Do not apply within 7 days of sloupter Spray Roosts, Litter, Building:

- 1. For control of Poultry Red Mite, Poultry Fleas, Northern Fux Mite, and Bedbugs: Mix 11 pounds of Super 85P per 25 gallon of water. Apply 1 to 2 gallons per 1000 square feet of wa litter or roost surface.
- 2. For control of Fowl Tick: Increase dosage to 5 pounds per 25 gallons of water. Force spray into cracks. Repeat as needed. Ventilate while spraying. Avoid contamination of nests, eggs, feeding and watering troughs. Do not apply these mixes to birds. Do not house birds in treated premises within 7 days of slaughter. COMPATIBILITY

Martin's Super 85 Sprayable Sevin forms a finely divided uniform suspension alone and when combined with many other pesticides. Incompatible mixtures curdle or precipitate and may injure animals or reduce control. Compatibility can be affected by water in your locality, various surfactants, and other formulation ingredients. Do not apply if physically incompatible. Unstable under highly alkaline conditions. Not effective if used with alkaline materials. CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANI-MALS. HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust or Spray. Do Not Take Internally. Avoid Prolonged or Repeated Contact with Skin.

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible. cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD. FEED. WATER SUPPLIES STREAMS, AND PONDS

U.S.D.A. Reg. No. 299-158

606 W; MAIN ST.

£

17317

NACOGDOCHES, TEXAS 75961

X AL



민포직

ACTIV Carb INER1

## **SUPER 85**P POULTRY SPRAYABLE SEVIN

For control of certain pests of poultry.

VE INGREDIENTS	
baryl (1-Naphthyl Methylcarbamate)	85% by weight
TINGREDIENTS	15% by weight

Super 85 Sprayable Sevin is a microfine, uniformly ground sprayable powder for dispersion in water and application as an insecticidal spray, READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

## NET WEIGHT: 5 POUNDS

## **CAUTION:** Keep Out of Reach of Children SEE BACK PANEL FOR OTHER CAUTIONS

PACKAGED BY C. J. MARTIN COMPANY, NACOGDOCHES, TEXAS 75961

(R) SEVIN is a registered trademark of Union Carbide Corporation for carbaryl. U. S. Pat Nos 2,903,478 3,009,855

