

or three inches outside the infestation. Repeat treatment twice weekly until infestation is controlled.

CAUTION: Harmful if swallowed. Avoid breathing spray mist. May be absorbed through skin. Do not get in eyes or on skin. Wash thoroughly after use.

Do not use on dairy animals. Do not use on newborn (very young) animals or calves under 3 months old. Do not use on dogs, cats, or other household pets. Some strains of screw worms may develop resistance to Lindane. This will be observed by lack of response to treatment.

WARNING: Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperature above 130° F. Do not spray into or near open flame.

E.P.A. REG. NO. 299-165

17428

636 W. MAIN ST.



BOMBANE JET STREAM

Ear Tick Control

Kills Screw Worms and Protects at the Same Time
For Beef Cattle, Horses, Swine, Sheep, and Goats

NET WEIGHT: 11 OUNCES

CAUTION: Keep Out of Reach of Children

SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:

Lindane (Gamma Isomer of Benzene Hexachloride) 3.00%

INERT INGREDIENTS 97.00%

DIRECTIONS

EAR TICK CONTROL:

Apply thoroughly to inner surface and base of ears, completely covering all areas and penetrating hair to skin. Repeat treatment as necessary.

TREATMENT OF SCREW-WORM INFESTATIONS IN SUPERFICIAL CUTS OR WOUNDS:

Spray directly into and one inch around infested wounds. Also spray drainage area below wound. Repeat treatment twice weekly until wound heals.

PROTECTION AGAINST SCREW WORMS:

Spray all wounds (wire cuts, dehorning, castration, docking, etc.) as soon as possible to prevent screwworm infestation. Repeat treatment twice weekly until wound heals.

FLEECE WORMS AND OTHER BLOW FLIES:

Spray directly on infested areas and to a distance of two

C. J. Martin Company

NACOGDOCHES, TEXAS 75467

299-165

8-16-72

1 of 2

ACCEPTED
AUG 16 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

299-165