

[299-170]



POULTRY HOUSE SPRAY

potential insect hiding places. Spray ant trails; treat door sills, window frames, and baseboards. Repeat as necessary. Not suitable for the control of stick-tight fleas which frequently infest poultry houses in the South. Do not apply directly to animals, including poultry.

CAUTION:

Harmful if swallowed. Do not contaminate feedstuffs and water supplies. Do not store, use, pour, or spill near heat or open flame.

Do not reuse empty container. Wash small containers thoroughly with water and detergent and discard in a safe place. Return drums to drum reconditioner, or destroy by perforating, or crushing and burying in a safe place away from water supplies.

E.P.A. Reg. No. 299-170

- Ready to Use
- For Fog or Spray Application

CAUTION: Keep Out of Reach of Children!

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

CONTENTS: 5 Gallons

ACTIVE INGREDIENTS:

Pyrethrins	0.30%
* Technical Piperonyl Butoxide	1.50%
Petroleum Distillate	98.20%

* Equivalent to 1.20% (butylcarbityl) (6-propylpiperonyl) ether and 0.30% related compounds.

ACCEPTED

Sept. 11, 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRY NO. 299-170-0001 TO ATTACHED COMMENTS

DIRECTIONS

FOR USE IN POULTRY HOUSES

FOG APPLICATION:

For flies, mosquitoes, wasps, and gnats: Fill room with mist. For best results, keep room tightly closed for 10 minutes. Ventilate building after fogging. For use in large commercial type thermal fogging machines, this product may be diluted 1 gallon to 6 gallons of diesel fuel.

SPRAY APPLICATION:

For poultry red mites, ticks, certain fleas (See Below), flies, mosquitoes, wasps, gnats, roaches, ants, spiders: Apply undiluted as a coarse spray to point of runoff on interior surfaces of buildings, thoroughly wetting all areas, hitting as many insects as possible. Force spray into cracks, crevices and

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75961