



# DAIRY DUST

**TO CONTROL LICE ON HOGS**  
Treat animals thoroughly. Make liberal application to pens and litter.

**CAUTION:** Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Do not contaminate feed water or foodstuffs. Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes. Apply this product only as specified on this label.

This product contains Malathion, a cholinesterase inhibitor. Do not use this product simultaneously or within a few days of treatment with other drugs, insecticides or chemicals having cholinesterase inhibiting activity.

EPA Reg. No. 299-174

## READY TO USE

Kills  
Hornflies, Lice, Ticks  
on  
Dairy Cattle  
and  
Other Livestock

Contains Malathion and Methoxychlor

NET WEIGHT 1 POUND

## CAUTION: Keep Out of Reach of Children

SEE BACK PANEL FOR OTHER CAUTIONS

### ACTIVE INGREDIENTS

1.1% Malathion  
1.1% Methoxychlor  
97.8% Inert Ingredients

### INERT INGREDIENTS

1.1% Malathion  
1.1% Methoxychlor  
97.8% Inert Ingredients

## DIRECTIONS

**TO CONTROL HORNFLIES ON DAIRY AND BEEF CATTLE:**  
Apply 1-2 ounces (2 rounded table spoonsful) per animal to the back and neck. Rub powder into hair. Do not apply within 5 hours of milking or during milking. Repeat at 3 week intervals if needed.

**TO CONTROL LICE AND TICKS ON NON-LACTATING DAIRY CATTLE AND BEEF CATTLE:**  
Treat entire animal thoroughly. Lightly rub dust into hair. Repeat at 14 day intervals if needed. Do not apply during lactation or within 2 weeks of freshening.

**TO CONTROL LICE AND KEDS (TICKS) ON SHEEP AND GOATS:**  
Treat animal thoroughly. Work dust into wool. Repeat as necessary. Do not apply to lactating goats.