

**TO CONTROL LICE AND TICKS ON NON-LACTATING DAIRY CATTLE AND BEEF CATTLE:**

Treat entire animal thoroughly. Lightly rub dust into hair. Repeat at 14 day intervals if needed. Do not apply during lactation or within 2 weeks of freshening.

**TO CONTROL LICE AND KEDS (TICKS) ON SHEEP AND GOATS:**

Treat animal thoroughly. Work dust into wool. Repeat as necessary. Do not apply to lactating goats.

**TO CONTROL LICE ON HOGS:**

Treat animals thoroughly. Make liberal application to pens and litter.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using.

**ENVIRONMENTAL HAZARDS**

Do not contaminate feed, water or foodstuffs.

**PHYSICAL OR CHEMICAL HAZARDS**

This product contains Malathion, cholinesterase inhibitor. Do not use this product simultaneously or within a few days of treatment with other drugs, insecticides, or chemicals having cholinesterase inhibiting activity.


E.P.A. Reg. No. 299-174

E.P.A. Est. 299-TX-1



# DAIRY DUST

READY TO USE



Kills Hornflies, Lice, Ticks on Dairy Cattle and Other Livestock  
Contains Malathion and Methoxychlor

NET WEIGHT: 1 POUND

**CAUTION: Keep Out of Reach of Children**  
**SEE BACK PANEL FOR OTHER CAUTIONS.**

**ACTIVE INGREDIENTS:**

\* Technical Methoxychlor ..... 5.00%  
\*\* Malathion ..... 4.00%

**INERT INGREDIENTS:** ..... 91.00%

\* Equivalent to 2.4% 2,4-dichlorophenyl-1,1,1-trichloroethane and 0.6% other isomers and related compounds.

\*\* 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE & DISPOSAL**

**PROHIBITIONS**

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. Open dumping is prohibited.

**PESTICIDE DISPOSAL**

Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL**


Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

**GENERAL**

Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

**TO CONTROL HORNFLIES ON DAIRY AND BEEF CATTLE:**

Apply 1-2 ounces (2 rounded table-spoonsful) per animal to the back and neck. Rub powder into hair. Repeat at 3 week intervals if needed.



606 W. Main St.

*C. J. Martin Company*

Nacogdoches, Texas 75961