

Spray or wash animals thoroughly. Do not apply within 28 days of slaughter. Do not use on animals under 3 months old. For Mange control, repeat treatment at 7-10 day intervals for two or more applications, as needed. In cases of light infestation one application may be adequate. Repeat as necessary for lice.

Replenishment of Dipping Vats: Add Stock Tox 6112 according to above directions for all fresh water added to vat or replenish by chemical tests.

WARNING

MAY BE FATAL IF SWALLOWED. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using.

Toxic to fish and wildlife. Do not contaminate streams, lakes, or ponds with this material. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty container. Wash glass containers thoroughly with water and detergent. Discard in a safe place. Return drums to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

ANTIDOTE: Give one tablespoonful of salt in a glass of warm water to induce vomiting. Repeat until vomit fluid is clear. Call a physician immediately.

Avoid spraying and dipping animals during cold, stormy weather. Avoid treating sick or overheated animals. Feed and water animals before treating. Spraying is recommended for young animals. If young animals must be dipped, individual hand immersion is essential to avoid swallowing of dip solution.

Do not use on dogs, cats, or other household pets or on poultry, or in poultry houses. Use only as directed on this label. Do not contaminate food or feedstuffs. **DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.**

Stock Tox forms a stable emulsion with water, but some settling may occur, especially in hard water. Thoroughly agitate mixture before dipping or spraying.

NOT FOR HOME USE.

E.P.A. Reg. No. 299-175

27528

606 W. Main St.

C. J. Martin Company

Nac



STOCK TOX 6112

TOXAPHENE EMULSIFIABLE CONCENTRATE

FOR USE AS A LIVESTOCK SPRAY OR DIP
TO CONTROL

SCABIES (PSOROPTIC, CHORIOPTIC AND SARCOPTIC
MANGE), HORNFLIES, LICE, TICKS

CONTENTS: 1 GALLON

WARNING: Keep Out of Reach of Children.


SEE BACK PANEL FOR OTHER WARNINGS

ACTIVE INGREDIENTS:

- ** Toxaphene 67.0%
- Aromatic Petroleum Distillate 26.8%

INERT INGREDIENTS:

- * Animal and Plant Health Service, U.S.D.A.
- ** Technical Chlorinated Camphene (Chlorine content 67% to 69%) 12.2%





DOCK TOX 6112

OXAPHENE EMULSIFIABLE CONCENTRATE

FOR USE AS A LIVESTOCK SPRAY OR DIP
TO CONTROL
SCABIES (PSOROPTIC, CHORIOPTIC AND SARCOPTIC
MANGE), HORNFLIES, LICE, TICKS


CONTENTS: 1 GALLON

WARNING: Keep Out of Reach of Children.

SEE BACK PANEL FOR OTHER WARNINGS

INGREDIENTS:

- Oxaphene 67.0%
 - Aromatic Petroleum Distillate 26.8%
 - Water 12.2%
- INGREDIENTS:
- and Plant Health Service, U.S.D.A.
- Technical Chlorinated Camphene (Chlorine content 67% to 69%)



DIRECTIONS

BEEF CATTLE AND HORSES

To control Scabies (Psoroptic, Sarcoptic and Chorioptic Mange), Lone Star Ticks, Cattle Ticks, Winter Ticks, Gulf Coast Ticks, Hornflies, Cattle Lice, Ear Ticks. Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for control of other insects. Do not apply within 28 days of slaughter. Do not use on dairy cattle. Do not use on animals under 3 months old. Do not dip animals under 6 months old. For calves between 3 and 6 months old must be treated, apply only as a spray.

BACK RUBBERS

As an aid in the control of Hornflies and Lice on beef cattle: Mix 1 quart with 5 gallons of kerosene or diesel fuel. Fill tank of backrubber device or soak burlap roll with solution so it is thoroughly wet but not to point of dripping. Recharge as necessary. Do not permit animals access to treatment within 28 days of slaughter. Do not permit access to backrubbers by dairy animals or calves under 6 months old.

SHEEP AND GOATS

To control Psoroptic Sheep Scab, Fleeceworms, Hornflies, Lone Star Ticks, Ear Ticks: Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use a mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for other insects.

To control Lice and Sheep Ticks (Keds): As a dip, use 1 gallon with 800 gallons of water. For spraying, 1 gallon with 200 gallons of water will give best results. Thoroughly wet animals. Repeat as necessary.

Do not apply within 28 days of slaughter. Do not use on dairy goats. Do not use on animals under 3 months old.

HOGS

To control Lice, Sarcoptic Mange: Mix 1 gallon with 150 gallons of water.

C. J. Martin Company

Nacogdoches, Texas 75961